



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 10, 2023 – 06:31 pm GMT

PDB ID : 1OFR
Title : CRYSTAL STRUCTURE OF THE TYROSINE-REGULATED 3-DEOXY-D-ARABINO-HEPTULOSONATE-7-PHOSPHATE SYNTHASE FROM SACCHAROMYCES CEREVISIAE COMPLEXED WITH PHENYLALANINE AND MANGANESE
Authors : Koenig, V.; Pfeil, A.; Heinrich, G.; Braus, G.; Schneider, T.R.
Deposited on : 2003-04-18
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	370	
1	B	370	
1	C	370	
1	D	370	
1	E	370	
1	F	370	
1	G	370	
1	H	370	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 20930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	350	Total 2631	C 1636	N 474	O 511	S 10	0	0	1
1	B	348	Total 2618	C 1629	N 469	O 510	S 10	0	0	1
1	C	349	Total 2626	C 1634	N 471	O 511	S 10	0	0	1
1	D	351	Total 2623	C 1630	N 470	O 513	S 10	0	0	1
1	E	339	Total 2560	C 1591	N 460	O 499	S 10	0	0	1
1	F	347	Total 2604	C 1620	N 467	O 507	S 10	0	0	1
1	G	344	Total 2581	C 1605	N 462	O 504	S 10	0	0	1
1	H	347	Total 2599	C 1614	N 466	O 509	S 10	0	0	1

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mn 1	0	0
2	B	1	Total 1	Mn 1	0	0
2	C	1	Total 1	Mn 1	0	0
2	D	1	Total 1	Mn 1	0	0
2	E	1	Total 1	Mn 1	0	0
2	F	1	Total 1	Mn 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Mn 1 1	0	0
2	H	1	Total Mn 1 1	0	0

- Molecule 3 is PHENYLALANINE (three-letter code: PHE) (formula: C₉H₁₁NO₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C N O 12 9 1 2	0	0
3	D	1	Total C N O 12 9 1 2	0	0
3	G	1	Total C N O 12 9 1 2	0	0
3	H	1	Total C N O 12 9 1 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	6	Total O 6 6	0	0
4	B	4	Total O 4 4	0	0
4	C	5	Total O 5 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	5	Total O 5 5	0	0
4	E	2	Total O 2 2	0	0
4	F	4	Total O 4 4	0	0
4	G	3	Total O 3 3	0	0
4	H	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain A: 




- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain B: 




- Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE


Chain C: 



• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE



• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE



M277	I276	Y280	S281	H282	S285	D288	F289	R290	N291	Q292	P293	K294	V295	N296	V299	C300	E301	Q302	I303	A304	M305	G306	E307	N308	M314	I315	E316	S317	N318	I319	N320	E321	G322	N323	Q324	G325	I326	P327	A328	GLU	GLY	LYS	A332	G337	V338	S339	D342	A343	C344	I345	G346	W347
E348	T349	I350	E351	D352	V353	L354	R355	K356	L357	V361	R362	Q363	R364	R365	E366	V367	N368	K369	LYS																																	

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	82.29Å 94.03Å 105.12Å 65.14° 85.68° 75.73°	Depositor
Resolution (Å)	37.20 – 2.70	Depositor
% Data completeness (in resolution range)	85.9 (37.20-2.70)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.232 , 0.274	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	20930	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2669	0.63	0/3613
1	B	0.42	0/2654	0.61	0/3590
1	C	0.42	0/2664	0.64	0/3605
1	D	0.43	0/2661	0.64	0/3605
1	E	0.39	0/2596	0.61	0/3513
1	F	0.41	0/2641	0.63	0/3575
1	G	0.40	0/2617	0.61	0/3544
1	H	0.39	0/2637	0.60	0/3577
All	All	0.41	0/21139	0.62	0/28622

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2631	0	2634	126	0
1	B	2618	0	2618	136	0
1	C	2626	0	2631	126	0
1	D	2623	0	2605	119	0
1	E	2560	0	2562	122	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2604	0	2596	105	0
1	G	2581	0	2564	146	0
1	H	2599	0	2576	132	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	C	12	0	8	0	0
3	D	12	0	8	0	0
3	G	12	0	8	1	0
3	H	12	0	8	1	0
4	A	6	0	0	1	0
4	B	4	0	0	0	0
4	C	5	0	0	1	0
4	D	5	0	0	1	0
4	E	2	0	0	0	0
4	F	4	0	0	1	0
4	G	3	0	0	0	0
4	H	3	0	0	0	0
All	All	20930	0	20818	964	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 964 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:ILE:HD12	1:C:134:ILE:H	1.19	1.05
1:B:134:ILE:H	1:B:134:ILE:HD12	1.19	1.05
1:E:324:GLN:HE21	1:E:338:VAL:HB	1.18	1.01
1:G:355:ARG:HH11	1:G:355:ARG:HG3	1.28	0.94
1:C:208:LEU:HD11	1:C:246:VAL:HG11	1.50	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/370 (94%)	298 (86%)	41 (12%)	7 (2%)	7	19
1	B	340/370 (92%)	296 (87%)	35 (10%)	9 (3%)	5	13
1	C	345/370 (93%)	302 (88%)	35 (10%)	8 (2%)	6	16
1	D	347/370 (94%)	299 (86%)	38 (11%)	10 (3%)	4	10
1	E	333/370 (90%)	294 (88%)	35 (10%)	4 (1%)	13	32
1	F	341/370 (92%)	307 (90%)	32 (9%)	2 (1%)	25	50
1	G	336/370 (91%)	287 (85%)	38 (11%)	11 (3%)	4	8
1	H	343/370 (93%)	298 (87%)	34 (10%)	11 (3%)	4	9
All	All	2731/2960 (92%)	2381 (87%)	288 (10%)	62 (2%)	6	16

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	111	GLU
1	B	22	ASP
1	B	286	ASN
1	B	324	GLN
1	C	323	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/298 (94%)	260 (93%)	20 (7%)	14	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	279/298 (94%)	267 (96%)	12 (4%)	29	57
1	C	280/298 (94%)	264 (94%)	16 (6%)	20	44
1	D	277/298 (93%)	266 (96%)	11 (4%)	31	60
1	E	275/298 (92%)	258 (94%)	17 (6%)	18	40
1	F	277/298 (93%)	258 (93%)	19 (7%)	15	35
1	G	274/298 (92%)	255 (93%)	19 (7%)	15	35
1	H	276/298 (93%)	266 (96%)	10 (4%)	35	64
All	All	2218/2384 (93%)	2094 (94%)	124 (6%)	21	45

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	318	ASN
1	G	324	GLN
1	E	300	CYS
1	G	318	ASN
1	H	127	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	286	ASN
1	H	305	ASN
1	F	216	GLN
1	H	243	HIS
1	G	320	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PHE	G	1002	-	11,12,12	0.71	0	14,15,15	0.56	0
3	PHE	D	1002	-	11,12,12	0.76	0	14,15,15	0.57	0
3	PHE	H	1002	-	11,12,12	0.65	0	14,15,15	0.60	0
3	PHE	C	1002	-	11,12,12	0.72	0	14,15,15	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PHE	G	1002	-	-	4/8/8/8	0/1/1/1
3	PHE	D	1002	-	-	2/8/8/8	0/1/1/1
3	PHE	H	1002	-	-	1/8/8/8	0/1/1/1
3	PHE	C	1002	-	-	3/8/8/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	1002	PHE	O-C-CA-N
3	G	1002	PHE	OXT-C-CA-N

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	1002	PHE	OXT-C-CA-N
3	G	1002	PHE	O-C-CA-CB
3	G	1002	PHE	OXT-C-CA-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1002	PHE	1	0
3	H	1002	PHE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.