



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 01:02 PM EST

PDB ID : 4OF1
Title : crystal structure of toxin from staphylococcus aureus Mu50
Authors : Park, J.Y.; Im, H.; Seok, S.H.; Lee, B.J.
Deposited on : 2014-01-14
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 3% 73% 12% •• 10%
1	B	128	 2% 73% 9% 6% • 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mRNA interferase MazF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	115	901	570	158	170	3	0	0	0
1	B	115	895	565	157	170	3	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	LEU	-	expression tag	UNP Q7A2N3
A	122	GLU	-	expression tag	UNP Q7A2N3
A	123	HIS	-	expression tag	UNP Q7A2N3
A	124	HIS	-	expression tag	UNP Q7A2N3
A	125	HIS	-	expression tag	UNP Q7A2N3
A	126	HIS	-	expression tag	UNP Q7A2N3
A	127	HIS	-	expression tag	UNP Q7A2N3
A	128	HIS	-	expression tag	UNP Q7A2N3
B	121	LEU	-	expression tag	UNP Q7A2N3
B	122	GLU	-	expression tag	UNP Q7A2N3
B	123	HIS	-	expression tag	UNP Q7A2N3
B	124	HIS	-	expression tag	UNP Q7A2N3
B	125	HIS	-	expression tag	UNP Q7A2N3
B	126	HIS	-	expression tag	UNP Q7A2N3
B	127	HIS	-	expression tag	UNP Q7A2N3
B	128	HIS	-	expression tag	UNP Q7A2N3



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	48	Total	O	0	0
			48	48		
2	B	29	Total	O	0	0
			29	29		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: mRNA interferase MazF



- Molecule 1: mRNA interferase MazF



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	127.27Å 127.27Å 41.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	110.22 – 2.45 29.55 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (110.22-2.45) 100.0 (29.55-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	23.03 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.164 , 0.225 0.163 , 0.223	Depositor DCC
R_{free} test set	734 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtrriage
Anisotropy	0.059	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1873	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	1/909 (0.1%)	1.09	4/1222 (0.3%)
1	B	0.91	1/903 (0.1%)	1.15	7/1215 (0.6%)
All	All	0.95	2/1812 (0.1%)	1.12	11/2437 (0.5%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	TYR	CG-CD2	5.36	1.46	1.39
1	B	19	GLU	CD-OE1	5.15	1.31	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	80	ARG	NE-CZ-NH2	-17.70	111.45	120.30
1	A	80	ARG	NE-CZ-NH2	-8.68	115.96	120.30
1	B	80	ARG	CG-CD-NE	-8.34	94.28	111.80
1	B	107	MET	CG-SD-CE	-8.21	87.07	100.20
1	B	80	ARG	NE-CZ-NH1	6.97	123.79	120.30
1	A	80	ARG	CB-CA-C	-5.94	98.52	110.40
1	A	106	LEU	CA-CB-CG	5.58	128.13	115.30
1	B	106	LEU	CA-CB-CG	5.41	127.75	115.30
1	A	49	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	24	ARG	NE-CZ-NH2	-5.35	117.62	120.30
1	B	24	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	901	0	969	18	0
1	B	895	0	951	12	0
2	A	48	0	0	2	0
2	B	29	0	0	1	0
All	All	1873	0	1920	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ARG:HD3	2:A:237:HOH:O	1.98	0.63
1:B:12:LEU:O	1:B:80:ARG:NH2	2.31	0.62
1:B:76:LEU:HD12	1:B:106:LEU:HD13	1.81	0.61
1:A:52:LYS:NZ	2:A:239:HOH:O	2.37	0.57
1:B:24:ARG:HH22	1:B:49:ARG:HH12	1.51	0.56
1:A:107:MET:HE2	1:A:112:LEU:CB	2.36	0.55
1:A:107:MET:HE2	1:A:112:LEU:HB2	1.89	0.55
1:A:43:VAL:HG21	1:A:82:LEU:HD22	1.90	0.54
1:A:107:MET:CE	1:A:112:LEU:CB	2.87	0.52
1:A:107:MET:CE	1:A:112:LEU:HB3	2.41	0.51
1:A:80:ARG:HD2	1:A:82:LEU:HD13	1.94	0.50
1:B:34:GLY:O	1:B:38:SER:HB3	2.12	0.48
1:B:80:ARG:HD2	2:B:202:HOH:O	2.13	0.47
1:A:76:LEU:HD12	1:A:106:LEU:HD13	1.97	0.47
1:A:107:MET:HE1	1:B:107:MET:SD	2.55	0.47
1:A:107:MET:CE	1:A:112:LEU:HB2	2.45	0.46
1:B:91:LEU:O	1:B:92:THR:HB	2.16	0.45
1:A:98:LYS:HE3	1:A:98:LYS:HA	1.99	0.44
1:A:113:ASN:C	1:A:115:VAL:N	2.69	0.44
1:B:113:ASN:O	1:B:114:ALA:CB	2.66	0.43
1:B:63:LYS:HE2	1:B:63:LYS:HB2	1.49	0.43
1:A:63:LYS:HE3	1:A:63:LYS:HB2	1.54	0.43
1:A:103:ASP:O	1:A:107:MET:HG3	2.18	0.43
1:B:24:ARG:HH22	1:B:49:ARG:NH1	2.16	0.42
1:A:24:ARG:HB2	1:A:25:PRO:CD	2.50	0.41
1:B:79:ILE:O	1:B:80:ARG:HG2	2.20	0.41
1:A:24:ARG:HB2	1:A:25:PRO:HD2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ALA:O	1:A:115:VAL:C	2.59	0.41
1:B:50:ILE:HG13	1:B:73:VAL:HG21	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	113/128 (88%)	111 (98%)	2 (2%)	0	100	100
1	B	113/128 (88%)	109 (96%)	1 (1%)	3 (3%)	5	2
All	All	226/256 (88%)	220 (97%)	3 (1%)	3 (1%)	12	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	92	THR
1	B	113	ASN
1	B	114	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/114 (90%)	94 (92%)	8 (8%)	12	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	100/114 (88%)	94 (94%)	6 (6%)	19	24
All	All	202/228 (89%)	188 (93%)	14 (7%)	15	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	SER
1	A	49	ARG
1	A	80	ARG
1	A	82	LEU
1	A	85	LYS
1	A	104	ASN
1	A	107	MET
1	A	115	VAL
1	B	20	GLN
1	B	49	ARG
1	B	63	LYS
1	B	70	LYS
1	B	80	ARG
1	B	88	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	115/128 (89%)	-0.39	4 (3%) 44 40	17, 30, 57, 91	0
1	B	115/128 (89%)	-0.14	3 (2%) 56 52	18, 32, 64, 115	0
All	All	230/256 (89%)	-0.26	7 (3%) 50 46	17, 32, 63, 115	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	114	ALA	6.8
1	B	115	VAL	6.1
1	A	115	VAL	3.9
1	A	114	ALA	3.8
1	B	37	TYR	2.3
1	A	113	ASN	2.2
1	A	62	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.