



Full wwPDB X-ray Structure Validation Report ⓘ

May 4, 2026 – 10:16 AM EDT

PDB ID : 9OFS / pdb_00009ofs
Title : Crystal structure of the human IGA2m2 FC fragment-FC-alpha receptor (CD89) complex
Authors : Chandravanshi, M.; Korzeniowski, M.; Tolbert, W.D.; Pazgier, M.
Deposited on : 2025-04-30
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>16%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>• •</div> </div> </div>
2	C	207	<div> <div>13%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>• 6%</div> </div> </div>
3	G	4	<div> <div>25%</div> <div>75%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3443 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunoglobulin heavy constant alpha 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	1	0
			1610	1013	281	308	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	451	LEU	MET	conflict	UNP P01877
A	454	LYS	SER	conflict	UNP P01877

- Molecule 2 is a protein called Immunoglobulin alpha Fc receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	195	Total	C	N	O	S	0	0	0
			1560	1000	266	286	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	196	ALA	-	expression tag	UNP P24071
C	197	ILE	-	expression tag	UNP P24071
C	198	ASP	-	expression tag	UNP P24071
C	199	GLY	-	expression tag	UNP P24071
C	200	ARG	-	expression tag	UNP P24071
C	201	ALA	-	expression tag	UNP P24071
C	202	HIS	-	expression tag	UNP P24071
C	203	HIS	-	expression tag	UNP P24071
C	204	HIS	-	expression tag	UNP P24071
C	205	HIS	-	expression tag	UNP P24071
C	206	HIS	-	expression tag	UNP P24071
C	207	HIS	-	expression tag	UNP P24071


- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyran

ose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Cl 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	109	Total 109	O 109	0	0
6	C	57	Total 57	O 57	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Immunoglobulin heavy constant alpha 2



- Molecule 2: Immunoglobulin alpha Fc receptor



- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	46.64Å 128.96Å 183.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.69 – 1.95 31.69 – 1.95	Depositor EDS
% Data completeness (in resolution range)	93.2 (31.69-1.95) 93.3 (31.69-1.95)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.227 , 0.253 0.228 , 0.255	Depositor DCC
R_{free} test set	1907 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	29.5	Xtriage
Anisotropy	0.858	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3443	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1651	0.49	0/2256
2	C	0.28	0/1604	0.59	1/2177 (0.0%)
All	All	0.28	0/3255	0.54	1/4433 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	PRO	N-CA-C	-9.99	91.89	112.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1610	0	1585	25	0
2	C	1560	0	1513	18	0
3	G	50	0	43	0	0
4	A	14	0	13	0	0
4	C	42	0	39	3	0
5	A	1	0	0	0	0
6	A	109	0	0	0	0
6	C	57	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3443	0	3193	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:5:PHE:HB3	6:C:409:HOH:O	1.68	0.94
2:C:178:ARG:HG3	2:C:178:ARG:HH11	1.35	0.89
2:C:178:ARG:HH11	2:C:178:ARG:CG	2.02	0.72
1:A:437:GLU:HA	2:C:85:HIS:CE1	2.27	0.69
2:C:178:ARG:HG3	2:C:178:ARG:NH1	2.10	0.66
1:A:362:ASN:HD21	1:A:425:LYS:NZ	1.95	0.65
1:A:327:HIS:CE1	1:A:329:GLU:HG3	2.33	0.64
1:A:246:LEU:HD11	1:A:266:CYS:SG	2.43	0.58
1:A:282:THR:OG1	1:A:283:PRO:HD3	2.05	0.55
2:C:44:ASN:H	4:C:301:NAG:H83	1.73	0.53
1:A:272:ARG:O	1:A:273:ASP:HB3	2.09	0.53
1:A:327:HIS:HE1	1:A:329:GLU:HG3	1.72	0.52
1:A:261:GLU:HG2	1:A:261:GLU:O	2.11	0.50
2:C:3:GLY:H	2:C:87:ARG:HH12	1.58	0.50
2:C:32:ARG:HD3	2:C:58:ASN:HA	1.94	0.48
1:A:251:PRO:HG3	1:A:264:LEU:HD12	1.96	0.48
1:A:314:PRO:HB2	1:A:321:PHE:HZ	1.79	0.47
2:C:59:GLU:HG3	4:C:302:NAG:C7	2.44	0.47
1:A:327:HIS:CD2	1:A:328:PRO:HD2	2.49	0.47
1:A:426:LYS:HB3	1:A:426:LYS:HE2	1.76	0.47
2:C:44:ASN:N	4:C:301:NAG:H83	2.30	0.47
2:C:83:ILE:HD13	2:C:89:ARG:HG3	1.97	0.46
1:A:268:LEU:HD22	1:A:325:ALA:HB1	1.97	0.46
1:A:362:ASN:HD21	1:A:425:LYS:HZ3	1.63	0.46
2:C:58:ASN:O	2:C:59:GLU:HG2	2.17	0.45
2:C:153:HIS:O	6:C:401:HOH:O	2.21	0.45
1:A:246:LEU:HD13	1:A:268:LEU:HD13	1.98	0.45
1:A:311:CYS:C	1:A:314:PRO:HD2	2.42	0.44
1:A:354:PRO:CB	1:A:365:VAL:HG13	2.48	0.44
2:C:134:ARG:HD3	2:C:175:TRP:CH2	2.52	0.43
1:A:298:LEU:C	1:A:300:GLY:H	2.26	0.43
1:A:268:LEU:HB3	1:A:271:LEU:HD11	2.00	0.43
2:C:134:ARG:HH21	2:C:149:GLN:HG3	1.83	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:LYS:HA	1:A:331:LYS:HD2	1.44	0.42
1:A:354:PRO:HB3	1:A:365:VAL:HG13	2.00	0.42
1:A:371:ALA:HB3	1:A:413:VAL:HG12	2.00	0.42
2:C:10:ILE:HG23	2:C:94:LEU:HB2	2.01	0.42
2:C:163:ASP:O	2:C:194:VAL:HG11	2.19	0.42
1:A:268:LEU:HD12	1:A:268:LEU:HA	1.87	0.41
1:A:287:LYS:HE3	1:A:287:LYS:HB2	1.72	0.41
2:C:49:GLU:OE2	6:C:402:HOH:O	2.21	0.41
1:A:433:MET:HG2	1:A:445:GLN:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/214 (97%)	187 (90%)	15 (7%)	6 (3%)	3	0
2	C	193/207 (93%)	182 (94%)	10 (5%)	1 (0%)	24	16
All	All	401/421 (95%)	369 (92%)	25 (6%)	7 (2%)	6	2

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	LYS
1	A	301	CYS
1	A	329	GLU
1	A	287	LYS
1	A	330	LEU
2	C	55	LYS
1	A	276	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/181 (99%)	171 (96%)	8 (4%)	24	14
2	C	170/180 (94%)	166 (98%)	4 (2%)	43	36
All	All	349/361 (97%)	337 (97%)	12 (3%)	34	23

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	243	HIS
1	A	272	ARG
1	A	285	SER
1	A	330	LEU
1	A	331	LYS
1	A	343[A]	ASN
1	A	343[B]	ASN
1	A	405	SER
2	C	5	PHE
2	C	52	ARG
2	C	79	CYS
2	C	178	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	362	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	G	1	3,1	14,14,15	0.76	0	17,19,21	1.82	3 (17%)
3	NAG	G	2	3	14,14,15	0.75	0	17,19,21	1.39	3 (17%)
3	MAN	G	3	3	11,11,12	0.76	0	15,15,17	0.88	0
3	MAN	G	4	3	11,11,12	0.66	0	15,15,17	1.49	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	1/6/23/26	0/1/1/1
3	MAN	G	3	3	-	2/2/19/22	0/1/1/1
3	MAN	G	4	3	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	4	MAN	C1-O5-C5	4.72	118.52	112.19
3	G	1	NAG	C2-N2-C7	4.25	128.59	122.90
3	G	1	NAG	C1-C2-N2	-3.73	104.56	110.43
3	G	1	NAG	C1-O5-C5	-3.45	107.57	112.19
3	G	2	NAG	C2-N2-C7	2.79	126.64	122.90
3	G	2	NAG	O5-C1-C2	-2.21	107.87	111.29
3	G	2	NAG	C1-C2-N2	2.10	113.74	110.43

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	3	MAN	C4-C5-C6-O6
3	G	3	MAN	O5-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
3	G	4	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	303	2	14,14,15	0.79	0	17,19,21	1.63	2 (11%)
4	NAG	C	302	2	14,14,15	0.85	1 (7%)	17,19,21	1.57	4 (23%)
4	NAG	A	701	1	14,14,15	0.69	0	17,19,21	1.42	1 (5%)
4	NAG	C	301	2	14,14,15	0.82	0	17,19,21	1.22	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	303	2	-	0/6/23/26	0/1/1/1
4	NAG	C	302	2	-	0/6/23/26	0/1/1/1
4	NAG	A	701	1	-	1/6/23/26	0/1/1/1
4	NAG	C	301	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	302	NAG	C1-C2	2.09	1.55	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	NAG	C1-O5-C5	4.82	118.65	112.19
4	C	303	NAG	O5-C1-C2	-4.37	104.52	111.29
4	C	303	NAG	C1-C2-N2	3.62	116.13	110.43
4	C	302	NAG	C2-N2-C7	3.61	127.74	122.90
4	C	301	NAG	C2-N2-C7	3.20	127.19	122.90
4	C	302	NAG	C4-C3-C2	3.04	115.48	111.02
4	C	302	NAG	O5-C5-C4	-2.09	105.74	110.83
4	C	302	NAG	C1-O5-C5	2.09	114.98	112.19

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	301	NAG	C8-C7-N2-C2
4	C	301	NAG	O7-C7-N2-C2
4	A	701	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	302	NAG	1	0
4	C	301	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/214 (97%)	0.82	35 (16%) 4 4	17, 48, 110, 131	1 (0%)
2	C	195/207 (94%)	0.94	27 (13%) 6 7	25, 47, 101, 115	0
All	All	404/421 (95%)	0.88	62 (15%) 5 6	17, 48, 106, 131	1 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343[A]	ASN	6.8
1	A	330	LEU	5.9
2	C	153	HIS	5.5
2	C	154	PRO	5.4
2	C	112	GLY	4.7
2	C	6	PRO	4.5
1	A	293	PRO	4.5
2	C	83	ILE	4.3
2	C	4	ASP	4.3
1	A	274	ALA	4.3
2	C	196	ALA	4.2
2	C	113	LEU	3.9
2	C	125	CYS	3.9
1	A	271	LEU	3.8
2	C	3	GLY	3.7
1	A	332	THR	3.7
1	A	359	LEU	3.6
1	A	451	LEU	3.5
1	A	282	THR	3.5
1	A	283	PRO	3.5
1	A	299	CYS	3.4
2	C	172	CYS	3.4
2	C	67	ASP	3.3
2	C	2	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	20	LEU	3.2
1	A	331	LYS	3.1
1	A	301	CYS	3.1
2	C	9	PHE	3.0
2	C	72	ASN	3.0
2	C	114	VAL	3.0
1	A	252	ALA	2.8
2	C	5	PHE	2.8
2	C	129	HIS	2.7
2	C	159	LEU	2.7
1	A	300	GLY	2.6
1	A	311	CYS	2.6
2	C	162	VAL	2.6
1	A	321	PHE	2.6
1	A	361	LEU	2.6
1	A	302	TYR	2.5
2	C	193	VAL	2.5
2	C	117	PRO	2.4
1	A	269	THR	2.4
1	A	324	THR	2.4
1	A	275	SER	2.4
1	A	292	GLY	2.3
1	A	327	HIS	2.3
1	A	279	PHE	2.3
1	A	278	THR	2.3
1	A	298	LEU	2.3
2	C	45	SER	2.2
1	A	408	THR	2.2
2	C	121	ILE	2.2
2	C	122	SER	2.2
1	A	277	ALA	2.2
2	C	164	LEU	2.2
1	A	294	PRO	2.2
1	A	328	PRO	2.2
1	A	304	VAL	2.1
1	A	273	ASP	2.1
1	A	334	LEU	2.1
1	A	243	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	G	1	14/15	-	-	68,83,96,98	0
3	NAG	G	2	14/15	-	-	89,112,129,142	0
3	MAN	G	3	11/12	-	-	125,141,145,147	0
3	MAN	G	4	11/12	-	-	132,149,159,160	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	C	301	14/15	0.04	0.18	84,110,125,127	0
4	NAG	C	303	14/15	0.30	0.16	75,98,108,118	0
4	NAG	C	302	14/15	0.46	0.14	67,88,96,105	0
4	NAG	A	701	14/15	0.51	0.15	82,100,108,114	0
5	CL	A	702	1/1	0.98	0.04	21,21,21,21	1

6.5 Other polymers [i](#)

There are no such residues in this entry.