



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 05:15 AM BST

PDB ID : 1OGQ
Title : The crystal structure of PGIP (polygalacturonase inhibiting protein), a leucine rich repeat protein involved in plant defense
Authors : Di Matteo, A.; Federici, L.; Mattei, B.; Salvi, G.; Johnson, K.A.; Savino, C.; De Lorenzo, G.; Tsernoglou, D.; Cervone, F.
Deposited on : 2003-05-08
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYGALACTURONASE INHIBITING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	2392	1524	406	452	10	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	320	Total O 320 320	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: POLYGALACTURONASE INHIBITING PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.84Å 65.45Å 34.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.70 27.88 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.1 (25.00-1.70) 94.1 (27.88-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 1.70Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.190 , 0.240 0.185 , 0.232	Depositor DCC
R_{free} test set	1650 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtrriage
Anisotropy	0.289	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2772	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/2444	1.50	27/3333 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	ARG	NE-CZ-NH2	-21.19	109.70	120.30
1	A	185	ARG	NH1-CZ-NH2	14.39	135.23	119.40
1	A	36	ARG	NE-CZ-NH1	-12.85	113.87	120.30
1	A	185	ARG	NE-CZ-NH1	-10.54	115.03	120.30
1	A	36	ARG	NE-CZ-NH2	10.14	125.37	120.30
1	A	160	ARG	NE-CZ-NH1	-9.60	115.50	120.30
1	A	183	ARG	NE-CZ-NH1	9.59	125.10	120.30
1	A	51	ARG	CD-NE-CZ	8.13	134.98	123.60
1	A	308	PRO	N-CA-CB	7.61	112.43	103.30
1	A	307	SER	CA-C-O	-7.50	104.36	120.10
1	A	107	TYR	CB-CG-CD2	-6.86	116.89	121.00
1	A	185	ARG	CG-CD-NE	-6.75	97.64	111.80
1	A	220	ASP	CB-CG-OD1	6.60	124.24	118.30
1	A	308	PRO	CA-N-CD	-6.55	102.32	111.50
1	A	294	ASP	CB-CG-OD2	6.45	124.11	118.30
1	A	252	ARG	NE-CZ-NH2	6.44	123.52	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	SER	N-CA-CB	-6.37	100.95	110.50
1	A	255	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	A	167	ASP	CB-CG-OD1	6.22	123.89	118.30
1	A	160	ARG	NH1-CZ-NH2	5.95	125.94	119.40
1	A	293	PHE	CB-CG-CD2	-5.54	116.92	120.80
1	A	36	ARG	CD-NE-CZ	5.53	131.34	123.60
1	A	295	VAL	CA-CB-CG1	5.34	118.91	110.90
1	A	156	PHE	CB-CG-CD1	-5.23	117.14	120.80
1	A	106	LEU	CB-CG-CD2	5.21	119.86	111.00
1	A	274	ASN	CB-CG-OD1	5.03	131.66	121.60
1	A	203	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	307	SER	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2413	56	0
2	A	56	0	52	12	0
3	A	4	0	3	0	0
4	A	320	0	0	17	1
All	All	2772	0	2468	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:ASN:HD21	2:A:1317:NAG:C1	1.45	1.29

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASN:HD21	2:A:1315:NAG:C1	1.71	1.02
1:A:274:ASN:ND2	2:A:1317:NAG:C1	2.30	0.94
1:A:189:LYS:HD3	4:A:2184:HOH:O	1.68	0.92
1:A:230:LYS:NZ	4:A:2234:HOH:O	2.05	0.88
1:A:189:LYS:O	1:A:189:LYS:HD2	1.84	0.78
1:A:30:THR:HG23	4:A:2017:HOH:O	1.83	0.77
1:A:233:LEU:H	1:A:254:ASN:HD22	1.32	0.76
1:A:287:GLY:H	1:A:291:GLN:HE22	1.34	0.76
1:A:112:ASN:ND2	2:A:1315:NAG:C1	2.49	0.74
1:A:274:ASN:HD21	2:A:1317:NAG:C2	2.02	0.71
1:A:271:HIS:CD2	4:A:2278:HOH:O	2.44	0.70
1:A:110:HIS:HD2	4:A:2135:HOH:O	1.75	0.70
1:A:159:ASN:HB2	1:A:184:ASN:HD21	1.60	0.66
1:A:231:ASN:HB2	1:A:254:ASN:HD21	1.60	0.65
1:A:174:LYS:CD	4:A:2190:HOH:O	2.45	0.63
1:A:112:ASN:HD21	2:A:1315:NAG:C2	2.14	0.61
1:A:227:HIS:NE2	4:A:2233:HOH:O	2.28	0.60
1:A:2:LEU:HD22	2:A:1314:NAG:H5	1.84	0.59
2:A:1317:NAG:O4	4:A:2315:HOH:O	1.93	0.59
1:A:87:ASN:HD21	2:A:1315:NAG:H4	1.71	0.56
1:A:189:LYS:CE	4:A:2184:HOH:O	2.55	0.55
1:A:233:LEU:H	1:A:254:ASN:ND2	2.02	0.54
1:A:73:ALA:HB1	1:A:99:LYS:HD3	1.88	0.54
1:A:120:PHE:O	1:A:123:GLN:HG2	2.07	0.54
1:A:53:ASN:HD22	1:A:79:ASN:HD22	1.56	0.54
1:A:295:VAL:HG23	1:A:309:LEU:HD23	1.90	0.53
1:A:309:LEU:HB3	1:A:310:PRO:HD2	1.91	0.53
1:A:161:ILE:H	1:A:184:ASN:HD22	1.57	0.53
1:A:111:THR:H	1:A:135:ASN:HD22	1.56	0.52
1:A:95:PRO:HA	1:A:120:PHE:CE2	2.45	0.51
1:A:274:ASN:ND2	2:A:1317:NAG:O5	2.42	0.50
1:A:295:VAL:CG2	1:A:309:LEU:HD23	2.41	0.50
1:A:103:LEU:HD13	1:A:106:LEU:HG	1.93	0.50
1:A:189:LYS:HD2	1:A:189:LYS:C	2.32	0.50
1:A:110:HIS:HE1	4:A:2073:HOH:O	1.96	0.49
1:A:274:ASN:ND2	2:A:1317:NAG:C2	2.69	0.48
1:A:287:GLY:H	1:A:291:GLN:NE2	2.07	0.48
1:A:12:LEU:O	1:A:16:LYS:HG3	2.14	0.47
1:A:195:ALA:HB2	1:A:215:VAL:HG23	1.96	0.47
1:A:112:ASN:ND2	2:A:1315:NAG:O5	2.45	0.47
1:A:16:LYS:HG3	4:A:2018:HOH:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:LYS:CD	4:A:2184:HOH:O	2.42	0.45
1:A:191:PRO:HA	1:A:192:PRO:HD3	1.89	0.44
1:A:309:LEU:HB3	1:A:310:PRO:CD	2.48	0.43
1:A:99:LYS:HG3	4:A:2126:HOH:O	2.18	0.43
1:A:111:THR:H	1:A:135:ASN:ND2	2.17	0.42
1:A:16:LYS:CG	4:A:2018:HOH:O	2.67	0.42
1:A:174:LYS:HD2	4:A:2190:HOH:O	2.12	0.42
1:A:146:SER:HA	1:A:172:PHE:CE2	2.55	0.42
1:A:252:ARG:HG3	1:A:253:ASN:HD22	1.85	0.41
1:A:253:ASN:ND2	4:A:2234:HOH:O	2.43	0.41
1:A:53:ASN:HD22	1:A:79:ASN:ND2	2.17	0.41
1:A:65:PRO:HA	1:A:88:ASN:O	2.22	0.40
1:A:84:GLY:HA2	1:A:109:THR:O	2.21	0.40
1:A:189:LYS:HE3	4:A:2184:HOH:O	2.19	0.40
1:A:68:ILE:CG2	1:A:72:LEU:HD12	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2202:HOH:O	4:A:2298:HOH:O 3_556	1.98	0.22

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	311/313 (99%)	296 (95%)	15 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	274/274 (100%)	264 (96%)	10 (4%)	35 16

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	GLU
1	A	24	LEU
1	A	28	LEU
1	A	33	CYS
1	A	64	LYS
1	A	106	LEU
1	A	121	LEU
1	A	174	LYS
1	A	189	LYS
1	A	303	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	74	ASN
1	A	79	ASN
1	A	87	ASN
1	A	110	HIS
1	A	112	ASN
1	A	135	ASN
1	A	184	ASN
1	A	253	ASN
1	A	254	ASN
1	A	271	HIS
1	A	274	ASN
1	A	291	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1314	-	14,14,15	1.21	1 (7%)	17,19,21	1.69	2 (11%)
2	NAG	A	1317	-	14,14,15	1.42	3 (21%)	17,19,21	3.99	9 (52%)
3	ACT	A	1318	-	1,3,3	4.48	1 (100%)	0,3,3	0.00	-
2	NAG	A	1315	-	14,14,15	1.42	2 (14%)	17,19,21	3.80	7 (41%)
2	NAG	A	1316	-	14,14,15	1.44	2 (14%)	17,19,21	2.53	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1314	-	-	0/6/23/26	0/1/1/1
2	NAG	A	1317	-	-	0/6/23/26	0/1/1/1
2	NAG	A	1315	-	-	0/6/23/26	0/1/1/1
2	NAG	A	1316	-	-	0/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1318	ACT	CH3-C	4.48	1.54	1.48
2	A	1314	NAG	O7-C7	-3.74	1.14	1.23
2	A	1316	NAG	O7-C7	-3.40	1.15	1.23
2	A	1315	NAG	O5-C1	3.24	1.48	1.43
2	A	1316	NAG	O5-C1	3.17	1.48	1.43
2	A	1317	NAG	O5-C1	3.05	1.48	1.43
2	A	1315	NAG	O7-C7	-3.01	1.16	1.23
2	A	1317	NAG	O7-C7	-2.61	1.17	1.23
2	A	1317	NAG	C1-C2	2.13	1.55	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1317	NAG	C1-O5-C5	-12.35	95.46	112.19
2	A	1315	NAG	C1-O5-C5	-12.12	95.77	112.19
2	A	1316	NAG	O5-C1-C2	-6.42	101.15	111.29
2	A	1315	NAG	O5-C1-C2	-5.63	102.40	111.29
2	A	1314	NAG	O5-C1-C2	-5.19	103.09	111.29
2	A	1317	NAG	O5-C1-C2	-5.18	103.10	111.29
2	A	1317	NAG	O5-C5-C6	5.04	115.10	107.20
2	A	1316	NAG	C1-O5-C5	-5.01	105.40	112.19
2	A	1316	NAG	C3-C4-C5	4.80	118.81	110.24
2	A	1315	NAG	O4-C4-C3	-4.45	100.05	110.35
2	A	1315	NAG	C3-C4-C5	4.39	118.07	110.24
2	A	1317	NAG	C4-C3-C2	-3.79	105.46	111.02
2	A	1314	NAG	C4-C3-C2	-3.62	105.72	111.02
2	A	1317	NAG	C2-N2-C7	3.61	128.04	122.90
2	A	1317	NAG	O3-C3-C4	2.88	117.01	110.35
2	A	1317	NAG	O4-C4-C5	-2.81	102.32	109.30
2	A	1317	NAG	C3-C4-C5	2.78	115.19	110.24
2	A	1315	NAG	O6-C6-C5	2.71	120.60	111.29
2	A	1315	NAG	C4-C3-C2	-2.23	107.75	111.02
2	A	1316	NAG	C1-C2-N2	2.23	114.30	110.49
2	A	1317	NAG	O5-C5-C4	-2.23	105.41	110.83
2	A	1315	NAG	O5-C5-C6	2.06	110.43	107.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1314	NAG	1	0
2	A	1317	NAG	6	0
2	A	1315	NAG	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/313 (100%)	0.03	5 (1%) 72 76	13, 19, 28, 38	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	PRO	3.0
1	A	6	GLN	2.9
1	A	174	LYS	2.4
1	A	97	ILE	2.4
1	A	50	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1314	14/15	0.68	0.34	89,89,90,90	0
2	NAG	A	1316	14/15	0.70	0.28	52,54,56,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	1315	14/15	0.75	0.17	27,33,35,36	0
2	NAG	A	1317	14/15	0.92	0.10	18,22,24,25	0
3	ACT	A	1318	4/4	0.97	0.07	21,21,22,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.