

wwPDB X-ray Structure Validation Summary Report (i)

Oct 3, 2023 – 06:22 AM EDT

PDB ID	:	6OHE
Title	:	Alpha-L-fucosidase AlfC D200A in complex with $Fuca(1,6)GlcNAc$
Authors	:	Klontz, E.H.; Sundberg, E.J.
Deposited on		
Resolution	:	3.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)		
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 3.14 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



6OHE

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 10367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	325	Total	С	Ν	0	\mathbf{S}	0	0	0
	А	323	2586	1651	425	498	12	0	0	0
1	В	322	Total	С	Ν	0	S	0	0	0
	D	322	2566	1639	421	494	12	0		
1	С	322	Total	С	Ν	0	S	0	0	0
	U	322	2565	1640	421	492	12	0	0	0
1	Л	320	Total	С	Ν	0	S	0	0	0
		520	2550	1629	419	490	12	0	0	0

• Molecule 1 is a protein called AlfC.

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	200	ALA	ASP	engineered mutation	UNP K0NB39
А	345	LEU	-	expression tag	UNP K0NB39
В	200	ALA	ASP	engineered mutation	UNP K0NB39
В	345	LEU	-	expression tag	UNP K0NB39
С	200	ALA	ASP	engineered mutation	UNP K0NB39
С	345	LEU	-	expression tag	UNP K0NB39
D	200	ALA	ASP	engineered mutation	UNP K0NB39
D	345	LEU	_	expression tag	UNP K0NB39

• Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-bet a-D-glucopyranose.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	Е	2	Total C N O	0	0	0
	Ľ	2	25 14 1 10	0	0	0
9	F	2	Total C N O	0	0	0
	Г	2	25 14 1 10		0	0
2	G	2	Total C N O	0	0	0
	G	2	25 14 1 10	0	0	0
0	Ц	2	Total C N O	0	0	0
	11	2	25 14 1 10	0	0	U



MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	89.02Å 139.25Å 263.38Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.76 - 3.14	Depositor
% Data completeness	99.7 (29.76-3.14)	Depositor
(in resolution range)		-
R _{merge}	0.19	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.39 (at 3.11 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.279 , 0.299	Depositor
Wilson B-factor $(Å^2)$	86.3	Xtriage
Anisotropy	0.479	Xtriage
L-test for twinning ²	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10367	wwPDB-VP
Average B, all atoms $(Å^2)$	126.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Turne	Chain	Res	Link	Bo	ond leng	\mathbf{ths}	В	ond ang	les
	Type	Ullain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	Е	1	2	$15,\!15,\!15$	0.29	0	21,21,21	0.34	0
2	FUC	Е	2	2	$10,\!10,\!11$	0.98	1 (10%)	14,14,16	0.99	0
2	NAG	F	1	2	$15,\!15,\!15$	0.33	0	21,21,21	0.31	0
2	FUC	F	2	2	10,10,11	1.15	1 (10%)	14,14,16	0.88	0
2	NAG	G	1	2	$15,\!15,\!15$	0.13	0	21,21,21	0.38	0
2	FUC	G	2	2	10,10,11	0.85	1 (10%)	14,14,16	1.03	1 (7%)
2	NAG	Н	1	2	$15,\!15,\!15$	0.41	0	21,21,21	0.45	0
2	FUC	Н	2	2	10,10,11	0.53	0	14,14,16	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	Е	1	2	-	2/6/26/26	0/1/1/1
2	FUC	Е	2	2	-	-	0/1/1/1
2	NAG	F	1	2	-	2/6/26/26	0/1/1/1
2	FUC	F	2	2	-	-	0/1/1/1
2	NAG	G	1	2	-	2/6/26/26	0/1/1/1
2	FUC	G	2	2	-	-	0/1/1/1
2	NAG	Н	1	2	-	2/6/26/26	0/1/1/1
2	FUC	Н	2	2	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
2	F	2	FUC	O5-C1	-2.64	1.39	1.43
2	G	2	FUC	O5-C1	-2.48	1.39	1.43
2	Е	2	FUC	O5-C1	-2.36	1.39	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	G	2	FUC	O2-C2-C1	2.61	114.48	109.15

There are no chirality outliers.

5 of 8 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	F	1	NAG	C1-C2-N2-C7
2	G	1	NAG	C1-C2-N2-C7
2	Н	1	NAG	C1-C2-N2-C7
2	Е	1	NAG	C4-C5-C6-O6
2	Е	1	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.
















4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

