



Full wwPDB EM Validation Report (i)

Dec 1, 2021 – 02:28 pm GMT

PDB ID : 7OKN
EMDB ID : EMD-12962
Title : Structure of the outer-membrane core complex (inner ring) from a conjugative type IV secretion system
Authors : Amin, H.; Ilangovan, A.; Costa, T.R.D.
Deposited on : 2021-05-18
Resolution : 3.34 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see [references \(1\)](#)) were used in the production of this report:


EMDB validation analysis : 0.0.0.dev97
MolProbitiy : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	S	461		37%	60%
1	U	461		37%	60%
1	W	461		37%	60%
1	Y	461		37%	60%
1	a	461		37%	60%
1	c	461		37%	60%
1	e	461		37%	60%
1	g	461		37%	60%
2	B	204		18%	81%
2	D	204		18%	81%
2	F	204		18%	81%
2	H	204		18%	81%
2	J	204		18%	81%
2	L	204		18%	81%
2	N	204		18%	81%
2	P	204		18%	81%
2	R	204		18%	81%
2	T	204		18%	81%
2	V	204		18%	81%
2	X	204		18%	81%
2	Z	204		18%	81%
2	b	204		18%	81%
2	d	204		18%	81%
2	f	204		18%	81%
2	h	204		18%	81%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 28067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TraB.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	C	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	E	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	G	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	I	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	K	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	M	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	O	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	Q	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	S	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	U	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	W	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	Y	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	a	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	c	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	e	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		
1	g	183	Total	C	N	O	S	0	0
			1386	871	240	269	6		


- Molecule 2 is a protein called Type IV conjugative transfer system lipoprotein TraV.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	38	265	155	48	57	5	0	0
2	D	38	265	155	48	57	5	0	0
2	F	38	265	155	48	57	5	0	0
2	H	38	265	155	48	57	5	0	0
2	J	38	265	155	48	57	5	0	0
2	L	38	265	155	48	57	5	0	0
2	N	38	265	155	48	57	5	0	0
2	P	38	265	155	48	57	5	0	0
2	R	38	265	155	48	57	5	0	0
2	T	38	265	155	48	57	5	0	0
2	V	38	265	155	48	57	5	0	0
2	X	38	265	155	48	57	5	0	0
2	Z	38	265	155	48	57	5	0	0
2	b	38	265	155	48	57	5	0	0
2	d	38	265	155	48	57	5	0	0
2	f	38	265	155	48	57	5	0	0
2	h	38	265	155	48	57	5	0	0


3 Residue-property plots [\(i\)](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TraB



- Molecule 1: TraB







- Molecule 1: TraB

Chain E: 38% • 60%





A horizontal progress bar for 'Chain E'. The bar is grey with a green segment representing the completed portion. A red dot marks the start of the bar. A black dot is located on the green segment, approximately one-third of the way from the start. To the right of the bar, the text '• 60%' is displayed.



- Molecule 1: TraB


Chain G: 37% • 60%

A horizontal progress bar for 'Chain G'. The bar is divided into three segments: a red segment on the left, a long green segment in the middle, and a grey segment on the right. A black dot is positioned on the green segment, approximately one-third of the way from the start. The text '37%' is centered above the green segment, and '60%' is centered above the grey segment.




- Molecule 1: TraB

Chain I:




- Molecule 1: TraB

Chain K:



- Molecule 1: TraB

Chain M:






- Molecule 1: TraB

Chain O: 38% 60%



A horizontal progress bar consisting of a red dot at the start, followed by a green segment representing 38% completion, and a grey segment representing the remaining 60%.



- Molecule 1: TraB






Chain Q: 37% 60%

A horizontal progress bar for 'Chain Q'. The bar is divided into two segments: a green segment representing 37% completion and a grey segment representing the remaining 60%. The percentage values are displayed in white text within their respective segments.







- Molecule 1: TraB

Chain S:



- Molecule 1: TraB


Chain U




- Molecule 1: TraB

Chain W








- Molecule 1: TraB


Chain c: 37% • 60%

A horizontal progress bar for 'Chain c'. The bar is divided into three segments: a red segment on the left, a green segment in the middle labeled '37%', and a grey segment on the right labeled '60%'. A small black dot is positioned between the green and grey segments.




- Molecule 1: TraB


Chain e: 37% • 60%




- Molecule 1: TraB



- Molecule 2: Type IV conjugative transfer system lipoprotein TraV




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV





- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain H: 18% : 81%




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain J: 18% 81%





- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain L: 18% : 81%




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain N:  18%  81%





- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain P: 18% : 81%




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain R:  18% : 81%



- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain T: 18% : 81%





- Molecule 2: Type IV conjugative transfer system lipoprotein TraV


Chain V: 18% 81%




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV



- Molecule 2: Type IV conjugative transfer system lipoprotein TraV




- Molecule 2: Type IV conjugative transfer system lipoprotein TraV



- Molecule 2: Type IV conjugative transfer system lipoprotein TraV






ARG	THR	VAL	SER	THR	PRO	PRO	VAL	SER	ALA	ALA	GLY	GLY	THR	SER	ALA	GLY	VAL	ASN	ASN	ASN	THR	THR	THR	THR	THR	SER	THR	LEU	THR	PRO	ARG	PRO	VAL	ALA	GLY	THR	PRO	SER	VAL	ALA	TYR	ARG	PRO	VAL	SER	VAL	VAL	VAL	THR	PRO	THR	PRO	SER	CYS	GLN	ASN
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain f:  18% 81%



- Molecule 2: Type IV conjugative transfer system lipoprotein TraV

Chain h: 18% . 81%



4 Experimental information i

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	74956	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1.3	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.778	Depositor
Minimum map value	-1.060	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.041	Depositor
Recommended contour level	0.265	Depositor
Map size (Å)	528.0, 528.0, 528.0	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.36	0/1408	0.55	0/1896
1	C	0.36	0/1408	0.56	0/1896
1	E	0.36	0/1408	0.55	0/1896
1	G	0.36	0/1408	0.55	0/1896
1	I	0.36	0/1408	0.55	0/1896
1	K	0.36	0/1408	0.55	0/1896
1	M	0.36	0/1408	0.55	0/1896
1	O	0.36	0/1408	0.55	0/1896
1	Q	0.36	0/1408	0.55	0/1896
1	S	0.36	0/1408	0.56	0/1896
1	U	0.36	0/1408	0.55	0/1896
1	W	0.36	0/1408	0.55	0/1896
1	Y	0.36	0/1408	0.56	0/1896
1	a	0.36	0/1408	0.56	0/1896
1	c	0.36	0/1408	0.55	0/1896
1	e	0.36	0/1408	0.55	0/1896
1	g	0.36	0/1408	0.56	0/1896
2	B	0.29	0/265	0.61	0/353
2	D	0.30	0/265	0.61	0/353
2	F	0.29	0/265	0.61	0/353
2	H	0.30	0/265	0.61	0/353
2	J	0.29	0/265	0.61	0/353
2	L	0.30	0/265	0.61	0/353
2	N	0.30	0/265	0.61	0/353
2	P	0.30	0/265	0.61	0/353
2	R	0.30	0/265	0.61	0/353
2	T	0.29	0/265	0.61	0/353
2	V	0.30	0/265	0.61	0/353
2	X	0.29	0/265	0.61	0/353
2	Z	0.29	0/265	0.61	0/353
2	b	0.29	0/265	0.61	0/353
2	d	0.29	0/265	0.61	0/353
2	f	0.29	0/265	0.61	0/353
2	h	0.30	0/265	0.61	0/353

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.35	0/28441	0.56	0/38233

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	C	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	E	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	G	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	I	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	K	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	M	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	O	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	Q	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	S	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	U	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	W	179/461 (39%)	163 (91%)	16 (9%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	Y	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	a	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	c	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	e	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
1	g	179/461 (39%)	163 (91%)	16 (9%)	0	100 100
2	B	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	D	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	F	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	H	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	J	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	L	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	N	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	P	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	R	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	T	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	V	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	X	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	Z	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	b	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	d	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	f	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
2	h	36/204 (18%)	32 (89%)	4 (11%)	0	100 100
All	All	3655/11305 (32%)	3315 (91%)	340 (9%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	C	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	E	144/361 (40%)	134 (93%)	10 (7%)	15	47
1	G	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	I	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	K	144/361 (40%)	134 (93%)	10 (7%)	15	47
1	M	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	O	144/361 (40%)	134 (93%)	10 (7%)	15	47
1	Q	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	S	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	U	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	W	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	Y	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	a	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	c	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	e	144/361 (40%)	133 (92%)	11 (8%)	13	42
1	g	144/361 (40%)	133 (92%)	11 (8%)	13	42
2	B	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	D	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	F	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	H	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	J	28/168 (17%)	27 (96%)	1 (4%)	35	65
2	L	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	N	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	P	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	R	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	T	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	V	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	X	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	Z	28/168 (17%)	26 (93%)	2 (7%)	14	45
2	b	28/168 (17%)	27 (96%)	1 (4%)	35	65
2	d	28/168 (17%)	26 (93%)	2 (7%)	14	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	f	28/168 (17%)	27 (96%)	1 (4%)	35 65
2	h	28/168 (17%)	26 (93%)	2 (7%)	14 45
All	All	2924/8993 (32%)	2709 (93%)	215 (7%)	17 43

All (215) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	199	TRP
1	A	224	LYS
1	A	242	ASP
1	A	270	ARG
1	A	271	SER
1	A	276	LEU
1	A	283	GLN
1	A	374	HIS
1	A	383	ASN
1	A	391	ASP
1	A	394	GLN
2	B	32	SER
2	B	48	LYS
1	C	199	TRP
1	C	224	LYS
1	C	242	ASP
1	C	270	ARG
1	C	271	SER
1	C	276	LEU
1	C	283	GLN
1	C	374	HIS
1	C	383	ASN
1	C	391	ASP
1	C	394	GLN
2	D	32	SER
2	D	48	LYS
1	E	199	TRP
1	E	224	LYS
1	E	242	ASP
1	E	270	ARG
1	E	271	SER
1	E	276	LEU
1	E	283	GLN
1	E	383	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	391	ASP
1	E	394	GLN
2	F	32	SER
2	F	48	LYS
1	G	199	TRP
1	G	224	LYS
1	G	242	ASP
1	G	270	ARG
1	G	271	SER
1	G	276	LEU
1	G	283	GLN
1	G	374	HIS
1	G	383	ASN
1	G	391	ASP
1	G	394	GLN
2	H	32	SER
2	H	48	LYS
1	I	199	TRP
1	I	224	LYS
1	I	242	ASP
1	I	270	ARG
1	I	271	SER
1	I	276	LEU
1	I	283	GLN
1	I	374	HIS
1	I	383	ASN
1	I	391	ASP
1	I	394	GLN
2	J	32	SER
1	K	199	TRP
1	K	224	LYS
1	K	242	ASP
1	K	270	ARG
1	K	271	SER
1	K	276	LEU
1	K	283	GLN
1	K	383	ASN
1	K	391	ASP
1	K	394	GLN
2	L	32	SER
2	L	48	LYS
1	M	199	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	M	224	LYS
1	M	242	ASP
1	M	270	ARG
1	M	271	SER
1	M	276	LEU
1	M	283	GLN
1	M	374	HIS
1	M	383	ASN
1	M	391	ASP
1	M	394	GLN
2	N	32	SER
2	N	48	LYS
1	O	199	TRP
1	O	224	LYS
1	O	242	ASP
1	O	270	ARG
1	O	271	SER
1	O	276	LEU
1	O	283	GLN
1	O	383	ASN
1	O	391	ASP
1	O	394	GLN
2	P	32	SER
2	P	48	LYS
1	Q	199	TRP
1	Q	224	LYS
1	Q	242	ASP
1	Q	270	ARG
1	Q	271	SER
1	Q	276	LEU
1	Q	283	GLN
1	Q	374	HIS
1	Q	383	ASN
1	Q	391	ASP
1	Q	394	GLN
2	R	32	SER
2	R	48	LYS
1	S	199	TRP
1	S	224	LYS
1	S	242	ASP
1	S	270	ARG
1	S	271	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	S	276	LEU
1	S	283	GLN
1	S	374	HIS
1	S	383	ASN
1	S	391	ASP
1	S	394	GLN
2	T	32	SER
2	T	48	LYS
1	U	199	TRP
1	U	224	LYS
1	U	242	ASP
1	U	270	ARG
1	U	271	SER
1	U	276	LEU
1	U	283	GLN
1	U	374	HIS
1	U	383	ASN
1	U	391	ASP
1	U	394	GLN
2	V	32	SER
2	V	48	LYS
1	W	199	TRP
1	W	224	LYS
1	W	242	ASP
1	W	270	ARG
1	W	271	SER
1	W	276	LEU
1	W	283	GLN
1	W	374	HIS
1	W	383	ASN
1	W	391	ASP
1	W	394	GLN
2	X	32	SER
2	X	48	LYS
1	Y	199	TRP
1	Y	224	LYS
1	Y	242	ASP
1	Y	270	ARG
1	Y	271	SER
1	Y	276	LEU
1	Y	283	GLN
1	Y	374	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Y	383	ASN
1	Y	391	ASP
1	Y	394	GLN
2	Z	32	SER
2	Z	48	LYS
1	a	199	TRP
1	a	224	LYS
1	a	242	ASP
1	a	270	ARG
1	a	271	SER
1	a	276	LEU
1	a	283	GLN
1	a	374	HIS
1	a	383	ASN
1	a	391	ASP
1	a	394	GLN
2	b	32	SER
1	c	199	TRP
1	c	224	LYS
1	c	242	ASP
1	c	270	ARG
1	c	271	SER
1	c	276	LEU
1	c	283	GLN
1	c	374	HIS
1	c	383	ASN
1	c	391	ASP
1	c	394	GLN
2	d	32	SER
2	d	48	LYS
1	e	199	TRP
1	e	224	LYS
1	e	242	ASP
1	e	270	ARG
1	e	271	SER
1	e	276	LEU
1	e	283	GLN
1	e	374	HIS
1	e	383	ASN
1	e	391	ASP
1	e	394	GLN
2	f	32	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	g	199	TRP
1	g	224	LYS
1	g	242	ASP
1	g	270	ARG
1	g	271	SER
1	g	276	LEU
1	g	283	GLN
1	g	374	HIS
1	g	383	ASN
1	g	391	ASP
1	g	394	GLN
2	h	32	SER
2	h	48	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	374	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-12962. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 283



Y Index: 282



Z Index: 253

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [\(i\)](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.265. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 186 nm³; this corresponds to an approximate mass of 168 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.299 \AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.299 \AA^{-1}

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.34	-	-
Author-provided FSC curve	3.34	3.85	3.41
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit (i)


This section contains information regarding the fit between EMDB map EMD-12962 and PDB model 7OKN. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.265 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Atom inclusion [\(i\)](#)



At the recommended contour level, 86% of all backbone atoms, 75% of all non-hydrogen atoms, are inside the map.