



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 6, 2023 – 10:16 am BST

PDB ID : 8OK8
Title : Variant Surface Glycoprotein VSG615
Authors : Zeelen, J.P.; Stebbins, C.E.; Chandra, M.
Deposited on : 2023-03-27
Resolution : 3.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1335 (3.24-3.20)
Clashscore	141614	1460 (3.24-3.20)
Ramachandran outliers	138981	1437 (3.24-3.20)
Sidechain outliers	138945	1436 (3.24-3.20)
RSRZ outliers	127900	1291 (3.24-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	 5% 54% 21% 24%
1	C	376	 2% 61% 26% 11%
1	E	376	 % 60% 24% 15%
1	G	376	 3% 60% 24% 14%
1	I	376	 2% 59% 23% 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	376	
2	F	2	
2	J	2	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 14117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Variant surface glycoprotein 615.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	Total 2123	C 1302	N 382	O 426	S 13	0	0	0
1	C	335	Total 2457	C 1502	N 438	O 503	S 14	0	0	0
1	E	319	Total 2345	C 1435	N 418	O 478	S 14	0	1	0
1	G	325	Total 2379	C 1452	N 428	O 485	S 14	0	0	0
1	I	314	Total 2311	C 1410	N 417	O 470	S 14	0	0	0
1	K	308	Total 2268	C 1390	N 402	O 462	S 14	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	J	2	Total 28	C 16	N 2	O 10	0	0	0
2	F	2	Total 28	C 16	N 2	O 10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	C	1	Total 14	C 8	N 1	O 5	0	0
3	G	1	Total 14	C 8	N 1	O 5	0	0
3	K	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 4 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 11 6 5	0	0
4	C	1	Total C O 11 6 5	0	0
4	E	1	Total C O 11 6 5	0	0
4	E	1	Total C O 11 6 5	0	0
4	G	1	Total C O 11 6 5	0	0
4	G	1	Total C O 11 6 5	0	0
4	I	1	Total C O 11 6 5	0	0
4	I	1	Total C O 11 6 5	0	0
4	K	1	Total C O 11 6 5	0	0
4	K	1	Total C O 11 6 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	2	Total O 2 2	0	0
5	E	1	Total O 1 1	0	0
5	G	2	Total O 2 2	0	0
5	I	3	Total O 3 3	0	0
5	K	4	Total O 4 4	0	0


• Molecule 1: Variant surface glycoprotein 615




• Molecule 1: Variant surface glycoprotein 615





• Molecule 1: Variant surface glycoprotein 615





• Molecule 1: Variant surface glycoprotein 615



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.28Å 111.23Å 108.59Å 90.00° 91.65° 90.00°	Depositor
Resolution (Å)	77.68 – 3.22 77.68 – 3.22	Depositor EDS
% Data completeness (in resolution range)	92.4 (77.68-3.22) 85.2 (77.68-3.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.307 , 0.325 0.306 , 0.324	Depositor DCC
R_{free} test set	1524 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	85.2	Xtrriage
Anisotropy	0.098	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.046 for -h,-l,-k 0.027 for -h,l,k 0.067 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	14117	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2147	0.51	0/2899
1	C	0.32	0/2487	0.53	0/3362
1	E	0.32	0/2374	0.54	0/3209
1	G	0.37	0/2410	0.54	0/3258
1	I	0.33	0/2338	0.56	0/3157
1	K	0.34	0/2292	0.54	0/3091
All	All	0.32	0/14048	0.54	0/18976

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	G	0	1
1	I	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	274	ARG	Sidechain
1	G	115	ARG	Sidechain
1	I	218	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2123	0	2052	55	0
1	C	2457	0	2362	69	0
1	E	2345	0	2266	65	0
1	G	2379	0	2274	66	0
1	I	2311	0	2241	54	0
1	K	2268	0	2171	73	0
2	F	28	0	25	1	0
2	J	28	0	25	0	0
3	A	14	0	13	1	0
3	C	14	0	13	0	0
3	G	14	0	13	1	0
3	K	14	0	13	1	0
4	C	22	0	20	1	0
4	E	22	0	20	0	0
4	G	22	0	20	3	0
4	I	22	0	20	3	0
4	K	22	0	20	0	0
5	C	2	0	0	0	0
5	E	1	0	0	0	0
5	G	2	0	0	0	0
5	I	3	0	0	0	0
5	K	4	0	0	0	0
All	All	14117	0	13568	369	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 369 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:217:TYR:H	1:I:226:ASN:HB3	1.41	0.84
1:C:303:LEU:HD21	1:C:311:ILE:HG13	1.59	0.83
1:C:268:PHE:HB3	1:C:273:SER:HB2	1.67	0.76
4:I:401:GLC:H3	4:I:402:GLC:H61	1.66	0.76
1:I:269:THR:N	1:I:285:SER:HG	1.84	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/376 (73%)	233 (85%)	41 (15%)	0	100	100
1	C	330/376 (88%)	279 (84%)	49 (15%)	2 (1%)	25	63
1	E	314/376 (84%)	269 (86%)	42 (13%)	3 (1%)	15	52
1	G	319/376 (85%)	268 (84%)	48 (15%)	3 (1%)	17	55
1	I	306/376 (81%)	253 (83%)	49 (16%)	4 (1%)	12	46
1	K	290/376 (77%)	246 (85%)	39 (13%)	5 (2%)	9	40
All	All	1833/2256 (81%)	1548 (84%)	268 (15%)	17 (1%)	17	55

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	202	ALA
1	K	279	LEU
1	K	297	VAL
1	C	268	PHE
1	C	276	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/281 (75%)	202 (96%)	8 (4%)	33	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	242/281 (86%)	231 (96%)	11 (4%)	27	62
1	E	232/281 (83%)	225 (97%)	7 (3%)	41	72
1	G	233/281 (83%)	216 (93%)	17 (7%)	14	46
1	I	229/281 (82%)	213 (93%)	16 (7%)	15	47
1	K	225/281 (80%)	206 (92%)	19 (8%)	11	38
All	All	1371/1686 (81%)	1293 (94%)	78 (6%)	20	55

5 of 78 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	293	SER
1	K	235	ILE
1	K	22	TYR
1	K	189	MET
1	K	281	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	119	HIS
1	I	187	GLN
1	I	301	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	F	1	2,1	14,14,15	0.28	0	17,19,21	0.57	0
2	NAG	F	2	2	14,14,15	0.41	0	17,19,21	0.52	0
2	NAG	J	1	2,1	14,14,15	0.28	0	17,19,21	0.60	0
2	NAG	J	2	2	14,14,15	0.35	0	17,19,21	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	F	2	2	-	1/6/23/26	0/1/1/1
2	NAG	J	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	1	NAG	O5-C5-C6-O6
2	J	2	NAG	C8-C7-N2-C2
2	J	2	NAG	O7-C7-N2-C2
2	J	1	NAG	C4-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	G	402	1	11,11,12	0.83	0	15,15,17	0.64	0
4	GLC	I	402	1	11,11,12	0.82	0	15,15,17	0.64	0
3	NAG	G	401	1	14,14,15	0.45	0	17,19,21	0.56	0
3	NAG	A	401	1	14,14,15	0.37	0	17,19,21	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GLC	E	402	1	11,11,12	0.38	0	15,15,17	0.70	0
3	NAG	C	401	1	14,14,15	0.41	0	17,19,21	0.44	0
4	GLC	K	403	1	11,11,12	0.80	0	15,15,17	0.67	0
4	GLC	C	402	1	11,11,12	0.76	0	15,15,17	0.62	0
4	GLC	E	401	1	11,11,12	0.43	0	15,15,17	0.83	1 (6%)
4	GLC	G	403	1	11,11,12	0.89	0	15,15,17	0.62	0
4	GLC	K	402	1	11,11,12	0.85	0	15,15,17	0.62	0
3	NAG	K	401	1	14,14,15	0.28	0	17,19,21	0.44	0
4	GLC	I	401	1	11,11,12	0.80	0	15,15,17	0.62	0
4	GLC	C	403	1	11,11,12	0.76	0	15,15,17	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	G	402	1	-	0/2/19/22	0/1/1/1
4	GLC	I	402	1	-	1/2/19/22	0/1/1/1
3	NAG	G	401	1	-	0/6/23/26	0/1/1/1
3	NAG	A	401	1	-	0/6/23/26	0/1/1/1
4	GLC	E	402	1	-	1/2/19/22	0/1/1/1
3	NAG	C	401	1	-	1/6/23/26	0/1/1/1
4	GLC	K	403	1	-	0/2/19/22	0/1/1/1
4	GLC	C	402	1	-	0/2/19/22	0/1/1/1
4	GLC	E	401	1	-	2/2/19/22	0/1/1/1
4	GLC	G	403	1	-	0/2/19/22	0/1/1/1
4	GLC	K	402	1	-	0/2/19/22	0/1/1/1
3	NAG	K	401	1	-	0/6/23/26	0/1/1/1
4	GLC	I	401	1	-	1/2/19/22	0/1/1/1
4	GLC	C	403	1	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	401	GLC	O5-C1-C2	-2.26	107.28	110.77

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	E	401	GLC	O5-C5-C6-O6
4	E	401	GLC	C4-C5-C6-O6
3	C	401	NAG	O5-C5-C6-O6
4	I	402	GLC	O5-C5-C6-O6
4	I	401	GLC	O5-C5-C6-O6


There are no ring outliers.



8 monomers are involved in 10 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	402	GLC	1	0
4	I	402	GLC	3	0
3	G	401	NAG	1	0
3	A	401	NAG	1	0
4	C	402	GLC	1	0
4	G	403	GLC	3	0
3	K	401	NAG	1	0
4	I	401	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/376 (76%)	0.35	18 (6%) 20 11	83, 100, 113, 114	0
1	C	335/376 (89%)	0.13	8 (2%) 59 45	62, 77, 116, 118	0
1	E	319/376 (84%)	0.10	4 (1%) 77 66	63, 75, 112, 116	0
1	G	325/376 (86%)	0.13	11 (3%) 45 31	75, 91, 109, 118	0
1	I	314/376 (83%)	0.01	6 (1%) 66 54	64, 74, 105, 111	0
1	K	308/376 (81%)	0.24	9 (2%) 51 37	80, 98, 109, 111	0
All	All	1887/2256 (83%)	0.16	56 (2%) 50 36	62, 88, 111, 118	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	126	PHE	3.8
1	A	69	SER	3.5
1	K	279	LEU	3.2
1	G	201	ALA	3.1
1	A	31	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	F	2	14/15	0.66	0.24	99,99,99,99	0
2	NAG	J	2	14/15	0.69	0.34	97,97,97,97	0
2	NAG	F	1	14/15	0.84	0.25	98,98,98,98	0
2	NAG	J	1	14/15	0.86	0.20	95,95,95,95	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GLC	K	403	11/12	0.51	0.36	108,108,108,108	0
3	NAG	G	401	14/15	0.70	0.32	97,97,97,97	0
4	GLC	G	403	11/12	0.71	0.24	95,95,95,95	0
3	NAG	C	401	14/15	0.77	0.27	94,94,94,94	0
4	GLC	K	402	11/12	0.77	0.22	107,107,107,107	0
4	GLC	C	403	11/12	0.77	0.28	74,74,74,74	0
4	GLC	E	402	11/12	0.83	0.22	80,80,80,80	0
3	NAG	K	401	14/15	0.83	0.26	94,94,94,94	0


Continued on next page...


Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GLC	I	402	11/12	0.84	0.27	77,77,77,77	0
4	GLC	G	402	11/12	0.84	0.16	98,98,98,98	0
4	GLC	I	401	11/12	0.84	0.35	77,77,77,77	0
3	NAG	A	401	14/15	0.85	0.21	88,88,88,88	0
4	GLC	C	402	11/12	0.86	0.21	75,75,75,75	0
4	GLC	E	401	11/12	0.90	0.19	76,76,76,76	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













Electron density around GLC K 402:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around GLC E 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG K 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GLC I 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GLC G 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GLC I 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GLC C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.