



wwPDB X-ray Structure Validation Summary Report

Jan 7, 2024 – 12:18 pm GMT

PDB ID : 5OLC
Title : Crystal structure of the 3,6-anhydro-D-galactonate cycloisomerase from *Zobellia galactanivorans*
Authors : Michel, G.; Czjzek, M.; Jam, M.
Deposited on : 2017-07-27
Resolution : 2.79 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 9% 70% 17% 11%
1	B	396	 7% 67% 19% 11%
1	C	396	 6% 67% 19% 11%
1	D	396	 3% 71% 16% 11%
1	E	396	 3% 68% 18% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	396	 4% 68% 19% • 11%
1	G	396	 4% 68% 17% • 11%
1	H	396	 4% 69% 17% • 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 21906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactonate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	351	2721	1749	446	513	13	0	0	0
1	B	351	2709	1742	444	510	13	0	0	0
1	C	351	2717	1748	446	510	13	0	0	0
1	D	351	2711	1745	443	510	13	0	0	0
1	E	351	2735	1758	451	513	13	0	0	0
1	F	351	2741	1764	451	513	13	0	0	0
1	G	351	2730	1755	449	513	13	0	0	0
1	H	351	2745	1767	452	513	13	0	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP G0L7B8
A	-6	HIS	-	expression tag	UNP G0L7B8
A	-5	HIS	-	expression tag	UNP G0L7B8
A	-4	HIS	-	expression tag	UNP G0L7B8
A	-3	HIS	-	expression tag	UNP G0L7B8
A	-2	HIS	-	expression tag	UNP G0L7B8
A	-1	HIS	-	expression tag	UNP G0L7B8
A	0	GLY	-	expression tag	UNP G0L7B8
A	1	SER	-	expression tag	UNP G0L7B8
B	-7	MET	-	initiating methionine	UNP G0L7B8
B	-6	HIS	-	expression tag	UNP G0L7B8
B	-5	HIS	-	expression tag	UNP G0L7B8
B	-4	HIS	-	expression tag	UNP G0L7B8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	HIS	-	expression tag	UNP G0L7B8
B	-2	HIS	-	expression tag	UNP G0L7B8
B	-1	HIS	-	expression tag	UNP G0L7B8
B	0	GLY	-	expression tag	UNP G0L7B8
B	1	SER	-	expression tag	UNP G0L7B8
C	-7	MET	-	initiating methionine	UNP G0L7B8
C	-6	HIS	-	expression tag	UNP G0L7B8
C	-5	HIS	-	expression tag	UNP G0L7B8
C	-4	HIS	-	expression tag	UNP G0L7B8
C	-3	HIS	-	expression tag	UNP G0L7B8
C	-2	HIS	-	expression tag	UNP G0L7B8
C	-1	HIS	-	expression tag	UNP G0L7B8
C	0	GLY	-	expression tag	UNP G0L7B8
C	1	SER	-	expression tag	UNP G0L7B8
D	-7	MET	-	initiating methionine	UNP G0L7B8
D	-6	HIS	-	expression tag	UNP G0L7B8
D	-5	HIS	-	expression tag	UNP G0L7B8
D	-4	HIS	-	expression tag	UNP G0L7B8
D	-3	HIS	-	expression tag	UNP G0L7B8
D	-2	HIS	-	expression tag	UNP G0L7B8
D	-1	HIS	-	expression tag	UNP G0L7B8
D	0	GLY	-	expression tag	UNP G0L7B8
D	1	SER	-	expression tag	UNP G0L7B8
E	-7	MET	-	initiating methionine	UNP G0L7B8
E	-6	HIS	-	expression tag	UNP G0L7B8
E	-5	HIS	-	expression tag	UNP G0L7B8
E	-4	HIS	-	expression tag	UNP G0L7B8
E	-3	HIS	-	expression tag	UNP G0L7B8
E	-2	HIS	-	expression tag	UNP G0L7B8
E	-1	HIS	-	expression tag	UNP G0L7B8
E	0	GLY	-	expression tag	UNP G0L7B8
E	1	SER	-	expression tag	UNP G0L7B8
F	-7	MET	-	initiating methionine	UNP G0L7B8
F	-6	HIS	-	expression tag	UNP G0L7B8
F	-5	HIS	-	expression tag	UNP G0L7B8
F	-4	HIS	-	expression tag	UNP G0L7B8
F	-3	HIS	-	expression tag	UNP G0L7B8
F	-2	HIS	-	expression tag	UNP G0L7B8
F	-1	HIS	-	expression tag	UNP G0L7B8
F	0	GLY	-	expression tag	UNP G0L7B8
F	1	SER	-	expression tag	UNP G0L7B8
G	-7	MET	-	initiating methionine	UNP G0L7B8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-6	HIS	-	expression tag	UNP G0L7B8
G	-5	HIS	-	expression tag	UNP G0L7B8
G	-4	HIS	-	expression tag	UNP G0L7B8
G	-3	HIS	-	expression tag	UNP G0L7B8
G	-2	HIS	-	expression tag	UNP G0L7B8
G	-1	HIS	-	expression tag	UNP G0L7B8
G	0	GLY	-	expression tag	UNP G0L7B8
G	1	SER	-	expression tag	UNP G0L7B8
H	-7	MET	-	initiating methionine	UNP G0L7B8
H	-6	HIS	-	expression tag	UNP G0L7B8
H	-5	HIS	-	expression tag	UNP G0L7B8
H	-4	HIS	-	expression tag	UNP G0L7B8
H	-3	HIS	-	expression tag	UNP G0L7B8
H	-2	HIS	-	expression tag	UNP G0L7B8
H	-1	HIS	-	expression tag	UNP G0L7B8
H	0	GLY	-	expression tag	UNP G0L7B8
H	1	SER	-	expression tag	UNP G0L7B8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0
2	G	1	Total Mg 1 1	0	0
2	H	1	Total Mg 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	13	Total O 13 13	0	0
3	B	9	Total O 9 9	0	0
3	C	12	Total O 12 12	0	0
3	D	8	Total O 8 8	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	9	Total O 9 9	0	0
3	F	11	Total O 11 11	0	0
3	G	19	Total O 19 19	0	0
3	H	11	Total O 11 11	0	0



• Molecule 1: Galactonate dehydratase



• Molecule 1: Galactonate dehydratase



D364				
K369				
Y373				
K376				
GLU				
ILE				
HIS				
LYS				
ASP				
GLY				
ALA				
THR				
LYS				
ILE				
ARG				
SER				

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.84Å 154.07Å 150.87Å 90.00° 104.38° 90.00°	Depositor
Resolution (Å)	49.17 – 2.79 49.17 – 2.79	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.17-2.79) 99.1 (49.17-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.81Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.231 , 0.266 0.263 , 0.299	Depositor DCC
R_{free} test set	4656 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	61.3	Xtrriage
Anisotropy	0.412	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 74.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	21906	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2777	0.72	0/3766
1	B	0.52	0/2765	0.73	0/3752
1	C	0.50	0/2773	0.72	0/3760
1	D	0.51	0/2767	0.72	0/3753
1	E	0.50	0/2791	0.71	0/3781
1	F	0.51	0/2797	0.73	0/3789
1	G	0.51	0/2786	0.72	0/3777
1	H	0.52	0/2801	0.73	0/3793
All	All	0.51	0/22257	0.72	0/30171

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2721	0	2689	31	0
1	B	2709	0	2668	34	0
1	C	2717	0	2690	38	0
1	D	2711	0	2679	26	0
1	E	2735	0	2722	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2741	0	2740	37	0
1	G	2730	0	2709	35	0
1	H	2745	0	2751	35	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	13	0	0	0	0
3	B	9	0	0	0	0
3	C	12	0	0	1	0
3	D	8	0	0	0	0
3	E	9	0	0	2	0
3	F	11	0	0	0	0
3	G	19	0	0	0	0
3	H	11	0	0	0	0
All	All	21906	0	21648	257	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 257 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:333:THR:O	1:F:338:ARG:NH2	2.01	0.93
1:F:33:ILE:HD12	1:F:302:ILE:HD12	1.61	0.82
1:H:1:SER:HB3	1:H:38:LEU:HD13	1.62	0.81
1:H:169:LEU:HD22	1:H:173:GLN:HG2	1.64	0.78
1:H:136:LEU:HB3	1:H:166:LYS:HD3	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/396 (87%)	322 (94%)	21 (6%)	0	100	100
1	B	343/396 (87%)	322 (94%)	21 (6%)	0	100	100
1	C	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	D	343/396 (87%)	323 (94%)	20 (6%)	0	100	100
1	E	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	F	343/396 (87%)	318 (93%)	24 (7%)	1 (0%)	41	72
1	G	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
1	H	343/396 (87%)	321 (94%)	22 (6%)	0	100	100
All	All	2744/3168 (87%)	2569 (94%)	174 (6%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	335	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/337 (84%)	250 (88%)	34 (12%)	5	15
1	B	281/337 (83%)	247 (88%)	34 (12%)	5	15
1	C	283/337 (84%)	248 (88%)	35 (12%)	4	14
1	D	282/337 (84%)	249 (88%)	33 (12%)	5	16
1	E	287/337 (85%)	254 (88%)	33 (12%)	5	17
1	F	289/337 (86%)	259 (90%)	30 (10%)	7	21
1	G	286/337 (85%)	250 (87%)	36 (13%)	4	14
1	H	290/337 (86%)	257 (89%)	33 (11%)	5	18
All	All	2282/2696 (85%)	2014 (88%)	268 (12%)	5	16

5 of 268 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	357	CYS
1	H	30	LYS
1	H	280	LYS
1	C	337	LEU
1	C	311	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	C	95	GLN
1	F	158	GLN
1	G	95	GLN
1	H	95	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/396 (88%)	0.46	37 (10%) 6 3	38, 69, 101, 116	0
1	B	351/396 (88%)	0.47	28 (7%) 12 6	39, 68, 99, 117	0
1	C	351/396 (88%)	0.49	25 (7%) 16 9	38, 68, 103, 132	0
1	D	351/396 (88%)	0.31	12 (3%) 45 35	39, 69, 102, 118	0
1	E	351/396 (88%)	0.30	11 (3%) 49 39	35, 64, 94, 118	0
1	F	351/396 (88%)	0.36	17 (4%) 30 21	35, 63, 93, 123	0
1	G	351/396 (88%)	0.42	15 (4%) 35 25	38, 64, 94, 106	0
1	H	351/396 (88%)	0.39	17 (4%) 30 21	38, 64, 94, 108	0
All	All	2808/3168 (88%)	0.40	162 (5%) 23 15	35, 66, 99, 132	0

The worst 5 of 162 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	169	LEU	6.3
1	C	137	TYR	5.9
1	B	159	GLY	5.0
1	B	137	TYR	5.0
1	C	177	TYR	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	F	401	1/1	0.81	0.13	64,64,64,64	0
2	MG	H	401	1/1	0.81	0.11	73,73,73,73	0
2	MG	C	401	1/1	0.82	0.14	59,59,59,59	0
2	MG	D	401	1/1	0.90	0.15	100,100,100,100	0
2	MG	G	401	1/1	0.93	0.06	61,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.