



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 05:14 am BST

PDB ID : 4OTN  
Title : Crystal structure of the C-terminal regulatory domain of murine GCN2  
Authors : He, H.; Georgiadis, M.M.  
Deposited on : 2014-02-13  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 40px;">4%      67%      21%      • 9%</p>
1	B	135	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 64%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 40px;">4%      64%      21%      • 12%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	1704	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 2-alpha kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	123	1135	719	192	223	1	0	13	0
1	B	119	1077	688	187	201	1	0	10	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1532	THR	ASN	SEE REMARK 999	UNP Q9QZ05
A	1534	SER	ILE	SEE REMARK 999	UNP Q9QZ05
A	1536	ILE	LEU	SEE REMARK 999	UNP Q9QZ05
A	1537	SER	ALA	SEE REMARK 999	UNP Q9QZ05
B	1532	THR	ASN	SEE REMARK 999	UNP Q9QZ05
B	1534	SER	ILE	SEE REMARK 999	UNP Q9QZ05
B	1536	ILE	LEU	SEE REMARK 999	UNP Q9QZ05
B	1537	SER	ALA	SEE REMARK 999	UNP Q9QZ05

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	79	Total O 79 79	0	0
4	B	90	Total O 90 90	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Eukaryotic translation initiation factor 2-alpha kinase 4



- Molecule 1: Eukaryotic translation initiation factor 2-alpha kinase 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.49Å 85.49Å 73.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.02 – 1.90 36.89 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (33.02-1.90) 99.6 (36.89-1.90)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.4_4	Depositor
R, $R_{free}$	0.195 , 0.236 0.194 , 0.229	Depositor DCC
$R_{free}$ test set	1252 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtrriage
Anisotropy	0.261	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 57.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2419	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/1152	0.51	0/1558
1	B	0.41	0/1094	0.56	0/1479
All	All	0.39	0/2246	0.54	0/3037

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1135	0	1136	30	0
1	B	1077	0	1104	36	1
2	A	10	0	0	0	0
3	A	8	0	12	0	0
3	B	20	0	30	6	0
4	A	79	0	0	7	0
4	B	90	0	0	6	2
All	All	2419	0	2282	61	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1546:ARG:O	4:B:1871:HOH:O	1.61	1.17
1:B:1543:ALA:N	3:B:1704:EDO:H12	1.75	1.01
1:B:1543:ALA:H	3:B:1704:EDO:H12	1.29	0.94
1:B:1613[B]:ARG:H	1:B:1613[B]:ARG:HD3	1.40	0.85
1:B:1549:HIS:N	4:B:1871:HOH:O	2.13	0.82
1:B:1542:SER:HB2	3:B:1704:EDO:H11	1.67	0.76
1:B:1567:LYS:HE3	4:B:1888:HOH:O	1.86	0.76
1:A:1592[B]:ASP:OD2	4:A:1860:HOH:O	2.08	0.70
1:A:1585[B]:GLN:OE1	4:A:1841:HOH:O	2.09	0.70
1:B:1613[B]:ARG:HD3	1:B:1613[B]:ARG:N	2.08	0.69
1:A:1640:ARG:HD2	4:A:1835:HOH:O	1.95	0.66
1:B:1612:GLN:O	4:B:1841:HOH:O	2.14	0.66
1:A:1553[B]:VAL:HG11	1:B:1573:ILE:HG21	1.79	0.65
1:A:1590:GLU:HG3	1:A:1601:THR:HG21	1.80	0.64
1:B:1576[A]:VAL:HG13	1:B:1614:TYR:OH	1.99	0.62
1:A:1539[A]:GLU:HG3	1:A:1540:LYS:N	2.14	0.62
1:B:1542:SER:HB2	3:B:1704:EDO:C1	2.29	0.62
1:B:1625:ILE:HG22	1:B:1631[B]:VAL:HG11	1.83	0.61
1:B:1625:ILE:HG22	1:B:1631[B]:VAL:CG1	2.34	0.58
1:B:1549:HIS:CA	4:B:1871:HOH:O	2.52	0.57
1:B:1625:ILE:CG2	1:B:1631[B]:VAL:HG11	2.34	0.57
1:A:1553[B]:VAL:HG22	1:B:1637:TYR:CE1	2.41	0.56
1:A:1637:TYR:OH	1:A:1642[A]:ASP:OD1	2.23	0.56
1:A:1541[B]:LEU:HD13	1:A:1546:ARG:NH2	2.20	0.55
1:A:1549:HIS:HD2	4:A:1830:HOH:O	1.90	0.54
1:B:1608:ARG:HD2	3:B:1702:EDO:H11	1.92	0.52
4:A:1810:HOH:O	1:B:1549:HIS:HE1	1.93	0.51
1:B:1550:GLU:N	4:B:1871:HOH:O	2.02	0.51
1:A:1527:THR:HG23	1:A:1528:THR:N	2.26	0.50
1:A:1604:GLN:O	1:A:1608[C]:ARG:HD3	2.11	0.50
1:A:1600:THR:O	1:A:1604:GLN:HG3	2.13	0.49
1:B:1605[A]:LEU:C	1:B:1605[A]:LEU:HD13	2.32	0.49
1:A:1553[A]:VAL:HG22	1:A:1557:LEU:HD22	1.95	0.48
1:B:1577:ASP:C	1:B:1640[A]:ARG:HH21	2.15	0.48
1:B:1561:LEU:O	1:B:1564:LEU:HB2	2.13	0.48
1:B:1537:SER:OG	1:B:1541:LEU:HG	2.13	0.47
1:B:1543:ALA:H	3:B:1704:EDO:C1	2.16	0.47
1:A:1637:TYR:CD2	1:A:1644:TYR:HB3	2.50	0.47
1:A:1594:ASP:HB3	1:A:1596:GLN:N	2.30	0.47
1:B:1577:ASP:O	1:B:1640[A]:ARG:NH2	2.38	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1541[B]:LEU:CD1	1:A:1546:ARG:NH2	2.79	0.46
1:A:1570[A]:GLU:O	1:A:1631:VAL:HG13	2.15	0.46
1:A:1549:HIS:O	1:A:1553[B]:VAL:HG23	2.15	0.46
1:B:1578:LEU:HA	1:B:1640[A]:ARG:NH2	2.31	0.46
1:B:1576[A]:VAL:CG1	1:B:1578:LEU:HG	2.46	0.46
1:A:1594:ASP:HB3	1:A:1596:GLN:H	1.79	0.46
1:A:1529:VAL:O	1:A:1531:PRO:HD3	2.16	0.45
1:B:1549:HIS:O	1:B:1553[A]:VAL:HG12	2.17	0.45
1:B:1576[A]:VAL:HG11	1:B:1578:LEU:HD12	1.98	0.45
1:A:1549:HIS:CD2	4:A:1830:HOH:O	2.69	0.44
1:A:1553[B]:VAL:CG1	1:B:1573:ILE:HG21	2.46	0.44
1:A:1539[A]:GLU:CG	1:A:1540:LYS:N	2.80	0.44
1:A:1637:TYR:HD2	1:A:1644:TYR:HB3	1.82	0.44
1:B:1637:TYR:OH	1:B:1642[A]:ASP:OD1	2.36	0.43
1:A:1553[B]:VAL:HG22	1:B:1637:TYR:CD1	2.53	0.43
1:B:1644:TYR:C	1:B:1644:TYR:CD2	2.92	0.43
1:A:1557:LEU:HD12	1:A:1557:LEU:HA	1.89	0.42
1:A:1606:LEU:CD2	1:A:1615:LEU:HB3	2.51	0.40
1:A:1567:LYS:HE2	4:A:1877:HOH:O	2.21	0.40
1:A:1617:LEU:HB3	1:B:1536:ILE:HD11	2.04	0.40
1:B:1606:LEU:HD11	1:B:1616:LYS:HD2	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1640[A]:ARG:NH2	4:B:1890:HOH:O[6_555]	1.66	0.54
4:B:1847:HOH:O	4:B:1882:HOH:O[5_555]	2.04	0.16

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/135 (100%)	131 (97%)	3 (2%)	1 (1%)	22	12
1	B	127/135 (94%)	125 (98%)	2 (2%)	0	100	100
All	All	262/270 (97%)	256 (98%)	5 (2%)	1 (0%)	34	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1593	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/127 (103%)	125 (95%)	6 (5%)	27	17
1	B	124/127 (98%)	115 (93%)	9 (7%)	14	6
All	All	255/254 (100%)	240 (94%)	15 (6%)	25	10

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1532	THR
1	A	1557	LEU
1	A	1592[A]	ASP
1	A	1592[B]	ASP
1	A	1594	ASP
1	A	1615	LEU
1	B	1556	ARG
1	B	1611	LYS
1	B	1613[A]	ARG
1	B	1613[B]	ARG
1	B	1615	LEU
1	B	1631[A]	VAL
1	B	1631[B]	VAL
1	B	1640[A]	ARG
1	B	1640[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1554	GLN
1	A	1563	ASN
1	B	1549	HIS
1	B	1563	ASN
1	B	1566	GLN
1	B	1599	ASN
1	B	1604	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	B	1703	-	3,3,3	0.48	0	2,2,2	0.11	0
2	SO4	A	1702	-	4,4,4	0.13	0	6,6,6	0.06	0
3	EDO	A	1704	-	3,3,3	0.63	0	2,2,2	1.06	0
3	EDO	B	1702	-	3,3,3	0.40	0	2,2,2	0.47	0
3	EDO	B	1705	-	3,3,3	0.56	0	2,2,2	2.35	1 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	1704	-	3,3,3	0.53	0	2,2,2	0.30	0
3	EDO	B	1701	-	3,3,3	0.49	0	2,2,2	0.28	0
2	SO4	A	1701	-	4,4,4	0.14	0	6,6,6	0.12	0
3	EDO	A	1703	-	3,3,3	0.59	0	2,2,2	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1703	-	-	0/1/1/1	-
3	EDO	A	1704	-	-	1/1/1/1	-
3	EDO	B	1702	-	-	1/1/1/1	-
3	EDO	B	1705	-	-	1/1/1/1	-
3	EDO	B	1704	-	-	1/1/1/1	-
3	EDO	B	1701	-	-	0/1/1/1	-
3	EDO	A	1703	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1705	EDO	O2-C2-C1	-2.82	91.59	111.91

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1704	EDO	O1-C1-C2-O2
3	B	1702	EDO	O1-C1-C2-O2
3	B	1705	EDO	O1-C1-C2-O2
3	B	1704	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1702	EDO	1	0
3	B	1704	EDO	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/135 (91%)	0.26	5 (4%) 37 40	25, 38, 66, 85	0
1	B	119/135 (88%)	0.18	6 (5%) 28 32	27, 38, 61, 84	0
All	All	242/270 (89%)	0.22	11 (4%) 33 36	25, 38, 65, 85	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1566	GLN	3.9
1	A	1541[A]	LEU	3.8
1	A	1540	LYS	3.4
1	B	1565	HIS	3.0
1	A	1526	GLY	2.7
1	B	1562	ALA	2.6
1	A	1543	ALA	2.5
1	A	1596	GLN	2.2
1	B	1596	GLN	2.2
1	B	1613[A]	ARG	2.0
1	B	1564	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	B	1704	4/4	0.65	0.20	43,45,49,52	0
3	EDO	B	1705	4/4	0.77	0.20	57,58,70,74	0
3	EDO	A	1704	4/4	0.77	0.22	57,65,67,69	0
2	SO4	A	1702	5/5	0.86	0.35	107,115,138,140	0
3	EDO	B	1702	4/4	0.90	0.20	36,48,54,70	0
2	SO4	A	1701	5/5	0.95	0.10	65,67,95,107	0
3	EDO	B	1701	4/4	0.96	0.28	57,62,66,81	0
3	EDO	B	1703	4/4	0.98	0.08	34,36,44,47	0
3	EDO	A	1703	4/4	0.98	0.09	32,33,35,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.