



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 4, 2023 – 01:38 am BST

PDB ID : 8OVR
Title : Clostridium perfringens chitinase CP56_3454 apo form
Authors : Bloch, Y.; Savvides, S.N.
Deposited on : 2023-04-26
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.33
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	630	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 9612 atoms, of which 4407 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chitinase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	572	8846	2852	4353	726	899	16	0	15	0

There are 31 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	600	PRO	-	expression tag	UNP F8UNI4
A	601	ARG	-	expression tag	UNP F8UNI4
A	602	ASN	-	expression tag	UNP F8UNI4
A	603	LYS	-	expression tag	UNP F8UNI4
A	604	GLY	-	expression tag	UNP F8UNI4
A	605	GLU	-	expression tag	UNP F8UNI4
A	606	LEU	-	expression tag	UNP F8UNI4
A	607	GLU	-	expression tag	UNP F8UNI4
A	608	GLY	-	expression tag	UNP F8UNI4
A	609	LYS	-	expression tag	UNP F8UNI4
A	610	PRO	-	expression tag	UNP F8UNI4
A	611	ILE	-	expression tag	UNP F8UNI4
A	612	PRO	-	expression tag	UNP F8UNI4
A	613	ASN	-	expression tag	UNP F8UNI4
A	614	PRO	-	expression tag	UNP F8UNI4
A	615	LEU	-	expression tag	UNP F8UNI4
A	616	LEU	-	expression tag	UNP F8UNI4
A	617	GLY	-	expression tag	UNP F8UNI4
A	618	LEU	-	expression tag	UNP F8UNI4
A	619	ASP	-	expression tag	UNP F8UNI4
A	620	SER	-	expression tag	UNP F8UNI4
A	621	THR	-	expression tag	UNP F8UNI4
A	622	ARG	-	expression tag	UNP F8UNI4
A	623	THR	-	expression tag	UNP F8UNI4
A	624	GLY	-	expression tag	UNP F8UNI4
A	625	HIS	-	expression tag	UNP F8UNI4
A	626	HIS	-	expression tag	UNP F8UNI4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	627	HIS	-	expression tag	UNP F8UNI4
A	628	HIS	-	expression tag	UNP F8UNI4
A	629	HIS	-	expression tag	UNP F8UNI4
A	630	HIS	-	expression tag	UNP F8UNI4

- Molecule 2 is PIPERAZINE-N,N'-BIS(2-ETHANESULFONIC ACID) (three-letter code: PIN) (formula: C₈H₁₈N₂O₆S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
2	A	1	Total	C	H	N	O	S	0	0
			36	8	18	2	6	2		
2	A	1	Total	C	H	N	O	S	0	0
			36	8	18	2	6	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Na	0	0
			2 2		

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	31	8	18	5	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	660	661	661	0	2

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	192.88Å 59.80Å 68.71Å 90.00° 108.18° 90.00°	Depositor
Resolution (Å)	45.09 – 1.60 45.09 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (45.09-1.60) 99.6 (45.09-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.153 , 0.172 0.152 , 0.172	Depositor DCC
R_{free} test set	2068 reflections (2.11%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtrriage
Anisotropy	0.280	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9612	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PIN, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.57	1/4638 (0.0%)	0.72	2/6288 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	160	GLU	CD-OE1	-6.37	1.18	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	186	ASP	CB-CG-OD2	-5.04	113.77	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4493	4353	4344	16	1
2	A	36	36	35	0	0
3	A	2	0	0	0	0
4	A	13	18	18	0	0
5	A	661	0	0	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5205	4407	4397	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ASN:ND2	5:A:802:HOH:O	2.31	0.63
1:A:458[B]:THR:HG22	5:A:1318:HOH:O	2.06	0.55
1:A:468:PRO:HB3	1:A:566:ILE:HG23	1.89	0.55
1:A:75:LYS:HD3	1:A:494:GLU:CD	2.32	0.49
1:A:282:LYS:HD3	5:A:1160:HOH:O	2.12	0.49
1:A:432:LYS:NZ	5:A:801:HOH:O	2.26	0.47
1:A:129[B]:ILE:HG13	1:A:130:ALA:N	2.32	0.45
1:A:583:GLN:HG3	5:A:998:HOH:O	2.19	0.43
1:A:518:ILE:HD11	1:A:571:ASN:CB	2.49	0.42
1:A:525:TYR:CG	1:A:526:PRO:HA	2.54	0.42
1:A:75:LYS:O	1:A:79:ILE:HG13	2.20	0.42
1:A:195:GLN:NE2	5:A:820:HOH:O	2.53	0.41
1:A:551:LYS:HE3	5:A:1111:HOH:O	2.21	0.41
1:A:170:LYS:O	1:A:348:GLY:HA2	2.21	0.40
1:A:63:ASP:HB3	1:A:71:ILE:O	2.21	0.40
1:A:79:ILE:C	1:A:79:ILE:HD12	2.40	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:LYS:HZ3	5:A:812:HOH:O[1_545]	1.60	0.00

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/630 (93%)	572 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/530 (92%)	482 (99%)	3 (1%)	86	77

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	LEU
1	A	464	CYS
1	A	488	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	ASN
1	A	195	GLN
1	A	460	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PG4	A	705	-	12,12,12	0.26	0	11,11,11	0.51	0
2	PIN	A	702	-	18,18,18	0.40	0	22,26,26	0.16	0
2	PIN	A	701	-	18,18,18	0.42	0	22,26,26	0.79	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG4	A	705	-	-	1/10/10/10	-
2	PIN	A	702	-	-	6/12/22/22	0/1/1/1
2	PIN	A	701	-	-	2/12/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	PIN	O3'-S1'-O1'	-2.16	106.00	111.27

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	PIN	C2'-C1'-S1'-O2'

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	701	PIN	C2'-C1'-S1'-O3'
2	A	702	PIN	C1-C2-N1-C3
2	A	702	PIN	C1-C2-N1-C4
2	A	702	PIN	C2-C1-S1-O3
2	A	702	PIN	C2-C1-S1-O1
2	A	702	PIN	C2-C1-S1-O2
4	A	705	PG4	C1-C2-O2-C3
2	A	702	PIN	C1'-C2'-N1'-C3'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/630 (90%)	0.10	25 (4%) 34 31	15, 24, 46, 71	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	472	SER	6.6
1	A	518	ILE	6.5
1	A	566	ILE	5.9
1	A	534	GLY	4.6
1	A	86	LEU	4.3
1	A	517	GLY	4.3
1	A	470	LYS	4.2
1	A	596	ASN	4.1
1	A	473	TRP	4.0
1	A	474	GLY	3.5
1	A	516	ASP	3.3
1	A	573	ILE	3.2
1	A	569	THR	3.2
1	A	476	GLY	3.1
1	A	568	ASP	3.1
1	A	572	ILE	2.9
1	A	389	GLU	2.8
1	A	563	ASP	2.6
1	A	536	TYR	2.6
1	A	535	ASP	2.4
1	A	312	GLY	2.4
1	A	570	ASN	2.4
1	A	370	THR	2.2
1	A	468	PRO	2.1
1	A	469	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PG4	A	705	13/13	0.88	0.20	34,44,72,72	0
2	PIN	A	701	18/18	0.93	0.12	28,38,49,76	0
2	PIN	A	702	18/18	0.94	0.30	45,54,65,71	36
3	NA	A	703	1/1	0.99	0.13	18,18,18,18	0
3	NA	A	704	1/1	1.00	0.07	16,16,16,16	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PIN A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PIN A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.