



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 13, 2021 – 09:08 pm GMT

PDB ID : 7OWI
Title : Crystal structure of dimeric chlorite dismutase variant R127A (CCld R127A)
from *Cyanothece* sp. PCC7425
Authors : Schmidt, D.; Mlynek, G.; Djinovic-Carugo, K.; Obinger, C.
Deposited on : 2021-06-18
Resolution : 1.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.24
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	 2% 90% 6% .
1	B	188	 4% 90% 6% .
1	C	188	 93% . .
1	D	188	 3% 93% . . 5%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13163 atoms, of which 6159 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chlorite dismutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	181	3026	970	1506	273	273	4	0	8	0
1	B	181	2978	957	1484	267	265	5	0	2	0
1	C	181	2960	952	1474	266	264	4	0	0	0
1	D	179	2942	946	1464	265	263	4	0	2	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP B8HNS6
A	-4	PRO	-	expression tag	UNP B8HNS6
A	-3	GLY	-	expression tag	UNP B8HNS6
A	-2	TYR	-	expression tag	UNP B8HNS6
A	-1	GLN	-	expression tag	UNP B8HNS6
A	0	ASP	-	expression tag	UNP B8HNS6
A	1	PRO	-	expression tag	UNP B8HNS6
A	127	ALA	ARG	engineered mutation	UNP B8HNS6
B	-5	GLY	-	expression tag	UNP B8HNS6
B	-4	PRO	-	expression tag	UNP B8HNS6
B	-3	GLY	-	expression tag	UNP B8HNS6
B	-2	TYR	-	expression tag	UNP B8HNS6
B	-1	GLN	-	expression tag	UNP B8HNS6
B	0	ASP	-	expression tag	UNP B8HNS6
B	1	PRO	-	expression tag	UNP B8HNS6
B	127	ALA	ARG	engineered mutation	UNP B8HNS6
C	-5	GLY	-	expression tag	UNP B8HNS6
C	-4	PRO	-	expression tag	UNP B8HNS6
C	-3	GLY	-	expression tag	UNP B8HNS6
C	-2	TYR	-	expression tag	UNP B8HNS6
C	-1	GLN	-	expression tag	UNP B8HNS6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	ASP	-	expression tag	UNP B8HNS6
C	1	PRO	-	expression tag	UNP B8HNS6
C	127	ALA	ARG	engineered mutation	UNP B8HNS6
D	-5	GLY	-	expression tag	UNP B8HNS6
D	-4	PRO	-	expression tag	UNP B8HNS6
D	-3	GLY	-	expression tag	UNP B8HNS6
D	-2	TYR	-	expression tag	UNP B8HNS6
D	-1	GLN	-	expression tag	UNP B8HNS6
D	0	ASP	-	expression tag	UNP B8HNS6
D	1	PRO	-	expression tag	UNP B8HNS6
D	127	ALA	ARG	engineered mutation	UNP B8HNS6

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	B	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	C	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	D	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	Total	C	H	O	0	0
			17	4	10	3		
3	C	1	Total	C	H	O	0	0
			17	4	10	3		
3	C	1	Total	C	H	O	0	0
			17	4	10	3		
3	D	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			22	6	14	2		
4	B	1	Total	C	H	O	0	0
			22	6	14	2		
4	C	1	Total	C	H	O	0	0
			22	6	14	2		
4	D	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			9	3	3	3		
5	B	1	Total	C	H	O	0	0
			9	3	3	3		
5	D	1	Total	C	H	O	0	0
			9	3	3	3		
5	D	1	Total	C	H	O	0	0
			9	3	3	3		
5	D	1	Total	C	H	O	0	0
			9	3	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	191	Total	O	0	0
			191	191		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	174	Total 174	O 174	0	0
6	C	224	Total 224	O 224	0	0
6	D	175	Total 175	O 175	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chlorite dismutase





- Molecule 1: Chlorite dismutase



- Molecule 1: Chlorite dismutase



- Molecule 1: Chlorite dismutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.45Å 72.67Å 112.12Å 90.00° 94.83° 90.00°	Depositor
Resolution (Å)	44.29 – 1.70 44.29 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.29-1.70) 99.9 (44.29-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.70Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.173 , 0.196 0.172 , 0.195	Depositor DCC
R_{free} test set	1177 reflections (1.23%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtrriage
Anisotropy	0.193	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13163	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 63.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.2113e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, MPD, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1584	0.61	0/2141
1	B	0.39	0/1540	0.59	0/2083
1	C	0.47	0/1521	0.62	0/2059
1	D	0.41	0/1517	0.59	0/2051
All	All	0.42	0/6162	0.60	0/8334

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1520	1506	1464	9	0
1	B	1494	1484	1470	6	0
1	C	1486	1474	1474	3	0
1	D	1478	1464	1458	5	0
2	A	43	30	30	1	0
2	B	43	30	30	2	0
2	C	43	30	30	1	0
2	D	43	30	30	1	0
3	A	7	10	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	14	20	20	0	0
3	D	7	10	10	0	0
4	A	8	14	14	0	0
4	B	8	14	14	3	0
4	C	8	14	14	0	0
4	D	8	14	14	2	0
5	B	12	6	16	0	0
5	D	18	9	24	0	0
6	A	191	0	0	2	2
6	B	174	0	0	1	0
6	C	224	0	0	0	0
6	D	175	0	0	3	1
All	All	7004	6159	6122	31	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:204:MPD:H12	4:B:204:MPD:H52	1.63	0.78
1:A:106[B]:ASP:OD1	1:A:110:ARG:NE	2.20	0.73
1:B:8:ILE:HD13	1:B:44:ILE:HG22	1.74	0.70
1:D:95:GLN:NE2	6:D:601:HOH:O	2.28	0.66
4:D:506:MPD:HM2	4:D:506:MPD:H52	1.77	0.66

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:772:HOH:O	6:D:714:HOH:O[2_445]	2.12	0.08
6:A:662:HOH:O	6:A:732:HOH:O[2_455]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/188 (99%)	183 (98%)	3 (2%)	0	100	100
1	B	181/188 (96%)	178 (98%)	3 (2%)	0	100	100
1	C	179/188 (95%)	175 (98%)	4 (2%)	0	100	100
1	D	177/188 (94%)	173 (98%)	4 (2%)	0	100	100
All	All	723/752 (96%)	709 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/158 (101%)	158 (99%)	2 (1%)	69	56
1	B	155/158 (98%)	152 (98%)	3 (2%)	57	41
1	C	153/158 (97%)	152 (99%)	1 (1%)	84	77
1	D	152/158 (96%)	150 (99%)	2 (1%)	69	56
All	All	620/632 (98%)	612 (99%)	8 (1%)	69	56

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	168	TRP
1	D	95	GLN
1	B	168	TRP
1	B	95	GLN
1	C	168	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	D	505	1	27,50,50	2.01	8 (29%)	17,82,82	1.46	3 (17%)
5	GOL	D	503	-	5,5,5	0.83	0	5,5,5	1.11	0
4	MPD	A	503	-	7,7,7	0.42	0	9,10,10	0.34	0
2	HEM	A	501	6,1	27,50,50	1.84	5 (18%)	17,82,82	1.80	4 (23%)
4	MPD	D	506	-	7,7,7	0.38	0	9,10,10	0.63	0
5	GOL	B	201	-	5,5,5	1.11	1 (20%)	5,5,5	1.11	0
3	PEG	C	503	-	6,6,6	0.23	0	5,5,5	0.03	0
5	GOL	B	202	-	5,5,5	0.84	0	5,5,5	0.89	0
3	PEG	C	501	-	6,6,6	0.22	0	5,5,5	0.12	0
5	GOL	D	502	-	5,5,5	0.89	0	5,5,5	1.02	0
2	HEM	B	203	1	27,50,50	2.00	6 (22%)	17,82,82	1.47	3 (17%)
2	HEM	C	502	6,1	27,50,50	1.92	6 (22%)	17,82,82	1.51	2 (11%)
3	PEG	D	504	-	6,6,6	0.20	0	5,5,5	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	C	504	-	7,7,7	0.32	0	9,10,10	0.33	0
4	MPD	B	204	-	7,7,7	0.35	0	9,10,10	0.53	0
5	GOL	D	501	-	5,5,5	0.69	0	5,5,5	0.98	0
3	PEG	A	502	-	6,6,6	0.22	0	5,5,5	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	D	505	1	-	0/6/54/54	-
5	GOL	D	503	-	-	2/4/4/4	-
4	MPD	A	503	-	-	0/5/5/5	-
2	HEM	A	501	6,1	-	0/6/54/54	-
4	MPD	D	506	-	-	1/5/5/5	-
5	GOL	B	201	-	-	0/4/4/4	-
3	PEG	C	503	-	-	1/4/4/4	-
5	GOL	B	202	-	-	2/4/4/4	-
3	PEG	C	501	-	-	1/4/4/4	-
5	GOL	D	502	-	-	2/4/4/4	-
2	HEM	B	203	1	-	0/6/54/54	-
2	HEM	C	502	6,1	-	0/6/54/54	-
3	PEG	D	504	-	-	2/4/4/4	-
4	MPD	C	504	-	-	0/5/5/5	-
4	MPD	B	204	-	-	1/5/5/5	-
5	GOL	D	501	-	-	2/4/4/4	-
3	PEG	A	502	-	-	2/4/4/4	-

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	502	HEM	C3B-C2B	-5.73	1.32	1.40
2	B	203	HEM	C3B-C2B	-4.97	1.33	1.40
2	D	505	HEM	C3B-C2B	-4.97	1.33	1.40
2	B	203	HEM	C3C-C2C	-4.53	1.34	1.40
2	A	501	HEM	C3B-C2B	-4.23	1.34	1.40

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	CAA-CBA-CGA	-3.42	106.93	112.67
2	C	502	HEM	CAA-CBA-CGA	-3.24	107.24	112.67
2	A	501	HEM	CMD-C2D-C1D	-2.83	124.11	128.46
2	A	501	HEM	CMA-C3A-C4A	-2.80	124.16	128.46
2	D	505	HEM	CMD-C2D-C1D	-2.69	124.33	128.46

There are no chirality outliers.

5 of 16 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	202	GOL	O1-C1-C2-C3
5	D	501	GOL	C1-C2-C3-O3
5	D	501	GOL	O2-C2-C3-O3
5	D	502	GOL	O1-C1-C2-C3
3	D	504	PEG	O2-C3-C4-O4


There are no ring outliers.


6 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	505	HEM	1	0
2	A	501	HEM	1	0
4	D	506	MPD	2	0
2	B	203	HEM	2	0
2	C	502	HEM	1	0
4	B	204	MPD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/188 (96%)	-0.20	3 (1%) 70 74	18, 28, 46, 76	0
1	B	181/188 (96%)	-0.20	7 (3%) 39 44	20, 31, 52, 100	0
1	C	181/188 (96%)	-0.31	0 100 100	18, 26, 39, 53	0
1	D	179/188 (95%)	-0.20	5 (2%) 53 57	18, 29, 49, 74	0
All	All	722/752 (96%)	-0.23	15 (2%) 63 67	18, 29, 47, 100	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	46	LEU	8.7
1	D	41	VAL	6.5
1	D	46	LEU	6.2
1	B	44	ILE	6.1
1	B	43	GLU	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	C	503	7/7	0.67	0.25	46,59,72,72	0
3	PEG	A	502	7/7	0.77	0.18	43,55,71,71	0
5	GOL	B	201	6/6	0.80	0.18	33,42,56,57	0
5	GOL	D	503	6/6	0.81	0.17	32,39,56,56	0
3	PEG	C	501	7/7	0.83	0.17	39,52,65,68	0
5	GOL	D	502	6/6	0.83	0.18	47,56,68,75	0
3	PEG	D	504	7/7	0.83	0.12	41,53,61,70	0
5	GOL	B	202	6/6	0.84	0.10	43,47,57,61	0
4	MPD	D	506	8/8	0.85	0.14	23,29,39,47	0
5	GOL	D	501	6/6	0.85	0.12	46,50,57,60	0
4	MPD	B	204	8/8	0.88	0.16	24,31,39,47	0
4	MPD	A	503	8/8	0.95	0.09	26,32,38,38	0
4	MPD	C	504	8/8	0.95	0.08	26,33,39,39	0
2	HEM	B	203	43/43	0.99	0.07	20,24,32,33	0
2	HEM	C	502	43/43	0.99	0.09	17,21,28,28	0
2	HEM	D	505	43/43	0.99	0.07	18,22,28,29	0
2	HEM	A	501	43/43	0.99	0.09	19,23,29,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 203:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.