



# Full wwPDB X-ray Structure Validation Report ⓘ

May 12, 2020 – 11:25 pm BST

PDB ID : 5OXZ  
Title : Crystal Structure of NeqN/NeqC complex from Nanoarchaeum equitans at 1.2Å  
Authors : Aparicio, D.; Perez-Luque, R.; Ribo, M.; Fita, I.  
Deposited on : 2017-09-07  
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	105	
2	B	33	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEQ068.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	90	865	562	133	169	1	0	16	0

- Molecule 2 is a protein called NEQ528.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	33	303	191	54	56	2	0	4	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	118	Total	O	0	0
			118	118		
3	B	33	Total	O	0	0
			33	33		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NEQ068



- Molecule 2: NEQ528



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.67Å 45.67Å 254.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.40 – 1.20 35.84 – 1.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.40-1.20) 99.9 (35.84-1.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.34 (at 1.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.137 , 0.156 0.137 , 0.156	Depositor DCC
$R_{free}$ test set	2482 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.1	Xtrriage
Anisotropy	0.437	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 57.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	1319	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.68	22/896 (2.5%)	1.46	18/1208 (1.5%)
2	B	1.38	2/307 (0.7%)	1.61	8/410 (2.0%)
All	All	1.61	24/1203 (2.0%)	1.50	26/1618 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	1
All	All	0	3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6[A]	GLU	CD-OE1	11.03	1.37	1.25
1	A	6[B]	GLU	CD-OE1	11.03	1.37	1.25
1	A	-7	PHE	CB-CG	10.31	1.68	1.51
1	A	8[A]	GLU	CD-OE1	-8.48	1.16	1.25
1	A	8[B]	GLU	CD-OE1	-8.48	1.16	1.25
1	A	-7	PHE	CG-CD2	8.10	1.50	1.38
1	A	69[A]	GLU	CD-OE1	-8.10	1.16	1.25
1	A	69[B]	GLU	CD-OE1	-8.10	1.16	1.25
2	B	17	GLU	CG-CD	-7.95	1.40	1.51
1	A	16	LYS	CA-CB	7.75	1.71	1.53
1	A	-7	PHE	CG-CD1	7.52	1.50	1.38
1	A	16	LYS	CD-CE	-7.24	1.33	1.51
1	A	56	TYR	CE2-CZ	-6.13	1.30	1.38
1	A	6[A]	GLU	CD-OE2	6.08	1.32	1.25
1	A	6[B]	GLU	CD-OE2	6.08	1.32	1.25
1	A	6[A]	GLU	CG-CD	5.61	1.60	1.51

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6[B]	GLU	CG-CD	5.61	1.60	1.51
1	A	61[A]	HIS	CB-CG	-5.58	1.40	1.50
1	A	61[B]	HIS	CB-CG	-5.58	1.40	1.50
2	B	32	ASP	CB-CG	5.56	1.63	1.51
1	A	64	GLU	CD-OE1	-5.29	1.19	1.25
1	A	70	ILE	CB-CG2	-5.28	1.36	1.52
1	A	16	LYS	CE-NZ	-5.17	1.36	1.49
1	A	16	LYS	CB-CG	-5.13	1.38	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-7	PHE	CB-CG-CD1	10.15	127.91	120.80
2	B	8	ARG	NE-CZ-NH2	-8.25	116.17	120.30
1	A	15[A]	LYS	CD-CE-NZ	7.81	129.66	111.70
1	A	15[B]	LYS	CD-CE-NZ	7.81	129.66	111.70
1	A	6[A]	GLU	CG-CD-OE1	7.41	133.11	118.30
1	A	6[B]	GLU	CG-CD-OE1	7.41	133.11	118.30
2	B	8	ARG	NE-CZ-NH1	7.30	123.95	120.30
1	A	8[A]	GLU	OE1-CD-OE2	7.27	132.03	123.30
1	A	8[B]	GLU	OE1-CD-OE2	7.27	132.03	123.30
2	B	2[A]	ARG	NE-CZ-NH1	-7.19	116.70	120.30
2	B	2[B]	ARG	NE-CZ-NH1	-7.19	116.70	120.30
1	A	22	ASP	CB-CG-OD1	6.32	123.99	118.30
1	A	-7	PHE	CB-CA-C	6.15	122.71	110.40
2	B	2[A]	ARG	NH1-CZ-NH2	5.52	125.47	119.40
2	B	2[B]	ARG	NH1-CZ-NH2	5.52	125.47	119.40
2	B	2[A]	ARG	NE-CZ-NH2	-5.40	117.60	120.30
2	B	2[B]	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	A	6[A]	GLU	OE1-CD-OE2	-5.37	116.85	123.30
1	A	6[B]	GLU	OE1-CD-OE2	-5.37	116.85	123.30
1	A	35[A]	GLU	CG-CD-OE1	5.33	128.97	118.30
1	A	35[B]	GLU	CG-CD-OE1	5.33	128.97	118.30
1	A	75[A]	LEU	CB-CG-CD1	5.14	119.74	111.00
1	A	75[B]	LEU	CB-CG-CD1	5.14	119.74	111.00
1	A	22	ASP	CB-CG-OD2	-5.08	113.72	118.30
1	A	66[A]	ASP	CB-CG-OD1	-5.07	113.73	118.30
1	A	66[B]	ASP	CB-CG-OD1	-5.07	113.73	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-7	PHE	Peptide
1	A	77	ALA	Mainchain
2	B	32	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	865	0	844	21	1
2	B	303	0	307	6	0
3	A	118	0	0	7	2
3	B	33	0	0	3	3
All	All	1319	0	1151	24	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8[B]:GLU:OE1	1:A:15[B]:LYS:CE	1.80	1.29
1:A:8[B]:GLU:OE1	1:A:15[B]:LYS:HE2	1.11	1.24
1:A:8[B]:GLU:CG	3:A:103:HOH:O	2.01	1.08
1:A:8[B]:GLU:HG2	3:A:103:HOH:O	1.61	1.00
1:A:6[B]:GLU:OE1	1:A:17:LYS:HE2	1.76	0.85
1:A:61[B]:HIS:HD2	3:B:102:HOH:O	1.58	0.84
1:A:8[B]:GLU:OE1	1:A:15[B]:LYS:HE3	1.85	0.75
1:A:61[B]:HIS:ND1	1:A:63[B]:THR:OG1	2.22	0.72
1:A:59:ILE:HB	2:B:13[B]:ASP:OD1	1.90	0.70
1:A:61[B]:HIS:CE1	1:A:63[B]:THR:HG1	2.17	0.62
1:A:61[B]:HIS:CD2	3:B:102:HOH:O	2.40	0.61
1:A:15[B]:LYS:HG2	3:A:103:HOH:O	2.01	0.59
2:B:20:LYS:NZ	3:B:101:HOH:O	2.40	0.54
1:A:61[B]:HIS:CE1	1:A:63[B]:THR:OG1	2.61	0.54
1:A:-4[B]:ILE:CD1	3:A:190:HOH:O	2.56	0.53
1:A:15[B]:LYS:CG	3:A:103:HOH:O	2.60	0.48
1:A:73:ILE:HD11	2:B:24[B]:ASN:ND2	2.29	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-4[B]:ILE:HD12	1:A:-3:TYR:N	2.31	0.45
2:B:1[B]:MET:HE3	2:B:1[B]:MET:HB3	1.47	0.44
2:B:13[A]:ASP:C	2:B:13[A]:ASP:OD1	2.56	0.44
1:A:8[B]:GLU:HG3	3:A:103:HOH:O	1.88	0.44
1:A:-4[B]:ILE:HD11	3:A:190:HOH:O	2.17	0.43
1:A:69[A]:GLU:HB3	2:B:2[A]:ARG:HB2	2.01	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:193:HOH:O	3:A:193:HOH:O[8_675]	1.04	1.16
3:B:106:HOH:O	3:B:106:HOH:O[12_564]	1.83	0.37
1:A:-1:ASP:OD1	3:B:129:HOH:O[12_564]	1.99	0.21
3:A:163:HOH:O	3:B:129:HOH:O[12_564]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/105 (100%)	101 (96%)	4 (4%)	0	100	100
2	B	34/33 (103%)	34 (100%)	0	0	100	100
All	All	139/138 (101%)	135 (97%)	4 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/94 (102%)	95 (99%)	1 (1%)	76	47
2	B	34/30 (113%)	30 (88%)	4 (12%)	5	0
All	All	130/124 (105%)	125 (96%)	5 (4%)	43	4

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-7	PHE
2	B	2[A]	ARG
2	B	2[B]	ARG
2	B	13[A]	ASP
2	B	13[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	90/105 (85%)	0.98	10 (11%) <b>5</b> <b>6</b>	8, 11, 35, 47	0
2	B	33/33 (100%)	0.16	2 (6%) <b>21</b> <b>21</b>	8, 11, 22, 51	0
All	All	123/138 (89%)	0.76	12 (9%) <b>7</b> <b>8</b>	8, 11, 35, 51	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-7	PHE	14.3
1	A	82	LEU	13.9
1	A	-5	VAL	11.5
1	A	81	VAL	9.2
1	A	-3	TYR	8.1
1	A	-4[A]	ILE	7.1
1	A	80	LEU	6.4
1	A	83	SER	6.4
1	A	-6	LYS	4.9
1	A	79	HIS	4.7
2	B	33	SER	3.7
2	B	32	ASP	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.