



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 05:39 am BST

PDB ID : 6OZU
Title : Crystal structure of the MIF4G domain of Trypanosoma cruzi translation initiation factor EIF4G5
Authors : Guimaraes, B.G.; Santos, L.P.C.
Deposited on : 2019-05-16
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	
1	B	241	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	409	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Eukaryotic translation initiation factor 4 gamma 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	235	Total	C	N	O	S	0	1	0
			1916	1224	345	340	7			
1	B	234	Total	C	N	O	S	0	0	0
			1880	1201	330	342	7			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	140	MET	-	initiating methionine	UNP A0A2V2WAU2
A	141	GLY	-	expression tag	UNP A0A2V2WAU2
A	372	ILE	-	expression tag	UNP A0A2V2WAU2
A	373	LEU	-	expression tag	UNP A0A2V2WAU2
A	374	GLU	-	expression tag	UNP A0A2V2WAU2
A	375	HIS	-	expression tag	UNP A0A2V2WAU2
A	376	HIS	-	expression tag	UNP A0A2V2WAU2
A	377	HIS	-	expression tag	UNP A0A2V2WAU2
A	378	HIS	-	expression tag	UNP A0A2V2WAU2
A	379	HIS	-	expression tag	UNP A0A2V2WAU2
A	380	HIS	-	expression tag	UNP A0A2V2WAU2
B	140	MET	-	initiating methionine	UNP A0A2V2WAU2
B	141	GLY	-	expression tag	UNP A0A2V2WAU2
B	372	ILE	-	expression tag	UNP A0A2V2WAU2
B	373	LEU	-	expression tag	UNP A0A2V2WAU2
B	374	GLU	-	expression tag	UNP A0A2V2WAU2
B	375	HIS	-	expression tag	UNP A0A2V2WAU2
B	376	HIS	-	expression tag	UNP A0A2V2WAU2
B	377	HIS	-	expression tag	UNP A0A2V2WAU2
B	378	HIS	-	expression tag	UNP A0A2V2WAU2
B	379	HIS	-	expression tag	UNP A0A2V2WAU2
B	380	HIS	-	expression tag	UNP A0A2V2WAU2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	51	Total O 51 51	0	0
4	B	44	Total O 44 44	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Eukaryotic translation initiation factor 4 gamma 5



- Molecule 1: Eukaryotic translation initiation factor 4 gamma 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	70.10Å 70.10Å 303.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.92 – 2.40 48.92 – 2.40	Depositor EDS
% Data completeness (in resolution range)	80.3 (48.92-2.40) 80.3 (48.92-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.39Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.201 , 0.243 0.217 , 0.259	Depositor DCC
R_{free} test set	1265 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	54.6	Xtrriage
Anisotropy	0.076	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3992	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1958	0.67	0/2645
1	B	0.50	0/1915	0.69	0/2590
All	All	0.51	0/3873	0.68	0/5235

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1916	0	1913	27	0
1	B	1880	0	1860	25	0
2	A	35	0	0	0	0
2	B	30	0	0	0	0
3	A	18	0	24	2	0
3	B	18	0	24	1	0
4	A	51	0	0	0	0
4	B	44	0	0	0	0
All	All	3992	0	3821	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:ASN:HB3	3:A:409:GOL:H32	1.72	0.71
1:B:224:ARG:HH11	3:B:407:GOL:H11	1.61	0.65
1:A:373:LEU:HD13	1:B:203:PRO:HB3	1.82	0.61
1:A:265:LEU:HD22	1:B:354:ARG:HB2	1.83	0.60
1:A:235:GLN:O	1:A:239:GLU:HB2	2.03	0.58
1:B:296:LEU:HD21	1:B:321:LEU:HD23	1.85	0.57
1:A:316:ILE:HG12	1:B:359:LEU:HD12	1.86	0.57
1:A:202:VAL:HG12	1:A:280:LEU:HD13	1.87	0.57
1:B:235:GLN:O	1:B:239:GLU:HB2	2.05	0.56
1:A:374:GLU:HG2	1:B:329:PHE:HD1	1.71	0.56
1:A:354:ARG:HB3	1:B:265:LEU:HD22	1.88	0.56
1:A:327:LYS:HD3	1:B:327:LYS:HB3	1.89	0.55
1:B:344:LEU:HD13	1:B:363:THR:HG22	1.90	0.53
1:A:352:PRO:HG2	1:A:357:GLN:HG3	1.91	0.52
1:A:359:LEU:HD12	1:B:316:ILE:HG12	1.91	0.51
1:A:372:ILE:HD11	1:B:203:PRO:HG2	1.94	0.50
1:A:300:LEU:HA	1:A:310:PRO:HB3	1.93	0.49
1:B:352:PRO:HG2	1:B:357:GLN:HG3	1.94	0.49
1:A:296:LEU:HD21	1:A:321:LEU:HD12	1.94	0.48
1:B:300:LEU:HA	1:B:310:PRO:HB3	1.94	0.48
1:A:315:LEU:HD13	1:B:315:LEU:HD13	1.95	0.48
1:B:315:LEU:O	1:B:319:MET:HG3	2.14	0.48
1:B:206:VAL:HG13	1:B:286:LEU:HD12	1.96	0.47
1:A:206:VAL:HG13	1:A:286:LEU:HD12	1.97	0.45
1:A:276:ASN:HB2	3:A:409:GOL:H2	1.99	0.45
1:A:315:LEU:O	1:A:319:MET:HG3	2.16	0.45
1:A:217:GLY:O	1:A:223:GLY:HA3	2.17	0.44
1:B:313:TYR:HA	1:B:316:ILE:HD12	1.99	0.44
1:B:345:GLU:O	1:B:348:MET:HB2	2.18	0.44
1:A:282:PHE:CD1	1:A:292:VAL:HG21	2.53	0.43
1:B:169:LEU:HD12	1:B:172:LEU:HD12	2.00	0.43
1:A:345:GLU:O	1:A:348:MET:HB2	2.18	0.43
1:A:366:THR:HG21	1:B:320:ASN:HA	2.00	0.43
1:B:309:PHE:HA	1:B:356:MET:HE1	2.00	0.42
1:A:327:LYS:HB2	1:B:323:LEU:HD11	2.00	0.42
1:A:169:LEU:HD12	1:A:172:LEU:HD12	2.02	0.42
1:A:248:LEU:O	1:A:252:ILE:HG12	2.20	0.41
1:B:309:PHE:CD1	1:B:310:PRO:HD2	2.56	0.41
1:B:344:LEU:HD13	1:B:363:THR:CG2	2.50	0.40
1:A:255:LEU:HD23	1:A:259:GLU:HB3	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:ASN:HD22	1:A:362:ASN:HA	1.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/241 (97%)	231 (99%)	3 (1%)	0	100	100
1	B	232/241 (96%)	226 (97%)	6 (3%)	0	100	100
All	All	466/482 (97%)	457 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/216 (95%)	198 (97%)	7 (3%)	37	56
1	B	199/216 (92%)	192 (96%)	7 (4%)	36	55
All	All	404/432 (94%)	390 (96%)	14 (4%)	36	55

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	200	ILE
1	A	201	PHE
1	A	255	LEU
1	A	290	LYS
1	A	321	LEU
1	A	357	GLN
1	A	359	LEU
1	B	180	LYS
1	B	198	ASP
1	B	200	ILE
1	B	249	GLU
1	B	255	LEU
1	B	357	GLN
1	B	359	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	228	ASN
1	A	362	ASN
1	A	375	HIS
1	B	195	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	406	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	A	407	-	4,4,4	0.11	0	6,6,6	0.13	0
2	SO4	B	402	-	4,4,4	0.20	0	6,6,6	0.20	0
3	GOL	B	409	-	5,5,5	0.09	0	5,5,5	0.25	0
3	GOL	A	409	-	5,5,5	0.13	0	5,5,5	0.48	0
2	SO4	B	405	-	4,4,4	0.20	0	6,6,6	0.09	0
2	SO4	A	405	-	4,4,4	0.18	0	6,6,6	0.12	0
3	GOL	B	407	-	5,5,5	0.05	0	5,5,5	0.31	0
3	GOL	A	410	-	5,5,5	0.07	0	5,5,5	0.20	0
2	SO4	B	401	-	4,4,4	0.22	0	6,6,6	0.09	0
2	SO4	A	401	-	4,4,4	0.13	0	6,6,6	0.14	0
2	SO4	A	406	-	4,4,4	0.17	0	6,6,6	0.06	0
2	SO4	A	402	-	4,4,4	0.21	0	6,6,6	0.24	0
3	GOL	A	408	-	5,5,5	0.11	0	5,5,5	0.21	0
2	SO4	A	403	-	4,4,4	0.13	0	6,6,6	0.25	0
2	SO4	B	403	-	4,4,4	0.16	0	6,6,6	0.15	0
3	GOL	B	408	-	5,5,5	0.09	0	5,5,5	0.39	0
2	SO4	B	404	-	4,4,4	0.20	0	6,6,6	0.10	0
2	SO4	A	404	-	4,4,4	0.12	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	409	-	-	0/4/4/4	-
3	GOL	B	407	-	-	1/4/4/4	-
3	GOL	A	410	-	-	0/4/4/4	-
3	GOL	B	409	-	-	0/4/4/4	-
3	GOL	A	408	-	-	2/4/4/4	-
3	GOL	B	408	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	408	GOL	O1-C1-C2-C3
3	B	407	GOL	O1-C1-C2-C3
3	A	408	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	409	GOL	2	0
3	B	407	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/241 (97%)	0.28	6 (2%) 56 54	41, 66, 97, 125	0
1	B	234/241 (97%)	0.47	22 (9%) 8 7	44, 76, 122, 144	0
All	All	469/482 (97%)	0.38	28 (5%) 21 20	41, 70, 111, 144	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	255	LEU	4.3
1	B	143	GLU	4.2
1	B	256	PRO	3.7
1	B	333	GLY	3.5
1	B	145	LYS	3.3
1	B	172	LEU	3.3
1	B	304	GLU	3.3
1	B	144	ARG	3.2
1	B	185	VAL	3.2
1	B	259	GLU	3.1
1	B	252	ILE	3.1
1	B	249	GLU	3.0
1	B	257	GLU	2.8
1	A	163	PRO	2.8
1	A	160	GLY	2.7
1	A	167	GLU	2.7
1	B	254	ASN	2.6
1	B	174	ILE	2.6
1	A	367	ILE	2.6
1	A	157	VAL	2.5
1	B	250	ARG	2.5
1	B	305	ARG	2.5
1	B	146	LEU	2.3
1	B	253	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	GLU	2.2
1	B	148	ASN	2.1
1	B	149	GLU	2.1
1	B	248	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	406	5/5	0.62	0.34	167,167,172,172	0
3	GOL	A	409	6/6	0.72	0.48	91,92,96,98	0
3	GOL	A	408	6/6	0.75	0.30	100,104,106,107	0
2	SO4	A	406	5/5	0.80	0.20	166,167,167,168	0
3	GOL	B	407	6/6	0.80	0.27	104,107,111,112	0
3	GOL	B	408	6/6	0.80	0.16	79,82,84,84	0
3	GOL	A	410	6/6	0.83	0.19	92,96,102,103	0
2	SO4	A	403	5/5	0.86	0.24	131,131,132,133	0
2	SO4	A	407	5/5	0.86	0.29	158,158,159,160	0
2	SO4	B	404	5/5	0.87	0.26	146,147,151,151	0
3	GOL	B	409	6/6	0.90	0.21	100,101,105,106	0
2	SO4	A	405	5/5	0.92	0.25	135,136,137,138	0
2	SO4	B	403	5/5	0.93	0.24	118,118,119,122	0
2	SO4	A	404	5/5	0.94	0.19	142,145,146,147	0
2	SO4	A	401	5/5	0.95	0.27	120,121,122,123	0
2	SO4	B	405	5/5	0.96	0.22	131,134,136,137	0
2	SO4	B	401	5/5	0.96	0.16	119,122,123,128	0
2	SO4	A	402	5/5	0.97	0.17	78,80,81,83	0
2	SO4	B	402	5/5	0.99	0.12	70,73,76,78	0

6.5 Other polymers

There are no such residues in this entry.