



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 08:56 AM EST

PDB ID : 2P2J
Title : Acetyl-CoA Synthetase, K609A mutation
Authors : Reger, A.S.; Gulick, A.M.
Deposited on : 2007-03-07
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	 2% 88% 8% ..
1	B	652	 % 88% 8% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetyl-coenzyme A synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	633	4923	3133	845	928	17	0	0	0
1	B	634	4913	3130	844	921	18	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	609	ALA	LYS	engineered mutation	UNP Q8ZKF6
B	609	ALA	LYS	engineered mutation	UNP Q8ZKF6

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	48	21	7	16	3	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	B	1	48	21	7	16	3	1	0	0

- Molecule 3 is ADENOSINE-5'-MONOPHOSPHATE-PROPYL ESTER (three-letter code: PRX) (formula: C₁₃H₂₀N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	26	13	5	7	1	0	0
3	B	1	26	13	5	7	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	208	208	208	0	0
4	B	171	171	171	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.67Å 94.50Å 167.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.30 47.69 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.3 (35.00-2.30) 95.1 (47.69-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.185 , 0.235 0.189 , 0.236	Depositor DCC
R_{free} test set	3147 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtrriage
Anisotropy	0.298	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 29.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10363	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, PRX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/5055	0.65	0/6893
1	B	0.57	0/5045	0.65	0/6881
All	All	0.59	0/10100	0.65	0/13774

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4923	0	4762	33	0
1	B	4913	0	4754	40	0
2	A	48	0	32	2	0
2	B	48	0	32	3	0
3	A	26	0	19	0	0
3	B	26	0	19	1	0
4	A	208	0	0	2	0
4	B	171	0	0	8	0
All	All	10363	0	9618	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:LEU:CD1	4:B:1055:HOH:O	2.16	0.93
1:B:402:ILE:HB	4:B:1055:HOH:O	1.71	0.91
1:B:383:LEU:HD13	4:B:1055:HOH:O	1.77	0.78
1:B:383:LEU:HD11	4:B:1055:HOH:O	1.81	0.76
1:A:191:ARG:NH1	2:A:990:COA:O4A	2.17	0.74
1:A:461:GLU:HB3	1:A:503:ARG:HB3	1.71	0.72
1:B:574:GLU:HB2	4:B:1164:HOH:O	1.93	0.67
1:B:398:TYR:O	4:B:1055:HOH:O	2.15	0.63
1:A:391:ASN:HD21	1:A:393:GLU:HG3	1.65	0.60
1:A:521:ASN:HD21	1:A:524:GLY:HA2	1.68	0.59
1:A:552:HIS:CD2	1:A:555:LYS:H	2.22	0.58
1:A:199:LYS:HG2	1:A:227:ILE:HD12	1.86	0.57
1:B:570:GLU:HG3	1:B:571:PRO:HD2	1.86	0.56
1:B:552:HIS:HD2	1:B:555:LYS:H	1.53	0.55
1:A:456:GLN:HB3	1:A:460:THR:HG21	1.89	0.55
1:B:221:LYS:HE3	1:B:224:GLY:HA2	1.88	0.54
1:A:391:ASN:HD21	1:A:393:GLU:CG	2.20	0.54
1:A:550:ILE:O	1:A:557:GLN:HA	2.09	0.53
1:A:313:HIS:HE1	4:A:1016:HOH:O	1.92	0.52
1:B:584:ARG:HH12	2:B:991:COA:P3B	2.32	0.52
1:B:611:MET:O	1:B:614:ILE:HG22	2.11	0.51
1:A:552:HIS:HD2	1:A:554:ILE:H	1.59	0.51
1:B:20:ASN:HB2	1:B:21:PRO:CD	2.40	0.51
1:B:544:GLU:HB3	1:B:564:THR:HB	1.93	0.49
1:B:76:ASN:HD22	1:B:79:ALA:H	1.59	0.49
1:B:397:TRP:CE2	1:B:401:LYS:HG3	2.48	0.48
1:A:391:ASN:ND2	1:A:393:GLU:CG	2.77	0.48
1:B:603:PRO:CD	1:B:615:LEU:HD21	2.44	0.48
1:B:584:ARG:NH1	2:B:991:COA:O7A	2.47	0.48
1:B:299:ASP:OD1	1:B:381:ARG:NH2	2.46	0.47
1:B:397:TRP:NE1	1:B:401:LYS:HG3	2.30	0.47
1:B:119:ARG:HD2	1:B:241:ILE:O	2.14	0.47
1:A:335:ASN:H	1:A:335:ASN:HD22	1.63	0.46
1:B:335:ASN:H	1:B:335:ASN:HD22	1.62	0.46
1:B:20:ASN:HB2	1:B:21:PRO:HD2	1.97	0.46
1:B:358:PRO:HG2	1:B:388:GLU:HG3	1.98	0.46
1:A:400:LYS:NZ	1:B:574:GLU:OE1	2.42	0.45
1:B:7:HIS:HB2	1:B:432:LEU:HB2	1.98	0.45
1:A:397:TRP:NE1	1:A:401:LYS:HG3	2.31	0.45
1:B:503:ARG:NH2	4:B:1128:HOH:O	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:THR:HG22	2:B:991:COA:H143	1.99	0.45
1:A:580:ARG:NH2	4:A:1064:HOH:O	2.49	0.44
1:A:20:ASN:HB2	1:A:21:PRO:CD	2.48	0.44
1:A:359:THR:HG22	2:A:990:COA:H143	2.00	0.43
1:B:552:HIS:CD2	1:B:555:LYS:H	2.32	0.43
1:A:604:LYS:HG2	1:A:610:ILE:HD13	2.00	0.43
1:B:611:MET:O	1:B:614:ILE:CG2	2.67	0.43
1:A:358:PRO:HG2	1:A:388:GLU:HG3	2.00	0.43
1:A:552:HIS:HD2	1:A:555:LYS:H	1.64	0.43
1:B:414:TRP:O	3:B:998:PRX:H3P3	2.19	0.42
1:A:552:HIS:CD2	1:A:554:ILE:H	2.38	0.42
1:B:552:HIS:CD2	1:B:554:ILE:H	2.38	0.42
1:A:76:ASN:HD21	1:A:78:ALA:HB3	1.85	0.42
1:A:161:VAL:HG12	1:A:308:GLY:HA2	2.02	0.42
1:B:603:PRO:HD3	1:B:615:LEU:HD21	2.02	0.42
1:B:370:LYS:HA	1:B:370:LYS:HD2	1.90	0.41
1:A:181:ARG:HA	1:A:181:ARG:HD3	1.90	0.41
1:A:539:HIS:HB3	1:A:542:ILE:HD12	2.02	0.41
1:B:50:THR:HG23	4:B:1001:HOH:O	2.20	0.41
1:B:159:HIS:CE1	1:B:260:PHE:HB3	2.55	0.41
1:B:552:HIS:HD2	1:B:554:ILE:H	1.68	0.41
1:A:74:THR:HB	1:A:252:ALA:HB1	2.02	0.41
1:B:490:SER:O	1:B:493:LYS:HE3	2.19	0.41
1:A:56:LYS:NZ	1:A:58:THR:OG1	2.45	0.41
1:B:206:LEU:HD22	1:B:211:VAL:HG11	2.03	0.41
1:A:390:ILE:O	1:A:390:ILE:HG23	2.21	0.41
1:B:603:PRO:HD3	1:B:641:LEU:HD11	2.03	0.41
1:A:515:ARG:HB3	1:A:517:ASP:OD2	2.20	0.40
1:A:391:ASN:ND2	1:A:393:GLU:HG2	2.35	0.40
1:A:300:ILE:HB	1:A:351:VAL:HA	2.03	0.40
1:B:634:ASP:O	1:B:637:VAL:HG22	2.20	0.40
1:A:261:ILE:HD13	1:A:261:ILE:HG21	1.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	629/652 (96%)	608 (97%)	20 (3%)	1 (0%)	47 58
1	B	630/652 (97%)	608 (96%)	20 (3%)	2 (0%)	41 50
All	All	1259/1304 (96%)	1216 (97%)	40 (3%)	3 (0%)	47 58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	633	ALA
1	A	438	THR
1	B	438	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	512/537 (95%)	499 (98%)	13 (2%)	47 65
1	B	509/537 (95%)	493 (97%)	16 (3%)	40 55
All	All	1021/1074 (95%)	992 (97%)	29 (3%)	43 60

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	HIS
1	A	76	ASN
1	A	127	LEU
1	A	247	GLU
1	A	316	LEU
1	A	333	VAL
1	A	335	ASN
1	A	494	ASN
1	A	503	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	506	GLU
1	A	521	ASN
1	A	612	ARG
1	A	646	GLN
1	B	50	THR
1	B	83	ASP
1	B	195	SER
1	B	262	LEU
1	B	267	SER
1	B	316	LEU
1	B	317	LEU
1	B	335	ASN
1	B	391	ASN
1	B	503	ARG
1	B	534	SER
1	B	570	GLU
1	B	585	LYS
1	B	614	ILE
1	B	632	LEU
1	B	645	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	104	GLN
1	A	313	HIS
1	A	335	ASN
1	A	344	GLN
1	A	391	ASN
1	A	454	HIS
1	A	521	ASN
1	A	552	HIS
1	A	646	GLN
1	B	53	GLN
1	B	76	ASN
1	B	335	ASN
1	B	344	GLN
1	B	456	GLN
1	B	552	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PRX	A	999	-	25,28,28	0.84	1 (4%)	27,41,41	1.15	1 (3%)
3	PRX	B	998	-	25,28,28	0.75	1 (4%)	27,41,41	1.33	1 (3%)
2	COA	A	990	-	41,50,50	1.71	3 (7%)	52,75,75	1.08	3 (5%)
2	COA	B	991	-	41,50,50	1.73	3 (7%)	52,75,75	1.16	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRX	A	999	-	-	0/11/31/31	0/3/3/3
3	PRX	B	998	-	-	1/11/31/31	0/3/3/3
2	COA	A	990	-	-	7/44/64/64	0/3/3/3
2	COA	B	991	-	-	8/44/64/64	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	991	COA	O9P-C9P	9.13	1.41	1.23
2	A	990	COA	O9P-C9P	8.98	1.41	1.23
2	B	991	COA	C2A-N3A	4.26	1.39	1.32
2	A	990	COA	C2A-N3A	4.18	1.38	1.32
2	A	990	COA	C2A-N1A	2.76	1.39	1.33
2	B	991	COA	C2A-N1A	2.53	1.38	1.33
3	B	998	PRX	O4'-C1'	2.34	1.44	1.41
3	A	999	PRX	O4'-C1'	2.01	1.43	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	998	PRX	N3-C2-N1	-5.51	120.06	128.68
2	B	991	COA	N3A-C2A-N1A	-5.07	120.76	128.68
2	A	990	COA	N3A-C2A-N1A	-4.64	121.42	128.68
3	A	999	PRX	N3-C2-N1	-4.38	121.83	128.68
2	B	991	COA	P2A-O3A-P1A	-3.93	119.34	132.83
2	A	990	COA	P2A-O3A-P1A	-3.21	121.80	132.83
2	B	991	COA	CEP-CBP-CAP	2.38	112.95	108.82
2	A	990	COA	CDP-CBP-CAP	2.08	112.43	108.82

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	990	COA	P1A-O3A-P2A-O6A
2	A	990	COA	C5P-C6P-C7P-N8P
2	A	990	COA	C2P-C3P-N4P-C5P
2	A	990	COA	S1P-C2P-C3P-N4P
2	B	991	COA	C3B-O3B-P3B-O8A
2	B	991	COA	C5B-O5B-P1A-O1A
2	B	991	COA	CAP-CBP-CCP-O6A
2	A	990	COA	O5P-C5P-N4P-C3P
2	A	990	COA	C6P-C5P-N4P-C3P
2	B	991	COA	CDP-CBP-CCP-O6A
2	B	991	COA	C5P-C6P-C7P-N8P
2	B	991	COA	CEP-CBP-CCP-O6A
2	B	991	COA	C5B-O5B-P1A-O3A
2	B	991	COA	C5B-O5B-P1A-O2A
3	B	998	PRX	O3P-C1P-C2P-C3P
2	A	990	COA	O4B-C4B-C5B-O5B



There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	998	PRX	1	0
2	A	990	COA	2	0
2	B	991	COA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	633/652 (97%)	-0.26	13 (2%) 63 70	14, 21, 37, 55	0
1	B	634/652 (97%)	-0.25	5 (0%) 86 89	16, 25, 36, 55	0
All	All	1267/1304 (97%)	-0.26	18 (1%) 75 80	14, 23, 36, 55	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	608	GLY	5.1
1	B	5	HIS	5.1
1	A	606	ARG	4.3
1	A	607	SER	4.1
1	A	635	PRO	3.9
1	A	609	ALA	3.7
1	A	605	THR	3.7
1	A	567	HIS	3.4
1	A	614	ILE	3.3
1	A	621	GLY	2.7
1	A	634	ASP	2.5
1	A	633	ALA	2.5
1	A	604	LYS	2.3
1	B	479	GLY	2.3
1	A	637	VAL	2.2
1	B	632	LEU	2.1
1	B	64	ASN	2.0
1	B	642	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	COA	A	990	48/48	0.86	0.25	27,48,64,66	0
2	COA	B	991	48/48	0.90	0.18	38,57,66,67	0
3	PRX	A	999	26/26	0.98	0.10	12,15,16,16	0
3	PRX	B	998	26/26	0.98	0.11	16,18,21,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [i](#)

There are no such residues in this entry.