



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 10:18 AM EST

PDB ID : 3P2N
Title : Discovery and structural characterization of a new glycoside hydrolase family abundant in coastal waters that was annotated as 'hypothetical protein'
Authors : Rebuffet, E.; Barbeyron, T.; Czjzek, M.; Michel, G.
Deposited on : 2010-10-03
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	 80% 8% • 11%
1	B	408	 75% 12% • 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3,6-anhydro-alpha-L-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	362	2922	1868	496	551	7	0	2	0
1	B	362	2925	1872	492	554	7	0	3	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	333	Total	O	0	0
			333	333		
4	B	339	Total	O	0	0
			339	339		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.99Å 96.80Å 126.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 1.95 45.22 – 1.95	Depositor EDS
% Data completeness (in resolution range)	94.8 (45.00-1.95) 94.8 (45.22-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.170 , 0.213 0.171 , 0.214	Depositor DCC
R_{free} test set	3182 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6523	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	4/3013 (0.1%)	0.96	8/4097 (0.2%)
1	B	1.17	3/3015 (0.1%)	0.99	7/4101 (0.2%)
All	All	1.15	7/6028 (0.1%)	0.97	15/8198 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	137	CYS	CB-SG	-6.42	1.71	1.82
1	A	368	GLU	CB-CG	-5.65	1.41	1.52
1	A	374	ARG	CG-CD	5.55	1.65	1.51
1	A	269	GLU	CB-CG	5.38	1.62	1.52
1	A	137	CYS	CB-SG	-5.36	1.73	1.81
1	B	254	CYS	CB-SG	5.31	1.91	1.82
1	B	269	GLU	CB-CG	5.08	1.61	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	374	ARG	NE-CZ-NH2	-18.18	111.21	120.30
1	B	374	ARG	NE-CZ-NH2	-15.67	112.47	120.30
1	A	374	ARG	NE-CZ-NH1	13.01	126.80	120.30
1	B	374	ARG	NE-CZ-NH1	11.98	126.29	120.30
1	A	374	ARG	CG-CD-NE	-7.67	95.70	111.80
1	B	236	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	399	ARG	NE-CZ-NH1	5.79	123.20	120.30
1	A	125	ASP	CB-CG-OD1	5.62	123.36	118.30
1	A	374	ARG	CD-NE-CZ	5.37	131.12	123.60
1	B	168	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	B	198	LYS	CD-CE-NZ	-5.22	99.70	111.70
1	B	236	ARG	NE-CZ-NH2	-5.16	117.72	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	ASP	CB-CG-OD1	5.12	122.91	118.30
1	B	358	GLY	N-CA-C	-5.09	100.37	113.10
1	A	242	LYS	CD-CE-NZ	-5.08	100.01	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2922	0	2792	23	0
1	B	2925	0	2797	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	333	0	0	4	0
4	B	339	0	0	4	0
All	All	6523	0	5589	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51:ASP:OD2	1:B:54:HIS:HD2	1.35	1.10
1:A:187:GLN:HE21	1:A:252:ASP:H	1.19	0.89
1:B:51:ASP:OD2	1:B:54:HIS:CD2	2.27	0.86
1:A:342:ASN:HD22	1:B:57:ALA:H	1.28	0.82
1:A:68:ASP:HB3	4:A:466:HOH:O	1.87	0.74
1:B:251:HIS:HB2	1:B:267:LYS:HE3	1.79	0.65
1:B:232:GLU:OE2	4:B:670:HOH:O	2.13	0.65
1:B:101[A]:ILE:HD11	1:B:377:LEU:HD12	1.80	0.64
1:A:187:GLN:NE2	1:A:252:ASP:H	1.93	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ILE:HD11	1:A:377:LEU:HG	1.79	0.63
1:A:139[B]:ILE:HD11	1:A:171:PHE:HA	1.81	0.61
1:A:309:HIS:HD2	1:A:327:ASP:OD2	1.87	0.57
1:B:198:LYS:HB3	1:B:223:PRO:HG2	1.86	0.57
1:B:218:GLU:HB3	1:B:219:PRO:HD2	1.87	0.57
1:A:198:LYS:HB3	1:A:223:PRO:HG2	1.88	0.56
1:B:38:THR:HG22	1:B:39:PHE:N	2.23	0.53
1:B:398:THR:HG22	1:B:398:THR:O	2.09	0.52
1:B:139[B]:ILE:HD11	1:B:171:PHE:HA	1.92	0.52
1:B:120:GLN:NE2	1:B:130:LYS:H	2.08	0.51
1:A:342:ASN:HD22	1:B:57:ALA:N	2.02	0.51
1:B:332:ASN:ND2	1:B:352:VAL:H	2.08	0.51
1:A:261:LYS:HD3	4:A:570:HOH:O	2.09	0.51
1:B:174:GLU:HG3	1:B:254:CYS:HB2	1.93	0.50
1:A:243:GLY:N	4:A:568:HOH:O	2.43	0.50
1:A:127:GLU:HG2	4:A:519:HOH:O	2.12	0.49
1:B:81:GLN:HG3	4:B:696:HOH:O	2.13	0.49
1:A:374:ARG:HD3	1:A:394:SER:O	2.13	0.48
1:B:120:GLN:HE21	1:B:130:LYS:H	1.62	0.48
1:A:399:ARG:HD2	1:B:237:PHE:CD1	2.49	0.48
1:B:111:TRP:CD1	1:B:141:TYR:HB3	2.48	0.47
1:B:353:ASN:HA	4:B:622:HOH:O	2.15	0.47
1:B:374:ARG:HD3	1:B:374:ARG:HA	1.63	0.47
1:A:73:TRP:CG	1:A:346:LYS:HB3	2.49	0.47
1:B:179:GLU:O	1:B:180:ASP:HB2	2.16	0.46
1:A:309:HIS:CD2	1:A:327:ASP:OD2	2.68	0.45
1:A:187:GLN:HE21	1:A:252:ASP:N	2.01	0.45
1:A:161:GLU:H	1:A:161:GLU:HG2	1.57	0.44
1:A:264:LEU:C	1:A:264:LEU:HD23	2.39	0.43
1:B:283:ARG:HG2	1:B:306:ASN:HA	1.99	0.43
1:B:101[A]:ILE:HD11	1:B:377:LEU:CD1	2.47	0.43
1:A:301:TYR:HA	1:B:54:HIS:O	2.20	0.42
1:A:177:LYS:HD2	1:A:182:TYR:CE1	2.54	0.42
1:B:101[B]:ILE:HG22	1:B:110:VAL:HB	2.02	0.42
1:B:245:PHE:CE2	1:B:302:ASN:HB3	2.55	0.41
1:B:230:LYS:NZ	4:B:669:HOH:O	2.08	0.41
1:B:336:TRP:HB3	1:B:346:LYS:HE3	2.03	0.41
1:A:64:LYS:HE3	1:A:64:LYS:HB2	1.89	0.41
1:A:107:LYS:HB2	1:A:107:LYS:HE2	1.51	0.41
1:B:140:TRP:CZ3	1:B:154:GLY:HA2	2.57	0.41
1:B:274:ALA:O	1:B:281:GLN:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:229:TRP:CH2	1:B:271:MET:HG2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/408 (89%)	348 (96%)	14 (4%)	0	100	100
1	B	363/408 (89%)	349 (96%)	13 (4%)	1 (0%)	41	30
All	All	725/816 (89%)	697 (96%)	27 (4%)	1 (0%)	51	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	273	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/348 (89%)	304 (98%)	5 (2%)	62	58
1	B	310/348 (89%)	305 (98%)	5 (2%)	62	58
All	All	619/696 (89%)	609 (98%)	10 (2%)	62	58

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	LYS
1	A	161	GLU
1	A	193	TYR
1	A	208	SER
1	A	374	ARG
1	B	76	GLN
1	B	161	GLU
1	B	171	PHE
1	B	193	TYR
1	B	398	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	76	GLN
1	A	187	GLN
1	A	309	HIS
1	A	342	ASN
1	A	382	ASN
1	B	54	HIS
1	B	120	GLN
1	B	194	ASN
1	B	332	ASN
1	B	335	GLN
1	B	382	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/408 (88%)	-0.37	3 (0%) 86 90	12, 20, 33, 54	0
1	B	362/408 (88%)	-0.44	3 (0%) 86 90	11, 18, 34, 52	0
All	All	724/816 (88%)	-0.40	6 (0%) 86 90	11, 19, 34, 54	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	398	THR	3.2
1	A	398	THR	2.8
1	A	38	THR	2.8
1	A	399	ARG	2.6
1	B	38	THR	2.6
1	B	37	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	900	1/1	1.00	0.03	29,29,29,29	0
2	ZN	B	900	1/1	1.00	0.02	27,27,27,27	0
3	CL	A	901	1/1	1.00	0.05	18,18,18,18	0
3	CL	B	901	1/1	1.00	0.06	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.