



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2021 – 06:45 PM EDT

PDB ID : 1P6K
Title : Rat neuronal NOS D597N mutant heme domain with L-N(omega)-nitroarginine-2,4-L-diaminobutyric amide bound
Authors : Flinspach, M.L.; Li, H.; Jamal, J.; Yang, W.; Huang, H.; Hah, J.-M.; Gomez-Vidal, J.A.; Litzinger, E.A.; Silverman, R.B.; Poulos, T.L.
Deposited on : 2003-04-29
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	
1	B	421	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric-oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	Total 3324	C 2124	N 570	O 608	S 22	0	6	0
1	B	410	Total 3357	C 2143	N 577	O 614	S 23	0	9	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	ASN	ASP	engineered mutation	UNP P29476
B	597	ASN	ASP	engineered mutation	UNP P29476

- Molecule 2 is D-MANNITOL (three-letter code: MTL) (formula: C₆H₁₄O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 12	C 6	O 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	B	1	12	6	6	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0
3	B	1	4	2	2	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	A	1	1	1	0	0

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 7 is L-N(OMEGA)-NITROARGININE-2,4-L-DIAMINOBTYRIC AMIDE (three-letter code: DP1) (formula: C₁₀H₂₂N₈O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			22	10	8	4		
7	B	1	Total	C	N	O	0	0
			22	10	8	4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	259	Total	O	0	0
			259	259		
8	B	314	Total	O	0	0
			314	314		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric-oxide synthase, brain



- Molecule 1: Nitric-oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.77Å 109.74Å 164.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.38 – 1.78 49.38 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.38-1.78) 97.6 (49.38-1.71)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.71Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.225 , 0.250 0.213 , 0.239	Depositor DCC
R_{free} test set	5019 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.9	Xtrriage
Anisotropy	0.647	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7451	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MTL, DP1, HEM, ZN, ACT, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3447	0.64	1/4675 (0.0%)
1	B	0.43	0/3495	0.66	1/4736 (0.0%)
All	All	0.42	0/6942	0.65	2/9411 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	589	MET	N-CA-C	-5.58	95.93	111.00
1	A	589	MET	N-CA-C	-5.49	96.18	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3237	67	0
1	B	3357	0	3279	39	0
2	A	12	0	14	1	0
2	B	12	0	14	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	43	0	30	4	0
5	B	43	0	30	0	0
6	A	17	0	15	0	0
6	B	17	0	15	0	0
7	A	22	0	21	2	0
7	B	22	0	21	1	0
8	A	259	0	0	7	0
8	B	314	0	0	5	0
All	All	7451	0	6682	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:373:GLY:H	1:B:377:HIS:HD2	1.11	0.92
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.51	0.91
1:A:396:LEU:HD22	1:A:577:LEU:HD12	1.54	0.86
1:A:350:THR:HB	1:A:353:GLN:HG3	1.62	0.82
1:B:373:GLY:H	1:B:377:HIS:CD2	2.01	0.74
1:A:307:GLU:HG3	8:B:941:HOH:O	1.88	0.74
1:A:396:LEU:CD2	1:A:577:LEU:HD12	2.21	0.70
1:A:503:GLU:HG3	8:A:1043:HOH:O	1.90	0.70
1:B:350:THR:HG22	1:B:352:ASP:H	1.55	0.70
1:A:350:THR:HG22	1:A:352:ASP:H	1.57	0.69
1:A:506:ILE:O	1:A:508:GLN:N	2.27	0.67
1:A:609:GLU:HG3	8:A:995:HOH:O	1.95	0.66
1:A:303:VAL:HG13	1:A:694:GLU:HB2	1.79	0.65
1:A:506:ILE:C	1:A:508:GLN:H	2.02	0.63
1:B:684:SER:HB3	1:B:687:ILE:HD11	1.81	0.61
1:A:299:ARG:HA	8:A:1045:HOH:O	2.00	0.61
1:B:513:PRO:HG2	1:B:518:ASP:CG	2.22	0.60
1:A:502:THR:O	1:A:506:ILE:HG12	2.02	0.60
1:A:361:PHE:O	1:A:364:GLN:HG2	2.02	0.59
1:B:355:PHE:N	1:B:356:PRO:HD2	2.18	0.59
1:B:621:THR:HG22	8:B:1165:HOH:O	2.03	0.59
1:A:545:PRO:HD3	8:A:1080:HOH:O	2.03	0.59
1:B:322:LEU:HB2	1:B:699:ARG:HB2	1.85	0.59
1:A:304:LYS:O	1:A:694:GLU:HG3	2.03	0.58
1:A:350:THR:HB	1:A:353:GLN:CG	2.33	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:614:MET:CE	1:A:632:GLU:HG3	2.34	0.57
1:A:664:ASN:OD1	1:A:667:ARG:NH2	2.38	0.56
1:A:684:SER:HB3	1:A:687:ILE:HD11	1.89	0.55
1:A:481[A]:ARG:NH2	7:A:790:DP1:O	2.40	0.55
1:A:306:TRP:NE1	1:B:336[B]:MET:HG3	2.22	0.55
1:A:473:ARG:NH2	1:A:710:PRO:HD3	2.23	0.54
5:A:750:HEM:HMC1	5:A:750:HEM:HBC2	1.89	0.54
1:B:684:SER:HB3	1:B:687:ILE:CG1	2.38	0.54
1:B:336[A]:MET:HG3	1:B:337:LEU:CD2	2.38	0.54
1:B:478:GLN:HB2	1:B:481[A]:ARG:HG3	1.90	0.53
1:A:545:PRO:HG2	1:A:547:ARG:NH1	2.24	0.52
1:B:546:ILE:HG12	1:B:560:LYS:HA	1.91	0.52
1:B:481[A]:ARG:NH2	7:B:791:DP1:O	2.44	0.51
1:B:485:TYR:CE1	1:B:514:ARG:HA	2.45	0.51
1:A:477[B]:SER:OG	1:A:481[B]:ARG:HD2	2.10	0.50
1:A:506:ILE:C	1:A:508:GLN:N	2.60	0.50
1:A:322:LEU:HD13	1:A:699:ARG:NH2	2.27	0.50
1:A:396:LEU:HD22	1:A:577:LEU:CD1	2.36	0.50
1:B:608:GLU:HG3	8:B:989:HOH:O	2.11	0.49
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.94	0.49
1:B:610:VAL:HG21	1:B:633:ILE:HD11	1.94	0.49
1:A:694:GLU:HB3	1:B:335:ILE:HD13	1.94	0.49
1:A:303:VAL:CG1	1:A:694:GLU:O	2.61	0.49
1:B:364:GLN:NE2	8:B:1140:HOH:O	2.46	0.48
1:A:355:PHE:N	1:A:356:PRO:HD2	2.28	0.48
1:A:501:PHE:HB2	2:A:870:MTL:H61	1.95	0.48
1:A:595:VAL:HG23	1:A:634:ASN:HD21	1.79	0.48
1:A:701:THR:HG23	8:A:994:HOH:O	2.13	0.48
1:A:516:ARG:HG2	1:A:517:PHE:CE1	2.48	0.48
1:A:684:SER:HB3	1:A:687:ILE:CG1	2.43	0.48
1:A:536:ILE:O	1:A:537:PRO:C	2.51	0.47
1:A:299:ARG:HE	1:A:318:LEU:HD22	1.79	0.47
1:B:684:SER:HB3	1:B:687:ILE:CD1	2.43	0.47
1:A:303:VAL:HG11	1:A:694:GLU:O	2.14	0.47
1:A:370:LYS:C	1:A:371:ARG:HG2	2.34	0.47
1:A:451:ASN:HB3	1:A:454:ASN:O	2.15	0.47
1:B:336[A]:MET:HG3	1:B:337:LEU:HD23	1.97	0.47
1:B:595:VAL:O	1:B:599:CYS:HB2	2.15	0.47
1:A:553:TRP:CE3	1:A:613:LYS:HD3	2.50	0.47
1:B:499:VAL:O	1:B:503:GLU:HG3	2.16	0.46
1:A:335:ILE:HB	1:A:338:PRO:HG3	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:316:LEU:HD23	1:A:319:LYS:HD2	1.97	0.46
1:A:436:HIS:HB2	8:A:1007:HOH:O	2.15	0.45
1:A:701:THR:HA	1:A:702:PRO:C	2.37	0.45
1:A:507:GLN:O	1:A:507:GLN:HG2	2.17	0.45
1:A:614:MET:HE1	1:A:632:GLU:HG3	1.98	0.45
1:B:355:PHE:CE1	1:B:385:ASN:HB2	2.52	0.45
1:A:715:VAL:O	1:A:715:VAL:HG23	2.16	0.45
1:B:475:TRP:HB2	1:B:523:LEU:HB3	1.99	0.45
1:A:447:LYS:HD2	8:A:1145:HOH:O	2.17	0.45
1:A:684:SER:HB3	1:A:687:ILE:CD1	2.47	0.44
1:B:606:ILE:HD11	1:B:633:ILE:HD13	1.98	0.44
1:A:301:LEU:CD1	1:B:330:ILE:HD13	2.47	0.44
1:A:524:LEU:O	1:A:531:PRO:HA	2.18	0.44
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.52	0.44
1:B:478:GLN:HB2	1:B:481[B]:ARG:CG	2.47	0.43
1:B:485:TYR:CZ	1:B:514:ARG:HA	2.54	0.43
1:B:528:GLY:O	1:B:717:LYS:HE2	2.19	0.43
1:B:565:PRO:HB3	1:B:588:TYR:CZ	2.54	0.42
1:B:321:THR:HG21	8:B:1063:HOH:O	2.19	0.42
1:A:308:THR:O	1:A:309:ASP:HB2	2.19	0.42
5:A:750:HEM:HBC2	5:A:750:HEM:CMC	2.49	0.42
1:A:684:SER:HB3	1:A:687:ILE:HG12	2.00	0.42
1:B:701:THR:HA	1:B:702:PRO:C	2.40	0.42
1:A:366:TYR:HA	1:A:369:ILE:HG12	2.02	0.42
1:B:299:ARG:HB3	1:B:299:ARG:HH11	1.85	0.42
1:A:622:SER:CB	1:B:642:GLN:HE22	2.33	0.42
5:A:750:HEM:HBB2	5:A:750:HEM:HHC	2.00	0.42
1:A:517:PHE:HB2	1:A:560:LYS:HE2	2.01	0.42
1:A:496:PRO:HA	1:A:499:VAL:HG23	2.01	0.42
5:A:750:HEM:O2A	7:A:790:DP1:HA	2.19	0.42
1:B:632:GLU:HA	1:B:635:ILE:HD12	2.01	0.41
1:A:595:VAL:O	1:A:599:CYS:HB2	2.20	0.41
1:A:676:TRP:CZ2	1:A:680:VAL:HG21	2.56	0.41
1:A:330:ILE:HD11	1:B:696:LEU:HB3	2.02	0.41
1:A:355:PHE:HD1	1:A:388:ILE:HD12	1.86	0.41
1:A:610:VAL:O	1:A:614:MET:HG3	2.21	0.40
1:B:355:PHE:N	1:B:356:PRO:CD	2.84	0.40
1:B:414:ARG:NH1	1:B:706:TYR:OH	2.53	0.40
1:A:485:TYR:CE2	1:A:512:ALA:HB1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/421 (97%)	388 (95%)	20 (5%)	1 (0%)	47	32
1	B	415/421 (99%)	404 (97%)	11 (3%)	0	100	100
All	All	824/842 (98%)	792 (96%)	31 (4%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	507	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/377 (98%)	361 (98%)	8 (2%)	52	36
1	B	375/377 (100%)	370 (99%)	5 (1%)	69	59
All	All	744/754 (99%)	731 (98%)	13 (2%)	59	48

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	316	LEU
1	A	396	LEU
1	A	485	TYR
1	A	493	LEU
1	A	523	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	527	ASN
1	A	547	ARG
1	A	552	ASP
1	B	316	LEU
1	B	454	ASN
1	B	535	GLN
1	B	547	ARG
1	B	552	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	451	ASN
1	A	454	ASN
1	A	500	GLN
1	A	508	GLN
1	A	634	ASN
1	A	697	ASN
1	B	364	GLN
1	B	377	HIS
1	B	425	GLN
1	B	451	ASN
1	B	454	ASN
1	B	507	GLN
1	B	527	ASN
1	B	535	GLN
1	B	634	ASN
1	B	642	GLN
1	B	697	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	H4B	B	761	-	16,18,18	2.09	4 (25%)	11,26,26	4.33	9 (81%)
3	ACT	A	860	-	1,3,3	2.75	1 (100%)	0,3,3	-	-
3	ACT	B	861	-	1,3,3	2.55	1 (100%)	0,3,3	-	-
5	HEM	B	750	1	27,50,50	1.61	5 (18%)	17,82,82	1.78	6 (35%)
7	DP1	B	791	-	17,21,21	0.87	1 (5%)	20,26,26	0.98	0
5	HEM	A	750	1	27,50,50	1.62	4 (14%)	17,82,82	1.76	6 (35%)
7	DP1	A	790	-	17,21,21	0.93	1 (5%)	20,26,26	0.88	0
2	MTL	B	871	-	11,11,11	0.86	0	14,14,14	0.77	1 (7%)
2	MTL	A	870	-	11,11,11	1.09	0	14,14,14	0.73	0
6	H4B	A	760	-	16,18,18	2.27	4 (25%)	11,26,26	4.17	9 (81%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	H4B	B	761	-	-	0/8/17/17	0/2/2/2
5	HEM	B	750	1	-	0/6/54/54	-
7	DP1	B	791	-	-	6/23/26/26	-
5	HEM	A	750	1	-	0/6/54/54	-
7	DP1	A	790	-	-	4/23/26/26	-
2	MTL	B	871	-	-	0/16/16/16	-
2	MTL	A	870	-	-	0/16/16/16	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	H4B	A	760	-	-	0/8/17/17	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	760	H4B	C4-N3	5.62	1.42	1.33
6	A	760	H4B	C4A-N5	4.75	1.47	1.38
6	B	761	H4B	C4-N3	4.69	1.41	1.33
5	B	750	HEM	C3C-C2C	-4.60	1.34	1.40
5	A	750	HEM	C3B-C2B	-4.35	1.34	1.40
6	B	761	H4B	C6-N5	4.29	1.54	1.45
6	A	760	H4B	C6-N5	3.93	1.53	1.45
5	A	750	HEM	C3C-CAC	-3.76	1.40	1.47
6	B	761	H4B	C4A-N5	3.69	1.45	1.38
5	B	750	HEM	C3B-C2B	-3.06	1.36	1.40
5	A	750	HEM	C3B-CAB	-2.78	1.42	1.47
3	A	860	ACT	CH3-C	2.75	1.52	1.48
5	B	750	HEM	C4A-NA	2.71	1.41	1.36
7	A	790	DP1	CZ-NH1	-2.65	1.28	1.35
5	A	750	HEM	C1A-NA	2.64	1.41	1.36
3	B	861	ACT	CH3-C	2.55	1.52	1.48
5	B	750	HEM	C3C-CAC	-2.41	1.42	1.47
7	B	791	DP1	CZ-NH1	-2.36	1.29	1.35
5	B	750	HEM	C3B-CAB	-2.35	1.43	1.47
6	B	761	H4B	C7-N8	2.18	1.48	1.44
6	A	760	H4B	C7-N8	2.02	1.48	1.44

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	761	H4B	C4-C4A-C8A	9.52	123.02	114.57
6	A	760	H4B	C4-C4A-C8A	9.09	122.64	114.57
6	B	761	H4B	C4-N3-C2	5.62	124.86	115.93
6	A	760	H4B	C4-N3-C2	5.04	123.94	115.93
6	A	760	H4B	N3-C2-N1	-4.58	118.23	125.42
6	B	761	H4B	N3-C2-N1	-4.57	118.25	125.42
6	A	760	H4B	C2-N1-C8A	3.81	123.08	114.54
6	B	761	H4B	C4A-C4-N3	-3.66	113.60	124.01
6	B	761	H4B	N2-C2-N1	3.58	122.81	117.25
6	A	760	H4B	C4A-C4-N3	-3.47	114.14	124.01
6	B	761	H4B	C4-C4A-N5	-3.39	116.28	119.12
6	A	760	H4B	N2-C2-N1	3.36	122.47	117.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	761	H4B	C2-N1-C8A	3.32	121.99	114.54
5	B	750	HEM	CBA-CAA-C2A	-3.26	106.47	112.49
5	A	750	HEM	CMD-C2D-C1D	-3.19	123.57	128.46
6	A	760	H4B	C4A-N5-C6	-3.16	112.56	121.16
5	A	750	HEM	CBD-CAD-C3D	-3.02	106.91	112.48
6	B	761	H4B	C4A-N5-C6	-3.00	113.00	121.16
5	B	750	HEM	C3B-C4B-NB	2.79	112.82	109.21
5	B	750	HEM	C4A-C3A-C2A	-2.65	105.15	107.00
6	A	760	H4B	C4-C4A-N5	-2.59	116.95	119.12
5	B	750	HEM	C4C-C3C-C2C	-2.56	105.11	106.90
5	A	750	HEM	CBA-CAA-C2A	-2.41	108.04	112.49
6	A	760	H4B	O9-C9-C6	2.38	114.66	108.98
5	A	750	HEM	CMA-C3A-C4A	-2.29	124.95	128.46
5	B	750	HEM	C3C-C4C-NC	2.23	115.15	110.94
5	A	750	HEM	CMD-C2D-C3D	2.14	128.98	124.94
5	B	750	HEM	CMA-C3A-C2A	2.05	128.81	124.94
6	B	761	H4B	O9-C9-C6	2.05	113.89	108.98
5	A	750	HEM	C3B-C4B-NB	2.05	111.86	109.21
2	B	871	MTL	C5-C4-C3	-2.04	109.28	112.47

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	790	DP1	O-C-CA-CB
7	A	790	DP1	N'-C-CA-CB
7	B	791	DP1	N'-C-CA-CB
7	B	791	DP1	O-C-CA-N
7	A	790	DP1	O-C-CA-N
7	B	791	DP1	O-C-CA-CB
7	A	790	DP1	N'-C-CA-N
7	B	791	DP1	N'-C-CA-N
7	B	791	DP1	CA-CB-CG-CD
7	B	791	DP1	C'-CA'-CB'-C1

There are no ring outliers.

4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	791	DP1	1	0
5	A	750	HEM	4	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	790	DP1	2	0
2	A	870	MTL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/421 (96%)	0.63	39 (9%) 8 7	24, 39, 59, 70	0
1	B	410/421 (97%)	0.36	18 (4%) 34 32	23, 33, 50, 66	0
All	All	817/842 (97%)	0.49	57 (6%) 16 16	23, 35, 56, 70	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	PHE	7.2
1	A	299	ARG	6.9
1	B	348	VAL	5.8
1	A	352	ASP	5.3
1	B	321	THR	5.0
1	A	488	PRO	4.7
1	B	352	ASP	4.6
1	A	355	PHE	4.3
1	A	321	THR	3.7
1	A	503	GLU	3.4
1	B	351	LYS	3.4
1	B	350	THR	3.4
1	B	300	PHE	3.4
1	A	489	ASP	3.1
1	A	391	THR	3.1
1	B	389	GLU	3.1
1	A	551	PHE	3.0
1	B	620	LYS	3.0
1	A	351	LYS	3.0
1	A	303	VAL	3.0
1	A	715	VAL	3.0
1	B	338	PRO	2.9
1	A	373	GLY	2.8
1	B	619	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	355	PHE	2.7
1	A	469	LYS	2.7
1	A	328	GLU	2.6
1	A	619	ARG	2.6
1	A	552	ASP	2.6
1	A	716	TRP	2.5
1	B	615	ASP	2.5
1	A	386	LYS	2.5
1	A	487	GLN	2.5
1	A	554	PHE	2.5
1	B	667	ARG	2.4
1	A	556	ASP	2.4
1	A	370	LYS	2.4
1	A	319	LYS	2.3
1	B	551	PHE	2.3
1	B	386	LYS	2.3
1	A	350	THR	2.3
1	A	490	GLY	2.3
1	A	385	ASN	2.3
1	A	510	TRP	2.2
1	A	371	ARG	2.2
1	A	393	THR	2.1
1	A	470	HIS	2.1
1	A	508	GLN	2.1
1	B	392[A]	SER	2.1
1	A	609	GLU	2.1
1	B	322	LEU	2.1
1	B	715	VAL	2.1
1	A	369	ILE	2.1
1	A	504	ILE	2.1
1	A	533	LEU	2.0
1	A	511	LYS	2.0
1	A	493	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MTL	A	870	12/12	0.80	0.17	51,51,52,52	0
7	DP1	A	790	22/22	0.85	0.16	39,43,45,45	0
7	DP1	B	791	22/22	0.86	0.16	41,43,44,44	0
2	MTL	B	871	12/12	0.92	0.09	45,46,47,47	0
3	ACT	A	860	4/4	0.95	0.18	60,60,60,60	0
3	ACT	B	861	4/4	0.96	0.13	45,45,45,45	0
6	H4B	A	760	17/17	0.96	0.09	24,25,26,26	0
6	H4B	B	761	17/17	0.97	0.09	24,25,27,27	0
5	HEM	A	750	43/43	0.98	0.09	25,26,27,27	0
5	HEM	B	750	43/43	0.98	0.10	23,23,26,28	0
4	ZN	A	900	1/1	0.99	0.10	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around DP1 B 791:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 750:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.