

# wwPDB X-ray Structure Validation Summary Report (i)

#### Jun 13, 2024 – 07:30 AM EDT

PDB ID : 1P8C

Title: Crystal structure of TM1620 (APC4843) from Thermotoga maritima

Authors: Kim, Y.; Joachimiak, A.; Brunzelle, J.S.; Korolev, S.V.; Edwards, A.; Xu, X.;

Savchenko, A.; Midwest Center for Structural Genomics (MCSG)

Deposited on : 2003-05-06

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467 Xtriage (Phenix): 1.13

EDS : 2.36.2

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Qua	ality of chain		
1	A	145	38%	35%	•	23%
1	В	145	37%	38%	·	21%
1	С	145	48%	31%	·	18%
1	D	145	37%	39%	6%	19%
1	Е	145	46%	34%		18%



Continued from previous page...

Mol	Chain	Length			Quality of chain		
			8%				
1	F	145		30%	46%	•	21%



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called conserved hypothetical protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	111	Total	С	N	О	S	0	0	0
1	A	111	878	549	159	165	5	0	0	0
1	В	114	Total	С	N	О	S	0	0	0
1	Ъ	114	910	574	164	167	5	0	0	0
1	С	119	Total	С	N	О	S	0	0	0
1		119	949	597	169	178	5	0	0	0
1	D	118	Total	С	N	О	S	0	0	0
1	ע	110	943	594	168	176	5	0	0	
1	Е	119	Total	С	N	О	S	0	0	0
1	12	119	949	597	169	178	5	0	0	0
1	1 F	115	Total	С	N	О	S	0	0	0
1		110	919	579	165	170	5		U	0

There are 144 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	CLONING ARTIFACT	UNP Q9X1V5
A	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
A	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
A	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
A	-17	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5
A	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
A	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
A	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
A	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
A	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
A	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
A	-3	PHE	_	CLONING ARTIFACT	UNP Q9X1V5
A	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
A	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
A	0	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
A	122	GLY	_	CLONING ARTIFACT	UNP Q9X1V5
A	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5
В	-21	MET	-	CLONING ARTIFACT	UNP Q9X1V5
В	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
В	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
В	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
В	-17	HIS	_	CLONING ARTIFACT	UNP Q9X1V5
В	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
В	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
В	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
В	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
В	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
В	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5
В	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
В	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
В	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
В	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
В	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
В	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5
В	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
В	-3	PHE	-	CLONING ARTIFACT	UNP Q9X1V5
В	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
В	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
В	0	HIS	_	CLONING ARTIFACT	UNP Q9X1V5
В	122	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
В	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5
С	-21	MET	_	CLONING ARTIFACT	UNP Q9X1V5
С	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
С	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
С	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
С	-17	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
С	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
С	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
С	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
С	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
С	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
С	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5
С	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
С	-3	PHE	-	CLONING ARTIFACT	UNP Q9X1V5
С	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
С	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
С	0	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
С	122	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
С	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5
D	-21	MET	-	CLONING ARTIFACT	UNP Q9X1V5
D	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
D	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
D	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
D	-17	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5
D	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
D	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
D	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
D	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
D	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
D	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5
D	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
D	-3	PHE	-	CLONING ARTIFACT	UNP Q9X1V5
D	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
D	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
D	0	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
D	122	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
D	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5
E	-21	MET	-	CLONING ARTIFACT	UNP Q9X1V5
E	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
E	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
E	-17	HIS	-	CLONING ARTIFACT	UNP Q9X1V5



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
Е	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-3	PHE	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
Е	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
Е	0	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
Е	122	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
Е	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5
F	-21	MET	-	CLONING ARTIFACT	UNP Q9X1V5
F	-20	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
F	-19	SER	-	CLONING ARTIFACT	UNP Q9X1V5
F	-18	SER	-	CLONING ARTIFACT	UNP Q9X1V5
F	-17	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-16	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-15	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-14	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-13	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-12	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	-11	SER	-	CLONING ARTIFACT	UNP Q9X1V5
F	-10	SER	-	CLONING ARTIFACT	UNP Q9X1V5
F	-9	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
F	-8	ARG	-	CLONING ARTIFACT	UNP Q9X1V5
F	-7	GLU	-	CLONING ARTIFACT	UNP Q9X1V5
F	-6	ASN	-	CLONING ARTIFACT	UNP Q9X1V5
F	-5	LEU	-	CLONING ARTIFACT	UNP Q9X1V5
F	-4	TYR	-	CLONING ARTIFACT	UNP Q9X1V5
F	-3	PHE	-	CLONING ARTIFACT	UNP Q9X1V5
F	-2	GLN	-	CLONING ARTIFACT	UNP Q9X1V5
F	-1	GLY	-	CLONING ARTIFACT	UNP Q9X1V5
F	0	HIS	-	CLONING ARTIFACT	UNP Q9X1V5
F	122	GLY	-	CLONING ARTIFACT	UNP Q9X1V5



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	123	SER	-	CLONING ARTIFACT	UNP Q9X1V5

### • Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	10	Total O 10 10	0	0
2	В	17	Total O 17 17	0	0
2	C	20	Total O 20 20	0	0
2	D	14	Total O 14 14	0	0
2	E	19	Total O 19 19	0	0
2	F	7	Total O 7 7	0	0




# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: conserved hypothetical protein











# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	65.82Å 82.54Å 135.40Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.10 - 2.30	Depositor
rtesolution (A)	48.10 - 2.15	EDS
% Data completeness	90.6 (48.10-2.30)	Depositor
(in resolution range)	87.3 (48.10-2.15)	EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.21 (at 2.16Å)	Xtriage
Refinement program	CNS 1.1	Depositor
P. P.	0.225 , 0.290	Depositor
$R, R_{free}$	0.225 , $0.289$	DCC
$R_{free}$ test set	3051 reflections (7.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.5	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 63.1	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5635	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.43	0/887	0.65	0/1191
1	В	0.44	0/921	0.69	0/1235
1	С	0.54	0/960	0.69	0/1288
1	D	0.47	0/954	0.64	0/1280
1	Е	0.50	0/960	0.68	0/1288
1	F	0.39	0/930	0.61	0/1247
All	All	0.47	0/5612	0.66	0/7529

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	878	0	899	54	0
1	В	910	0	934	72	0
1	С	949	0	969	48	0
1	D	943	0	964	70	0
1	Ε	949	0	969	46	0
1	F	919	0	940	92	0
2	A	10	0	0	0	0
2	В	17	0	0	5	0
2	С	20	0	0	0	0
2	D	14	0	0	3	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Е	19	0	0	0	0
2	F	7	0	0	1	0
All	All	5635	0	5675	308	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 308 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:D:107:GLU:HB3	1:F:20:ARG:HH12	1.29	0.98
1:F:22:THR:HG22	1:F:23:LEU:H	1.26	0.97
1:A:22:THR:HG22	1:A:24:ASN:H	1.29	0.95
1:E:92:GLY:O	1:E:95:VAL:HG12	1.70	0.91
1:D:22:THR:HG22	1:D:24:ASN:H	1.36	0.90

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	109/145 (75%)	97 (89%)	10 (9%)	2 (2%)	8 7
1	В	112/145~(77%)	101 (90%)	10 (9%)	1 (1%)	17 20
1	С	117/145 (81%)	109 (93%)	4 (3%)	4 (3%)	3 2
1	D	116/145 (80%)	106 (91%)	8 (7%)	2 (2%)	9 8
1	E	117/145 (81%)	109 (93%)	8 (7%)	0	100 100
1	F	113/145 (78%)	105 (93%)	7 (6%)	1 (1%)	17 20
All	All	684/870 (79%)	627 (92%)	47 (7%)	10 (2%)	10 10



5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	VAL
1	A	116	GLY
1	D	116	GLY
1	F	10	ARG
1	С	6	PHE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	F	erce	entiles
1	A	97/126 (77%)	89 (92%)	8 (8%)		11	14
1	В	99/126 (79%)	94 (95%)	5 (5%)		24	33
1	С	104/126 (82%)	99 (95%)	5 (5%)		25	36
1	D	103/126 (82%)	94 (91%)	9 (9%)		10	12
1	Е	104/126 (82%)	99 (95%)	5 (5%)		25	36
1	F	100/126 (79%)	95 (95%)	5 (5%)		24	34
All	All	607/756 (80%)	570 (94%)	37 (6%)		18	25

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	25	THR
1	F	81	PHE
1	Е	68	VAL
1	F	12	GLU
1	В	95	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	98	HIS
1	Е	30	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	F	72	GLN
1	F	30	ASN
1	С	30	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	111/145 (76%)	0.55	13 (11%) 4 6	30, 58, 129, 160	0
1	В	114/145 (78%)	0.51	17 (14%) 2 3	26, 53, 141, 179	0
1	С	119/145 (82%)	-0.03	1 (0%) 86 89	21, 43, 102, 127	0
1	D	118/145 (81%)	0.05	4 (3%) 45 52	30, 49, 104, 153	0
1	E	119/145 (82%)	-0.03	1 (0%) 86 89	30, 48, 93, 118	0
1	F	115/145 (79%)	0.58	11 (9%) 8 10	30, 64, 136, 161	0
All	All	696/870 (80%)	0.27	47 (6%) 17 22	21, 52, 123, 179	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	9	ALA	13.3
1	A	116	GLY	5.8
1	A	118	THR	5.8
1	D	5	LYS	5.5
1	A	13	LEU	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.



# 6.5 Other polymers (i)

There are no such residues in this entry.

