



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2024 – 05:03 pm BST

PDB ID : 8PBO  
Title : Deep interactome learning for generative drug design  
Authors : Hakansson, M.; Focht, D.; Atz, K.; Schneider, G.  
Deposited on : 2023-06-09  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
1	B	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	508	-	-	X	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 9061 atoms, of which 4512 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	265	4384	1394	2233	346	400	11	46	6	0
1	B	263	4370	1389	2228	346	396	11	50	6	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	GLY	-	expression tag	UNP P37231
B	203	GLY	-	expression tag	UNP P37231

- Molecule 2 is 3-[2-fluoranyl-4-[3-[2-fluoranyl-4-(5-methyl-1,3,4-thiadiazol-2-yl)phenoxy]propoxy]phenyl]propanoic acid (three-letter code: Y5I) (formula: C<sub>21</sub>H<sub>20</sub>F<sub>2</sub>N<sub>2</sub>O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	F	H	N	O			S
2	A	1	49	21	2	19	2	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	14	3	8	3	2	0
3	A	1	14	3	8	3	2	0
3	A	1	14	3	8	3	2	0
3	B	1	14	3	8	3	2	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	109	Total	O	0	1
			109	109		
5	B	68	Total	O	0	0
			68	68		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.41Å 60.61Å 117.94Å 90.00° 102.86° 90.00°	Depositor
Resolution (Å)	44.28 – 1.85 44.24 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.28-1.85) 100.0 (44.24-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.8.0405	Depositor
R, $R_{free}$	0.171 , 0.213 0.180 , 0.219	Depositor DCC
$R_{free}$ test set	2778 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.2	Xtrriage
Anisotropy	0.450	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 52.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9061	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, Y5I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.80	4/2205 (0.2%)	1.02	1/2969 (0.0%)
1	B	0.71	3/2195 (0.1%)	0.98	2/2952 (0.1%)
All	All	0.76	7/4400 (0.2%)	1.00	3/5921 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	365	GLU	CD-OE2	-8.70	1.16	1.25
1	B	324	GLU	CD-OE2	-7.19	1.17	1.25
1	A	342	SER	CA-CB	-6.14	1.43	1.52
1	B	378	GLU	CD-OE2	-5.88	1.19	1.25
1	B	427	GLU	CD-OE2	-5.67	1.19	1.25
1	A	418	GLU	CD-OE1	-5.52	1.19	1.25
1	A	428	SER	CA-CB	-5.01	1.45	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212	ARG	NE-CZ-NH2	-6.41	117.10	120.30
1	B	212	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	A	350	ARG	NE-CZ-NH1	-5.53	117.54	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	212	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2151	2233	2230	15	0
1	B	2142	2228	2225	16	0
2	A	30	19	0	0	0
3	A	18	24	24	0	0
3	B	6	8	8	0	0
4	A	20	0	0	2	0
4	B	5	0	0	1	0
5	A	109	0	0	1	0
5	B	68	0	0	3	0
All	All	4549	4512	4487	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:429:SER:OG	4:B:502:SO4:O4	1.98	0.80
1:A:325[B]:ILE:HD11	1:A:392:ILE:HG13	1.65	0.79
1:A:276:GLU:CD	1:A:357:ARG:HE	2.03	0.61
1:A:325[B]:ILE:HD11	1:A:392:ILE:CG1	2.32	0.59
1:B:255:LEU:HD21	1:B:277:VAL:HG13	1.85	0.58
1:A:325[A]:ILE:HD11	1:A:391:ILE:HB	1.85	0.57
1:B:367:LYS:HE3	5:B:657:HOH:O	2.04	0.57
1:A:263:LYS:O	1:A:264:PHE:HB2	2.08	0.53
1:A:466:HIS:ND1	1:A:467:PRO:HD2	2.24	0.52
1:B:357:ARG:HG3	1:B:357:ARG:HH21	1.74	0.52
1:B:276:GLU:HG3	1:B:357:ARG:HE	1.74	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:LEU:CD2	1:B:277:VAL:HG13	2.39	0.52
1:B:325:ILE:HD11	1:B:392:ILE:HG13	1.91	0.51
1:A:205:ASN:HB2	1:A:206:PRO:CD	2.42	0.49
1:B:277:VAL:HA	1:B:280:ARG:NH2	2.28	0.49
1:B:449:HIS:CD2	5:B:657:HOH:O	2.67	0.48
1:B:247:PHE:CZ	1:B:257:MET:HG3	2.49	0.47
1:B:449:HIS:HD2	5:B:657:HOH:O	1.97	0.47
1:A:427:GLU:HB2	4:A:508:SO4:O1	2.14	0.47
1:B:327:TYR:CE1	1:B:367:LYS:HE2	2.50	0.46
1:B:459:THR:C	1:B:460:GLU:HG2	2.35	0.46
1:A:320:TYR:CE1	1:A:476:LEU:HD12	2.52	0.45
1:B:325:ILE:HD11	1:B:392:ILE:CG1	2.46	0.45
1:B:255:LEU:CD2	1:B:277:VAL:CG1	2.95	0.44
1:A:203:GLY:HA3	5:A:697:HOH:O	2.19	0.43
1:B:449:HIS:CE1	1:B:453:LEU:HG	2.54	0.42
1:A:262:ILE:HG22	1:A:264:PHE:H	1.85	0.41
1:A:325[A]:ILE:HD13	1:A:391:ILE:HG21	2.03	0.41
1:A:427:GLU:HB2	4:A:508:SO4:S	2.60	0.41
1:A:204:LEU:O	1:A:204:LEU:HD13	2.21	0.41
1:A:466:HIS:CE1	1:A:467:PRO:HD2	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/275 (97%)	264 (99%)	3 (1%)	0	100	100
1	B	263/275 (96%)	259 (98%)	4 (2%)	0	100	100
All	All	530/550 (96%)	523 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	243/247 (98%)	242 (100%)	1 (0%)	91 89
1	B	242/247 (98%)	236 (98%)	6 (2%)	47 31
All	All	485/494 (98%)	478 (99%)	7 (1%)	65 55

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LEU
1	B	208	SER
1	B	245	SER
1	B	283	GLN
1	B	362	ASP
1	B	461	THR
1	B	463	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	501	-	5,5,5	0.28	0	5,5,5	0.21	0
3	GOL	A	503	-	5,5,5	0.20	0	5,5,5	0.46	0
3	GOL	A	504	-	5,5,5	0.27	0	5,5,5	1.15	0
4	SO4	B	502	-	4,4,4	0.93	0	6,6,6	0.19	0
4	SO4	A	507	-	4,4,4	0.61	0	6,6,6	0.17	0
2	Y5I	A	501	-	29,32,32	0.74	1 (3%)	33,43,43	1.60	4 (12%)
4	SO4	A	505	-	4,4,4	0.27	0	6,6,6	0.43	0
4	SO4	A	506	-	4,4,4	0.50	0	6,6,6	0.49	0
4	SO4	A	508	-	4,4,4	0.68	0	6,6,6	0.50	0
3	GOL	A	502	-	5,5,5	0.25	0	5,5,5	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	501	-	-	2/4/4/4	-
3	GOL	A	503	-	-	4/4/4/4	-
3	GOL	A	504	-	-	0/4/4/4	-
2	Y5I	A	501	-	-	2/17/17/17	0/3/3/3
3	GOL	A	502	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	Y5I	N2-N1	2.16	1.41	1.37

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	Y5I	C1-C2-S1	6.72	129.16	120.12
2	A	501	Y5I	C3-N2-N1	3.93	113.39	105.29
2	A	501	Y5I	O3-C17-C16	-2.17	116.12	123.08
2	A	501	Y5I	O4-C17-C16	2.13	120.87	114.03

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	503	GOL	C1-C2-C3-O3
3	A	503	GOL	O2-C2-C3-O3
3	A	503	GOL	O1-C1-C2-C3
3	B	501	GOL	O1-C1-C2-C3
3	A	503	GOL	O1-C1-C2-O2
3	B	501	GOL	O1-C1-C2-O2
2	A	501	Y5I	C15-C16-C17-O3
2	A	501	Y5I	C15-C16-C17-O4

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	502	SO4	1	0
4	A	508	SO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/275 (96%)	-0.03	8 (3%) 50 48	26, 39, 68, 104	0
1	B	263/275 (95%)	0.72	44 (16%) 1 1	29, 48, 90, 128	0
All	All	528/550 (96%)	0.34	52 (9%) 7 7	26, 43, 82, 128	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	456	ILE	7.9
1	A	264	PHE	7.1
1	B	241	THR	6.7
1	B	242	THR	6.2
1	A	203	GLY	5.7
1	B	458	LYS	5.7
1	A	204	LEU	5.5
1	B	263	LYS	5.0
1	B	463	MET	5.0
1	B	244	LYS	4.8
1	B	457	LYS	4.8
1	B	462	ASP	4.5
1	A	263	LYS	4.5
1	B	360	PHE	4.3
1	B	357	ARG	4.2
1	B	454	GLN	4.1
1	B	461	THR	4.0
1	B	464	SER	3.9
1	B	240	LYS	3.8
1	B	277	VAL	3.7
1	B	439[A]	MET	3.6
1	B	252	MET	3.4
1	B	465	LEU	3.2
1	B	256	MET	3.0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	358	LYS	3.0
1	B	359	PRO	3.0
1	B	356	LEU	2.9
1	B	279	ILE	2.9
1	B	260[A]	ASP	2.7
1	B	328	THR	2.5
1	B	322	VAL	2.5
1	B	460	GLU	2.5
1	A	465	LEU	2.5
1	B	391	ILE	2.5
1	B	361	GLY	2.4
1	B	239	GLY	2.4
1	B	442	LEU	2.3
1	B	354	LYS	2.3
1	A	260	ASP	2.3
1	A	275	LYS	2.3
1	B	327	TYR	2.2
1	B	276	GLU	2.2
1	A	206	PRO	2.2
1	B	459	THR	2.1
1	B	326	ILE	2.1
1	B	392	ILE	2.1
1	B	208	SER	2.1
1	B	268	THR	2.1
1	B	293	VAL	2.1
1	B	325	ILE	2.1
1	B	259	GLU	2.0
1	B	389	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	504	6/6	0.61	0.26	54,68,76,76	2
3	GOL	A	503	6/6	0.85	0.13	51,68,78,84	2
4	SO4	A	508	5/5	0.89	0.23	44,47,62,62	5
3	GOL	B	501	6/6	0.92	0.12	67,69,74,77	2
4	SO4	B	502	5/5	0.93	0.12	32,36,47,51	5
2	Y5I	A	501	30/30	0.94	0.09	38,46,72,74	0
3	GOL	A	502	6/6	0.95	0.09	55,66,79,86	2
4	SO4	A	506	5/5	0.95	0.17	54,66,75,98	0
4	SO4	A	507	5/5	0.97	0.18	48,54,58,64	5
4	SO4	A	505	5/5	0.98	0.15	57,61,65,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.