

Full wwPDB X-ray Structure Validation Report (i)

Sep 17, 2023 – 08:41 AM EDT

PDB ID : 4PF7

Title : Crystal structure of insulin degrading enzyme complexed with inhibitor

Authors: Wang, Y.; Guo, S.

Deposited on : 2014-04-28

Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	989	90%	6% •
1	В	989	89%	7% • •



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Insulin-degrading enzyme.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	955	Total 7812	C 5035	N 1310	O 1445	S 22	0	1	0
1	В	959	Total 7848		N 1317	O 1455	S 22	0	1	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	MET	-	initiating methionine	UNP P14735
A	32	GLY	_	expression tag	UNP P14735
A	33	HIS	_	expression tag	UNP P14735
A	34	HIS	-	expression tag	UNP P14735
A	35	HIS	-	expression tag	UNP P14735
A	36	HIS	-	expression tag	UNP P14735
A	37	HIS	_	expression tag	UNP P14735
A	38	HIS	-	expression tag	UNP P14735
A	39	GLY	-	expression tag	UNP P14735
A	40	ARG	-	expression tag	UNP P14735
A	41	ALA	-	expression tag	UNP P14735
A	110	LEU	CYS	engineered mutation	UNP P14735
A	111	GLN	GLU	engineered mutation	UNP P14735
A	171	SER	CYS	engineered mutation	UNP P14735
A	178	ALA	CYS	engineered mutation	UNP P14735
A	257	VAL	CYS	engineered mutation	UNP P14735
A	414	LEU	CYS	engineered mutation	UNP P14735
A	573	ASN	CYS	engineered mutation	UNP P14735
A	590	SER	CYS	engineered mutation	UNP P14735
A	789	SER	CYS	engineered mutation	UNP P14735
A	812	ALA	CYS	engineered mutation	UNP P14735
A	819	ALA	CYS	engineered mutation	UNP P14735
A	904	SER	CYS	engineered mutation	UNP P14735
A	966	ASN	CYS	engineered mutation	UNP P14735
A	974	ALA	CYS	engineered mutation	UNP P14735




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	31	MET	-	initiating methionine	UNP P14735
В	32	GLY	-	- expression tag	
В	33	HIS	-	expression tag	UNP P14735
В	34	HIS	-	expression tag	UNP P14735
В	35	HIS	-	expression tag	UNP P14735
В	36	HIS	-	expression tag	UNP P14735
В	37	HIS	-	expression tag	UNP P14735
В	38	HIS	-	expression tag	UNP P14735
В	39	GLY	-	expression tag	UNP P14735
В	40	ARG	-	expression tag	UNP P14735
В	41	ALA	-	expression tag	UNP P14735
В	110	LEU	CYS	engineered mutation	UNP P14735
В	111	GLN	GLU	engineered mutation	UNP P14735
В	171	SER	CYS	engineered mutation	UNP P14735
В	178	ALA	CYS	engineered mutation	UNP P14735
В	257	VAL	CYS	engineered mutation	UNP P14735
В	414	LEU	CYS	engineered mutation	UNP P14735
В	573	ASN	CYS	engineered mutation	UNP P14735
В	590	SER	CYS	engineered mutation	UNP P14735
В	789	SER	CYS	engineered mutation	UNP P14735
В	812	ALA	CYS	engineered mutation	UNP P14735
В	819	ALA	CYS	engineered mutation	UNP P14735
В	904	SER	CYS	engineered mutation	UNP P14735
В	966	ASN	CYS	engineered mutation	UNP P14735
В	974	ALA	CYS	engineered mutation	UNP P14735

• Molecule 2 is (2S)-2-amino-N-{(1S)-1-cyclohexyl-2-[(4-methylphenyl)amino]-2-oxoethyl}-4-(methylselanyl)butanamide (three-letter code: 2QW) (formula: $C_{20}H_{31}N_3O_2Se$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Λ	1	Total	С	N	О	Se	0	0
2	A	1	26	20	3	2	1	U	
2	D	1	Total	С	N	О	Se	0	0
	Б	1	26	20	3	2	1	U	U

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	В	1	Total Zn 1 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	289	Total O 289 289	0	0
4	В	289	Total O 289 289	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Insulin-degrading enzyme





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.84Å 115.96Å 124.28Å	Donositor
a, b, c, α , β , γ	90.00° 97.83° 90.00°	Depositor
Resolution (Å)	19.94 - 2.33	Depositor
Resolution (A)	19.91 - 2.33	EDS
% Data completeness	99.7 (19.94-2.33)	Depositor
(in resolution range)	99.7 (19.91-2.33)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$< I/\sigma(I) > 1$	2.22 (at 2.33Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
D D.	0.199 , 0.236	Depositor
R, R_{free}	0.205 , 0.244	DCC
R_{free} test set	1050 reflections $(1.12%)$	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 42.4	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16292	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 2QW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.48	0/8008	0.65	0/10835
1	В	0.47	0/8046	0.65	0/10884
All	All	0.47	0/16054	0.65	0/21719

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7812	0	7741	35	0
1	В	7848	0	7791	28	0
2	A	26	0	26	0	0
2	В	26	0	26	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	289	0	0	1	0
4	В	289	0	0	0	0
All	All	16292	0	15584	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

		Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap(A)
1:A:44:ASN:HB3	1:A:45:PRO:HD3	1.57	0.86
1:B:643:LYS:HA	1:B:744:MET:HE1	1.66	0.77
1:A:648:LYS:O	1:A:652:PHE:HB3	1.88	0.73
1:A:317:PHE:CE2	1:A:476:VAL:HG13	2.31	0.66
1:A:236:ASP:HB3	1:A:239:GLN:HG2	1.81	0.61
1:A:124:GLU:OE2	1:A:181:ARG:NH1	2.31	0.60
1:A:440:ILE:HG22	1:A:449:VAL:HG23	1.85	0.59
1:A:680:GLN:HA	1:A:683:MET:HE3	1.86	0.57
1:A:473:PRO:O	1:A:476:VAL:HG22	2.05	0.56
1:B:298:LEU:HD21	1:B:318:PRO:HG3	1.90	0.54
1:B:425:LYS:HD3	1:B:454:TYR:OH	2.08	0.54
1:B:236:ASP:HB2	1:B:239:GLN:HB3	1.90	0.52
1:A:655:ASP:OD2	1:A:658:ARG:HG2	2.08	0.52
1:A:437:ILE:HG23	1:A:449:VAL:CG2	2.39	0.52
1:A:277:GLU:HB2	4:A:1313:HOH:O	2.10	0.52
1:A:94:ILE:HG13	1:A:248:TYR:HB3	1.92	0.51
1:A:960:ALA:HB3	1:A:963:MET:HB2	1.91	0.51
1:B:557:SER:HB2	1:B:742:MET:HE3	1.92	0.51
1:B:413:GLU:HG2	1:B:531:ILE:HD11	1.93	0.50
1:B:778:VAL:HG22	1:B:955:SER:HB2	1.92	0.50
1:B:960:ALA:HB3	1:B:963:MET:HB2	1.92	0.50
1:A:437:ILE:HG23	1:A:449:VAL:HG21	1.94	0.50
1:B:224:TYR:HA	1:B:228:THR:HB	1.93	0.49
1:A:648:LYS:HG3	1:A:652:PHE:HB2	1.94	0.49
1:A:771:LEU:HB2	1:A:952:HIS:HB3	1.95	0.49
1:B:367:ALA:HB3	1:B:370:PHE:CE2	2.47	0.49
1:B:771:LEU:HB2	1:B:952:HIS:HB3	1.95	0.48
1:A:44:ASN:HB3	1:A:45:PRO:CD	2.37	0.48
1:A:778:VAL:HG22	1:A:955:SER:HB2	1.95	0.48
1:A:425:LYS:HE2	1:A:454:TYR:OH	2.15	0.47
1:B:441:LEU:HD23	1:B:449:VAL:HG11	1.96	0.47
1:A:367:ALA:HB3	1:A:370:PHE:CE2	2.50	0.47
1:B:685:TYR:HB2	1:B:956:VAL:HG11	1.96	0.46
1:A:188:SER:HB3	1:A:831:TYR:HD1	1.80	0.46
1:A:413:GLU:HG2	1:A:531:ILE:HD11	1.98	0.46
1:B:491:ARG:HH11	1:B:500:TYR:HE2	1.61	0.46
1:A:648:LYS:HG3	1:A:652:PHE:CB	2.46	0.45
1:B:887:GLN:O	1:B:891:ILE:HG12	2.17	0.45
1:B:91:ASP:OD2	1:B:146:HIS:ND1	2.45	0.44
1:A:100:PRO:HG2	1:A:103:ILE:HD12	1.99	0.44



Continued from previous page...

A + 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:A:685:TYR:HB2	1:A:956:VAL:HG11	1.99	0.44
1:A:538:LEU:HD13	1:A:734:THR:HG23	1.99	0.44
1:B:538:LEU:HD13	1:B:734:THR:HG23	1.99	0.44
1:A:88:ALA:HB3	1:A:151:PHE:CE2	2.53	0.43
1:A:823:LEU:HB3	1:A:829:LEU:HD12	2.01	0.43
1:B:88:ALA:HB3	1:B:151:PHE:CE2	2.53	0.43
1:B:823:LEU:HB3	1:B:829:LEU:HD12	2.00	0.43
1:B:577:GLU:HG2	1:B:579:PHE:CZ	2.54	0.43
1:B:803:SER:HA	1:B:927:TYR:CE2	2.53	0.43
1:B:44:ASN:HD22	1:B:46:ALA:H	1.66	0.43
1:A:291:HIS:CD2	1:A:370:PHE:HB2	2.53	0.42
1:A:624:ILE:HD12	1:A:766:TYR:HD1	1.84	0.42
1:A:200:ARG:HG2	1:A:496:TYR:HB3	2.02	0.42
1:A:803:SER:HA	1:A:927:TYR:CE2	2.54	0.42
1:B:600:LEU:HD21	1:B:649:MET:HG3	2.01	0.42
1:B:94:ILE:HG13	1:B:248:TYR:HB3	2.01	0.42
1:B:113:MET:HA	1:B:116:LEU:HD12	2.01	0.42
1:A:91:ASP:OD2	1:A:146:HIS:ND1	2.46	0.41
1:B:616:LEU:HD11	1:B:638:GLN:HG3	2.02	0.41
1:A:230:PRO:HB3	1:A:235:ILE:HB	2.02	0.41
1:B:624:ILE:HD12	1:B:766:TYR:HD1	1.84	0.41
1:A:121:TYR:HB3	1:A:126:GLU:HG2	2.01	0.41
1:B:44:ASN:HB3	1:B:47:ILE:HD12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percenti	les
1	A	952/989~(96%)	918 (96%)	33 (4%)	1 (0%)	51 62	2
1	В	956/989 (97%)	924 (97%)	30 (3%)	2 (0%)	47 55	5



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1908/1978 (96%)	1842 (96%)	63 (3%)	3 (0%)	47 55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	175	ASP
1	В	452	ALA
1	В	978	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	849/878 (97%)	837 (99%)	12 (1%)	67 78		
1	В	856/878 (98%)	832 (97%)	24 (3%)	43 53		
All	All	1705/1756 (97%)	1669 (98%)	36 (2%)	53 65		

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	ILE
1	A	181	ARG
1	A	228	THR
1	A	287	GLU
1	A	476	VAL
1	A	518	LEU
1	A	604	LEU
1	A	652	PHE
1	A	676	GLU
1	A	789	SER
1	A	799	MET
1	A	821	ASN
1	В	44	ASN
1	В	83	THR
1	В	118	THR



Continued from previous page...

Mol	Chain	Res	Type
1	В	124	GLU
1	В	125	ASN
1	В	128	SER
1	В	181	ARG
1	В	236	ASP
1	В	238	ARG
1	В	287	GLU
1	В	476	VAL
1	В	508	GLU
1	В	518	LEU
1	В	676	GLU
1	В	742	MET
1	В	744	MET
1	В	751	GLU
1	В	789	SER
1	В	799	MET
1	В	824	ARG
1	В	868	ILE
1	В	978	ILE
1	В	988	GLN
1	В	997	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	ASN
1	A	781	GLN
1	A	813	GLN
1	A	988	GLN
1	В	44	ASN
1	В	231	ASN
1	В	781	GLN
1	В	813	GLN
1	В	988	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		$_{ m n} \mid _{ m Res}$	Res Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2QW	A	1101	-	25,27,27	1.29	4 (16%)	31,35,35	1.03	1 (3%)
2	2QW	В	1101	-	25,27,27	1.38	2 (8%)	31,35,35	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2QW	A	1101	-	-	2/23/32/32	0/2/2/2
2	2QW	В	1101	-	-	3/23/32/32	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$Ideal(\AA)$
2	В	1101	2QW	C4-N1	3.63	1.42	1.34
2	A	1101	2QW	C4-N1	3.60	1.42	1.34
2	В	1101	2QW	C12-N2	2.95	1.42	1.35
2	A	1101	2QW	C14-C13	2.24	1.43	1.39
2	A	1101	2QW	C19-C13	2.22	1.43	1.39
2	A	1101	2QW	C11-C6	2.06	1.58	1.53



All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	1101	2QW	C7-C6-C5	2.14	114.72	111.97

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	В	1101	2QW	C4-C-C1-C2
2	В	1101	2QW	N-C-C1-C2
2	A	1101	2QW	N-C-C1-C2
2	В	1101	2QW	O1-C12-N2-C13
2	A	1101	2QW	C4-C-C1-C2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	955/989~(96%)	0.16	41 (4%) 35 46	24, 43, 72, 93	0
1	В	959/989 (96%)	0.10	28 (2%) 51 61	26, 42, 68, 117	0
All	All	1914/1978 (96%)	0.13	69 (3%) 42 53	24, 43, 71, 117	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1012	ILE	6.3
1	В	979	ASN	5.7
1	В	978	ILE	5.5
1	В	977	ASP	5.3
1	A	517	ASP	5.1
1	A	45	PRO	4.9
1	В	43	ASN	4.9
1	В	518	LEU	4.6
1	В	964	ASP	4.4
1	A	980	LEU	4.3
1	A	44	ASN	4.0
1	A	985	ALA	3.8
1	A	947	ASP	3.8
1	A	982	GLN	3.8
1	A	1011	HIS	3.7
1	A	983	ALA	3.7
1	A	988	GLN	3.7
1	В	980	LEU	3.5
1	В	962	GLU	3.4
1	В	985	ALA	3.4
1	A	857	HIS	3.3
1	A	259	LEU	3.3
1	A	787	ASN	3.2
1	В	235	ILE	3.1



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ	
1	A	817	GLU	3.1	
1	A	258	VAL	3.1	
1	A	518	LEU	3.1	
1	В	45	PRO	3.0	
1	В	1012	ILE	3.0	
1	A	965	SER	2.8	
1	A	876	ASP	2.8	
1	A	791	ILE	2.8	
1	A	119	LYS	2.8	
1	В	123	LYS	2.7	
1	В	44	ASN	2.7	
1	В	983	ALA	2.7	
1	A	831	TYR	2.7	
1	A	125	ASN	2.7	
1	A	774	ARG	2.7	
1	В	129	GLN	2.6	
1	A	261	ARG	2.5	
1	В	517	ASP	2.5	
1	В	728	LEU	2.5	
1	В	236	ASP	2.5	
1	В	507	ASP	2.5	
1	A	962	GLU	2.4	
1	В	965	SER	2.4	
1	A	579[A]	PHE	2.4	
1	A	880	GLU	2.4	
1	В	1011	HIS	2.3	
1	A	46	ALA	2.3	
1	В	114	LEU	2.3	
1	A	964	ASP	2.3	
1	A	989	PRO	2.3	
1	В	774	ARG	2.2	
1	В	1009	LYS	2.2	
1	A	936	ILE	2.2	
1	A	956	VAL	2.2	
1	A	257	VAL	2.1	
1	A	543	GLU	2.1	
1	A	863	VAL	2.1	
1	В	1010	PRO	2.1	
1	A	877	MET	2.1	
1	В	982	GLN	2.1	
1	A	955	SER	2.1	
1	A	867	LEU	2.1	



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	76	LEU	2.1
1	A	109	PHE	2.0
1	A	810	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	2QW	В	1101	26/26	0.88	0.20	30,34,45,53	0
2	2QW	A	1101	26/26	0.89	0.20	32,38,50,56	0
3	ZN	В	1102	1/1	0.97	0.06	86,86,86,86	0
3	ZN	A	1102	1/1	0.98	0.03	72,72,72,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

