



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2023 – 07:39 AM EDT

PDB ID : 4PFJ  
Title : The structure of bi-acetylated SAHH  
Authors : Kavran, J.M.; Wang, Y.; Cole, P.A.; Leahy, D.J.  
Deposited on : 2014-04-29  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	
1	B	432	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13833 atoms, of which 6729 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenosylhomocysteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	430	6677	2118	3338	572	623	26	1	1	0
1	B	427	6626	2101	3313	569	617	26	0	1	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	86	ASN	ASP	variant	UNP P23526
A	396	CYS	GLU	engineered mutation	UNP P23526
B	86	ASN	ASP	variant	UNP P23526
B	396	CYS	GLU	engineered mutation	UNP P23526

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	A	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		
2	B	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

- Molecule 3 is ADENOSINE (three-letter code: ADN) (formula: C<sub>10</sub>H<sub>13</sub>N<sub>5</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
3	A	1	Total	C	H	N	O	0	0
			32	10	13	5	4		
3	B	1	Total	C	H	N	O	0	0
			32	10	13	5	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	194	Total	O	0	0
			194	194		
4	B	132	Total	O	0	0
			132	132		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosylhomocysteinase



- Molecule 1: Adenosylhomocysteinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.37Å 102.73Å 176.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.18 – 2.30 49.18 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.18-2.30) 99.7 (49.18-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.194 , 0.245 0.197 , 0.247	Depositor DCC
$R_{free}$ test set	2008 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.1	Xtrriage
Anisotropy	0.512	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 32.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13833	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADN, NAD, ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/3380	0.44	0/4573
1	B	0.27	0/3354	0.43	0/4538
All	All	0.27	0/6734	0.44	0/9111

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3339	3338	3351	6	0
1	B	3313	3313	3324	22	0
2	A	44	26	26	1	0
2	B	44	26	26	0	0
3	A	19	13	13	1	0
3	B	19	13	13	1	0
4	A	194	0	0	1	0
4	B	132	0	0	5	0
All	All	7104	6729	6753	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:ASN:ND2	1:B:182:ASP:OD2	2.09	0.84
1:B:94:LYS:NZ	4:B:604:HOH:O	2.30	0.64
1:B:168:MET:HE2	1:B:173:LEU:HD23	1.84	0.60
1:B:26:GLU:OE1	4:B:601:HOH:O	2.17	0.60
1:B:301:HIS:ND1	3:B:502:ADN:O5'	2.37	0.58
1:A:233:ARG:NH2	4:A:610:HOH:O	2.38	0.56
1:B:72:ALA:O	4:B:602:HOH:O	2.18	0.55
1:B:431:ARG:O	4:B:603:HOH:O	2.18	0.54
1:A:151:ARG:HD3	1:A:376:PRO:HG3	1.89	0.53
1:B:30:PRO:HA	1:B:401:ALY:HH32	1.89	0.53
1:B:131:ASP:HB3	1:B:156:GLU:HB3	1.92	0.52
1:A:34:ARG:NH2	1:A:401:ALY:OH	2.44	0.50
1:B:131:ASP:CB	1:B:156:GLU:HB3	2.41	0.49
1:A:168:MET:HE2	1:A:173:LEU:HD23	1.94	0.49
1:B:188:LYS:NZ	4:B:603:HOH:O	2.40	0.48
1:B:189:PHE:HA	1:B:193:TYR:CD2	2.49	0.47
1:B:131:ASP:HA	1:B:155:GLU:OE2	2.15	0.46
1:B:185:THR:HG22	1:B:360:ASN:HB3	1.99	0.45
1:B:185:THR:HG21	1:B:394:VAL:CG1	2.47	0.44
1:A:179:ASN:ND2	1:A:182:ASP:OD2	2.38	0.43
1:B:250:LEU:O	1:B:254:MET:HG2	2.18	0.43
1:B:318:LYS:HD3	1:B:328:TYR:CE1	2.53	0.42
2:A:501:NAD:C4N	3:A:502:ADN:H3'	2.49	0.42
1:B:82:PHE:CE1	1:B:104:GLY:HA2	2.55	0.42
1:B:59:GLU:HG3	1:B:355:SER:HA	2.01	0.41
1:A:30:PRO:O	1:A:34:ARG:HG2	2.19	0.41
1:B:147:LEU:N	1:B:148:PRO:CD	2.84	0.41
1:B:408:ALY:HH31	1:B:408:ALY:HE2	1.83	0.41
1:B:22:LEU:HD21	1:B:61:ALA:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/432 (99%)	412 (96%)	15 (4%)	0	100	100
1	B	424/432 (98%)	407 (96%)	16 (4%)	1 (0%)	47	58
All	All	851/864 (98%)	819 (96%)	31 (4%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	381	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/354 (100%)	352 (100%)	1 (0%)	92	97
1	B	350/354 (99%)	348 (99%)	2 (1%)	86	94
All	All	703/708 (99%)	700 (100%)	3 (0%)	91	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	335	ARG
1	B	34	ARG
1	B	87	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	ALY	A	401	1	10,11,12	0.84	0	7,12,14	0.60	0
1	ALY	B	408	1	10,11,12	0.89	0	7,12,14	1.47	1 (14%)
1	ALY	B	401	1	10,11,12	0.86	0	7,12,14	0.78	0
1	ALY	A	408	1	10,11,12	0.82	0	7,12,14	1.22	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	A	401	1	-	4/9/10/12	-
1	ALY	B	408	1	-	4/9/10/12	-
1	ALY	B	401	1	-	1/9/10/12	-
1	ALY	A	408	1	-	4/9/10/12	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	408	ALY	CE-NZ-CH	2.87	126.97	122.56
1	A	408	ALY	CE-NZ-CH	2.26	126.04	122.56

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	401	ALY	N-CA-CB-CG
1	A	401	ALY	C-CA-CB-CG

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	401	ALY	CG-CD-CE-NZ
1	A	408	ALY	OH-CH-NZ-CE
1	A	408	ALY	CH3-CH-NZ-CE
1	B	408	ALY	OH-CH-NZ-CE
1	B	408	ALY	CH3-CH-NZ-CE
1	A	408	ALY	CG-CD-CE-NZ
1	A	408	ALY	CA-CB-CG-CD
1	B	408	ALY	CG-CD-CE-NZ
1	A	401	ALY	CD-CE-NZ-CH
1	B	401	ALY	CA-CB-CG-CD
1	B	408	ALY	CE-CD-CG-CB

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	401	ALY	1	0
1	B	408	ALY	1	0
1	B	401	ALY	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	B	501	-	42,48,48	1.51	3 (7%)	50,73,73	1.61	10 (20%)
3	ADN	A	502	-	18,21,21	0.71	1 (5%)	18,31,31	0.89	0
3	ADN	B	502	-	18,21,21	0.86	1 (5%)	18,31,31	0.92	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	A	501	-	42,48,48	1.64	8 (19%)	50,73,73	1.62	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	501	-	-	8/26/62/62	0/5/5/5
3	ADN	A	502	-	-	2/2/22/22	0/3/3/3
3	ADN	B	502	-	-	2/2/22/22	0/3/3/3
2	NAD	A	501	-	-	6/26/62/62	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAD	PN-O5D	4.91	1.79	1.59
2	A	501	NAD	PN-O5D	4.60	1.77	1.59
2	B	501	NAD	PA-O5B	4.24	1.76	1.59
2	A	501	NAD	PA-O5B	3.85	1.74	1.59
2	A	501	NAD	C2N-N1N	3.28	1.38	1.35
2	B	501	NAD	C7N-N7N	2.93	1.38	1.33
2	A	501	NAD	C7N-N7N	2.92	1.38	1.33
2	A	501	NAD	C2D-C1D	2.65	1.57	1.53
3	B	502	ADN	C2-N3	2.46	1.36	1.32
2	A	501	NAD	C2A-N1A	2.15	1.37	1.33
2	A	501	NAD	C2A-N3A	2.14	1.35	1.32
2	A	501	NAD	O2B-C2B	-2.08	1.38	1.43
3	A	502	ADN	C2-N3	2.02	1.35	1.32

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAD	PN-O3-PA	-5.33	114.54	132.83
2	A	501	NAD	PN-O3-PA	-5.25	114.80	132.83
2	B	501	NAD	O2A-PA-O1A	3.52	129.66	112.24
2	A	501	NAD	O2A-PA-O1A	3.46	129.33	112.24
2	A	501	NAD	O2N-PN-O1N	3.44	129.24	112.24
2	B	501	NAD	O2N-PN-O1N	3.29	128.52	112.24
2	A	501	NAD	O4B-C1B-C2B	-2.63	103.08	106.93
2	A	501	NAD	O5D-PN-O1N	-2.61	98.86	109.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAD	C5B-C4B-C3B	-2.61	105.40	115.18
2	B	501	NAD	O5D-PN-O1N	-2.55	99.12	109.07
2	B	501	NAD	PN-O5D-C5D	-2.47	107.22	121.68
2	A	501	NAD	PN-O5D-C5D	-2.40	107.58	121.68
2	B	501	NAD	C5B-C4B-C3B	-2.35	106.39	115.18
2	A	501	NAD	O2N-PN-O5D	-2.33	96.93	107.75
2	B	501	NAD	C1B-N9A-C4A	-2.27	122.64	126.64
2	A	501	NAD	C1B-N9A-C4A	-2.17	122.83	126.64
3	B	502	ADN	C1'-N9-C4	-2.11	122.94	126.64
2	B	501	NAD	O2N-PN-O5D	-2.10	98.01	107.75
2	B	501	NAD	C2N-N1N-C1D	-2.08	114.51	119.14
2	B	501	NAD	C3D-C2D-C1D	-2.03	97.92	100.98

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NAD	O4D-C1D-N1N-C2N
2	A	501	NAD	O4D-C1D-N1N-C6N
2	A	501	NAD	C2D-C1D-N1N-C2N
2	A	501	NAD	C2D-C1D-N1N-C6N
2	B	501	NAD	O4D-C1D-N1N-C2N
2	B	501	NAD	O4D-C1D-N1N-C6N
2	B	501	NAD	C2D-C1D-N1N-C2N
2	B	501	NAD	C2D-C1D-N1N-C6N
3	A	502	ADN	O4'-C4'-C5'-O5'
3	B	502	ADN	O4'-C4'-C5'-O5'
3	A	502	ADN	C3'-C4'-C5'-O5'
3	B	502	ADN	C3'-C4'-C5'-O5'
2	B	501	NAD	O4B-C4B-C5B-O5B
2	B	501	NAD	C3B-C4B-C5B-O5B
2	B	501	NAD	O4D-C4D-C5D-O5D
2	A	501	NAD	O4B-C4B-C5B-O5B
2	A	501	NAD	C5D-O5D-PN-O1N
2	B	501	NAD	C5D-O5D-PN-O1N

There are no ring outliers.

3 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	ADN	1	0



Continued on next page...


*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	ADN	1	0
2	A	501	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	428/432 (99%)	0.09	6 (1%) 75 80	27, 35, 45, 61	0
1	B	425/432 (98%)	0.93	74 (17%) 1 1	27, 44, 74, 204	0
All	All	853/864 (98%)	0.51	80 (9%) 8 11	27, 37, 71, 204	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	118	LEU	10.6
1	B	175	VAL	8.1
1	B	55	HIS	7.0
1	B	125	LEU	6.2
1	B	63	LEU	5.9
1	B	68	VAL	5.5
1	B	86	ASN	5.1
1	B	111	LEU	5.0
1	B	54	LEU	4.8
1	B	153	ILE	4.8
1	B	82	PHE	4.5
1	B	377	ASP	4.5
1	B	39	TYR	4.4
1	B	119	TYR	4.3
1	B	379	TYR	4.1
1	B	135	LEU	4.1
1	B	124	PRO	4.1
1	B	45	LEU	4.1
1	B	110	TYR	4.0
1	B	67	LEU	4.0
1	B	302	PHE	3.9
1	B	170	ASN	3.8
1	B	123	GLY	3.8
1	B	121	LYS	3.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	64	ILE	3.7
1	B	142	LYS	3.7
1	B	143	TYR	3.6
1	B	114	ILE	3.5
1	B	131	ASP	3.5
1	B	165	TYR	3.5
1	B	141	THR	3.3
1	B	147	LEU	3.3
1	B	70	LEU	3.3
1	B	150	ILE	3.3
1	B	41	ALA	3.2
1	B	101	ALA	3.2
1	B	301	HIS	3.2
1	B	366	VAL	3.2
1	B	84	THR	3.2
1	B	35	MET	3.2
1	B	129	LEU	3.2
1	A	229	ALA	3.1
1	B	149	GLY	3.1
1	B	378	LYS	3.1
1	A	231	ALA	3.0
1	B	81	ILE	2.9
1	B	415	GLN	2.9
1	B	102	TRP	2.7
1	A	234	GLY	2.7
1	B	166	LYS	2.7
1	B	189	PHE	2.7
1	B	115	GLU	2.6
1	B	44	PRO	2.5
1	B	83	SER	2.5
1	B	237	ALA	2.5
1	B	99	VAL	2.4
1	B	92	ILE	2.4
1	B	172	ILE	2.4
1	B	79	CYS	2.4
1	B	97	ILE	2.4
1	A	232	LEU	2.3
1	B	47	GLY	2.3
1	B	368	ALA	2.2
1	B	7	TYR	2.2
1	A	237	ALA	2.2
1	B	256	GLY	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	10	ALA	2.2
1	B	157	THR	2.2
1	B	46	LYS	2.2
1	B	37	GLU	2.2
1	B	139	ILE	2.1
1	B	36	ARG	2.1
1	B	43	LYS	2.1
1	B	48	ALA	2.1
1	B	236	GLY	2.1
1	A	228	CYS	2.1
1	B	112	TRP	2.1
1	B	188	LYS	2.1
1	B	8	LYS	2.0
1	B	253	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	ALY	B	401	12/13	0.81	0.26	36,48,55,55	0
1	ALY	A	401	12/13	0.82	0.25	33,43,53,54	0
1	ALY	B	408	12/13	0.90	0.24	32,42,51,52	0
1	ALY	A	408	12/13	0.95	0.12	33,39,47,47	9

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ADN	B	502	19/19	0.77	0.33	50,56,67,70	0
3	ADN	A	502	19/19	0.89	0.21	33,39,47,50	0
2	NAD	B	501	44/44	0.96	0.13	27,37,54,56	0
2	NAD	A	501	44/44	0.97	0.14	28,34,41,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around ADN A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.