



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 30, 2023 – 06:25 pm GMT

PDB ID : 8PFH  
Title : Crystal structure of the yeast septin complex Shs1-Cdc12-Cdc3-Cdc10  
Authors : Grupp, B.; Denkhaus, L.; Gerhardt, S.; Gronemeyer, T.  
Deposited on : 2023-06-16  
Resolution : 3.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| $R_{free}$            | 130704                      | 1619 (3.28-3.20)                                      |
| Clashscore            | 141614                      | 1755 (3.28-3.20)                                      |
| Ramachandran outliers | 138981                      | 1728 (3.28-3.20)                                      |
| Sidechain outliers    | 138945                      | 1727 (3.28-3.20)                                      |
| RSRZ outliers         | 127900                      | 1567 (3.28-3.20)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 300    | <br>78% 13% 9%   |
| 2   | B     | 332    | <br>73% 14% 14%  |
| 3   | C     | 330    | <br>75% 12% 12%  |
| 4   | D     | 320    | <br>79% 12% 9%   |

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 17801 atoms, of which 8715 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CDC10 isoform 1.

| Mol | Chain | Residues | Atoms |      |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C    | H    | N   | O   | S |         |         |       |
| 1   | A     | 272      | 4308  | 1380 | 2129 | 375 | 421 | 3 | 0       | 0       | 0     |

There are 7 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment               | Reference      |
|-------|---------|----------|--------|-----------------------|----------------|
| A     | 23      | MET      | -      | initiating methionine | UNP A0A8H4BSX4 |
| A     | 24      | ALA      | -      | expression tag        | UNP A0A8H4BSX4 |
| A     | 25      | ASP      | -      | expression tag        | UNP A0A8H4BSX4 |
| A     | 26      | LEU      | -      | expression tag        | UNP A0A8H4BSX4 |
| A     | 27      | ASN      | -      | expression tag        | UNP A0A8H4BSX4 |
| A     | 28      | TRP      | -      | expression tag        | UNP A0A8H4BSX4 |
| A     | 29      | MET      | -      | expression tag        | UNP A0A8H4BSX4 |

- Molecule 2 is a protein called Cell division control protein 3.

| Mol | Chain | Residues | Atoms |      |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C    | H    | N   | O   | S |         |         |       |
| 2   | B     | 287      | 4416  | 1437 | 2156 | 380 | 436 | 7 | 0       | 0       | 0     |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment               | Reference  |
|-------|---------|----------|--------|-----------------------|------------|
| B     | 79      | MET      | -      | initiating methionine | UNP P32457 |
| B     | 80      | LEU      | -      | expression tag        | UNP P32457 |

- Molecule 3 is a protein called CDC12 isoform 1.

| Mol | Chain | Residues | Atoms |      |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C    | H    | N   | O   | S |         |         |       |
| 3   | C     | 289      | 4491  | 1441 | 2220 | 402 | 421 | 7 | 0       | 0       | 0     |

There are 16 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment               | Reference      |
|-------|---------|----------|--------|-----------------------|----------------|
| C     | -15     | MET      | -      | initiating methionine | UNP A0A6A5PZQ6 |
| C     | -14     | GLY      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -13     | SER      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -12     | SER      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -11     | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -10     | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -9      | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -8      | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -7      | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -6      | HIS      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -5      | SER      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -4      | GLN      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -3      | ASP      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -2      | PRO      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | -1      | ASN      | -      | expression tag        | UNP A0A6A5PZQ6 |
| C     | 0       | SER      | -      | expression tag        | UNP A0A6A5PZQ6 |


- Molecule 4 is a protein called SHS1 isoform 1.

| Mol | Chain | Residues | Atoms |      |      |     |     | ZeroOcc | AltConf | Trace |   |
|-----|-------|----------|-------|------|------|-----|-----|---------|---------|-------|---|
|     |       |          | Total | C    | H    | N   | O   |         |         |       | S |
| 4   | D     | 291      | 4426  | 1436 | 2162 | 366 | 454 | 8       | 0       | 0     | 0 |

There is a discrepancy between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment               | Reference      |
|-------|---------|----------|--------|-----------------------|----------------|
| D     | 20      | MET      | -      | initiating methionine | UNP A0A8H8UM78 |

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms |    |    |   |    | ZeroOcc | AltConf |   |
|-----|-------|----------|-------|----|----|---|----|---------|---------|---|
|     |       |          | Total | C  | H  | N | O  |         |         | P |
| 5   | A     | 1        | Total | C  | H  | N | O  | P       | 0       | 0 |
|     |       |          | 40    | 10 | 12 | 5 | 11 | 2       |         |   |
| 5   | B     | 1        | Total | C  | H  | N | O  | P       | 0       | 0 |
|     |       |          | 40    | 10 | 12 | 5 | 11 | 2       |         |   |
| 5   | C     | 1        | Total | C  | H  | N | O  | P       | 0       | 0 |
|     |       |          | 40    | 10 | 12 | 5 | 11 | 2       |         |   |
| 5   | D     | 1        | Total | C  | H  | N | O  | P       | 0       | 0 |
|     |       |          | 40    | 10 | 12 | 5 | 11 | 2       |         |   |


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CDC10 isoform 1




- Molecule 2: Cell division control protein 3




- Molecule 3: CDC12 isoform 1



- Molecule 4: SHS1 isoform 1

Chain D:  79% 12% 9%



## 4 Data and refinement statistics i

| Property  | Value   | Source           |
|---|---|------------------|
| Space group   | C 1 2 1   | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 258.18Å 69.37Å 92.53Å<br>90.00° 109.68° 90.00°              | Depositor        |
| Resolution (Å)  | 121.55 – 3.24<br>121.55 – 3.24                              | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 63.2 (121.55-3.24)<br>55.5 (121.55-3.24)                    | Depositor<br>EDS |
| $R_{merge}$   | (Not available)   | Depositor        |
| $R_{sym}$   | (Not available)   | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 1.64 (at 3.26Å)   | Xtrriage         |
| Refinement program  | PHENIX 1.20.1_4487  | Depositor        |
| R, $R_{free}$   | 0.282 , 0.290<br>0.285 , 0.294                              | Depositor<br>DCC |
| $R_{free}$ test set   | 779 reflections (4.96%)                                     | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 30.0  | Xtrriage         |
| Anisotropy  | 0.222   | Xtrriage         |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.38 , 24.8   | EDS              |
| L-test for twinning <sup>2</sup>  | $\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$ | Xtrriage         |
| Estimated twinning fraction   | 0.026 for -h-2*1,-k,l                                       | Xtrriage         |
| $F_o, F_c$ correlation  | 0.78  | EDS              |
| Total number of atoms   | 17801   | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 45.0  | wwPDB-VP         |

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.66 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.2733e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |         | Bond angles |         |
|-----|-------|--------------|---------|-------------|---------|
|     |       | RMSZ         | # Z  >5 | RMSZ        | # Z  >5 |
| 1   | A     | 0.26         | 0/2216  | 0.50        | 0/3005  |
| 2   | B     | 0.25         | 0/2305  | 0.44        | 0/3125  |
| 3   | C     | 0.28         | 0/2315  | 0.50        | 0/3144  |
| 4   | D     | 0.25         | 0/2305  | 0.45        | 0/3129  |
| All | All   | 0.26         | 0/9141  | 0.47        | 0/12403 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 2179  | 2129     | 2128     | 36      | 0            |
| 2   | B     | 2260  | 2156     | 2155     | 45      | 1            |
| 3   | C     | 2271  | 2220     | 2217     | 29      | 1            |
| 4   | D     | 2264  | 2162     | 2159     | 23      | 1            |
| 5   | A     | 28    | 12       | 12       | 2       | 0            |
| 5   | B     | 28    | 12       | 12       | 0       | 0            |
| 5   | C     | 28    | 12       | 12       | 0       | 0            |
| 5   | D     | 28    | 12       | 12       | 0       | 0            |
| All | All   | 9086  | 8715     | 8707     | 115     | 2            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1           | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 2:B:223:ILE:HD11 | 2:B:275:VAL:HG22 | 1.61                     | 0.82              |
| 1:A:118:ILE:HG12 | 1:A:143:VAL:HG21 | 1.74                     | 0.68              |
| 3:C:267:TRP:CE2  | 4:D:295:ILE:HD11 | 2.34                     | 0.62              |
| 2:B:342:VAL:HG11 | 2:B:380:LEU:HD22 | 1.79                     | 0.62              |
| 2:B:135:LEU:O    | 2:B:139:LEU:HD23 | 2.00                     | 0.61              |

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1          | Atom-2                 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------------|--------------------------|-------------------|
| 3:C:228:SER:OG  | 3:C:232:GLU:OE1[2_556] | 2.11                     | 0.09              |
| 2:B:356:GLN:OE1 | 4:D:61:SER:OG[2_566]   | 2.17                     | 0.03              |

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Favoured   | Allowed | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1   | A     | 268/300 (89%)   | 259 (97%)  | 9 (3%)  | 0        | 100         | 100 |
| 2   | B     | 283/332 (85%)   | 277 (98%)  | 6 (2%)  | 0        | 100         | 100 |
| 3   | C     | 281/330 (85%)   | 272 (97%)  | 9 (3%)  | 0        | 100         | 100 |
| 4   | D     | 283/320 (88%)   | 277 (98%)  | 6 (2%)  | 0        | 100         | 100 |
| All | All   | 1115/1282 (87%) | 1085 (97%) | 30 (3%) | 0        | 100         | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed       | Rotameric  | Outliers | Percentiles |     |
|-----|-------|----------------|------------|----------|-------------|-----|
| 1   | A     | 237/271 (88%)  | 237 (100%) | 0        | 100         | 100 |
| 2   | B     | 246/306 (80%)  | 246 (100%) | 0        | 100         | 100 |
| 3   | C     | 240/289 (83%)  | 240 (100%) | 0        | 100         | 100 |
| 4   | D     | 249/291 (86%)  | 249 (100%) | 0        | 100         | 100 |
| All | All   | 972/1157 (84%) | 972 (100%) | 0        | 100         | 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3   | C     | 23  | GLN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths |      |          | Bond angles |      |          |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
|     |      |       |     |      | Counts       | RMSZ | # Z  > 2 | Counts      | RMSZ | # Z  > 2 |
| 5   | GDP  | B     | 501 | -    | 24,30,30     | 0.86 | 1 (4%)   | 30,47,47    | 0.99 | 2 (6%)   |
| 5   | GDP  | C     | 401 | -    | 24,30,30     | 0.83 | 0        | 30,47,47    | 0.79 | 1 (3%)   |
| 5   | GDP  | D     | 401 | -    | 24,30,30     | 0.86 | 0        | 30,47,47    | 0.86 | 1 (3%)   |
| 5   | GDP  | A     | 401 | -    | 24,30,30     | 0.85 | 1 (4%)   | 30,47,47    | 0.96 | 2 (6%)   |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions   | Rings   |
|-----|------|-------|-----|------|---------|------------|---------|
| 5   | GDP  | B     | 501 | -    | -       | 1/12/32/32 | 0/3/3/3 |
| 5   | GDP  | C     | 401 | -    | -       | 2/12/32/32 | 0/3/3/3 |
| 5   | GDP  | D     | 401 | -    | -       | 4/12/32/32 | 0/3/3/3 |
| 5   | GDP  | A     | 401 | -    | -       | 2/12/32/32 | 0/3/3/3 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z     | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 5   | B     | 501 | GDP  | C5-C6 | -2.14 | 1.43        | 1.47     |
| 5   | A     | 401 | GDP  | C5-C6 | -2.12 | 1.43        | 1.47     |

The worst 5 of 6 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms       | Z    | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|------|-------------|----------|
| 5   | B     | 501 | GDP  | PA-O3A-PB   | 3.38 | 144.41      | 132.83   |
| 5   | D     | 401 | GDP  | PA-O3A-PB   | 3.26 | 144.02      | 132.83   |
| 5   | A     | 401 | GDP  | PA-O3A-PB   | 2.99 | 143.10      | 132.83   |
| 5   | B     | 501 | GDP  | C3'-C2'-C1' | 2.46 | 104.67      | 100.98   |
| 5   | A     | 401 | GDP  | C3'-C2'-C1' | 2.36 | 104.53      | 100.98   |

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


| Mol | Chain | Res | Type | Atoms          |
|-----|-------|-----|------|----------------|
| 5   | B     | 501 | GDP  | PA-O3A-PB-O2B  |
| 5   | C     | 401 | GDP  | PA-O3A-PB-O2B  |
| 5   | C     | 401 | GDP  | PA-O3A-PB-O3B  |
| 5   | D     | 401 | GDP  | C5'-O5'-PA-O2A |
| 5   | D     | 401 | GDP  | C5'-O5'-PA-O3A |


There are no ring outliers.


1 monomer is involved in 2 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 5   | A     | 401 | GDP  | 2       | 0            |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed        | <RSRZ> | #RSRZ>2   | OWAB(Å <sup>2</sup> ) | Q<0.9 |
|-----|-------|-----------------|--------|-----------|-----------------------|-------|
| 1   | A     | 272/300 (90%)   | -0.17  | 0 100 100 | 37, 48, 65, 76        | 0     |
| 2   | B     | 287/332 (86%)   | -0.20  | 0 100 100 | 36, 48, 65, 79        | 0     |
| 3   | C     | 289/330 (87%)   | -0.13  | 0 100 100 | 36, 47, 64, 80        | 0     |
| 4   | D     | 291/320 (90%)   | -0.13  | 0 100 100 | 36, 47, 63, 75        | 0     |
| All | All   | 1139/1282 (88%) | -0.16  | 0 100 100 | 36, 48, 65, 80        | 0     |

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR  | B-factors(Å <sup>2</sup> ) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 5   | GDP  | C     | 401 | 28/28 | 0.95 | 0.22 | 32,40,46,47                | 0     |
| 5   | GDP  | B     | 501 | 28/28 | 0.96 | 0.19 | 37,39,45,47                | 0     |
| 5   | GDP  | A     | 401 | 28/28 | 0.97 | 0.18 | 39,43,46,48                | 0     |
| 5   | GDP  | D     | 401 | 28/28 | 0.97 | 0.21 | 38,41,46,46                | 0     |






The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around GDP B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GDP A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.