



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2023 – 03:49 PM JST

PDB ID : 4PH6
Title : Structure of 3-Dehydroquinate Dehydratase from *Enterococcus faecalis*
Authors : Xue, B.; Cheung, V.W.; Yew, W.S.; Robinson, R.C.
Deposited on : 2014-05-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-dehydroquinase dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	1873	1190	312	358	13	0	0	0
1	B	240	1873	1190	312	358	13	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P36923
A	-1	SER	-	expression tag	UNP P36923
A	0	HIS	-	expression tag	UNP P36923
B	-2	GLY	-	expression tag	UNP P36923
B	-1	SER	-	expression tag	UNP P36923
B	0	HIS	-	expression tag	UNP P36923


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	162	Total	O	0	0
			162	162		
2	B	149	Total	O	0	0
			149	149		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-dehydroquinate dehydratase

Chain A:  80% 13% 6%



- Molecule 1: 3-dehydroquinate dehydratase

Chain B:  81% 12% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.97Å 48.29Å 67.43Å 108.07° 92.86° 111.83°	Depositor
Resolution (Å)	19.79 – 2.20 19.79 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.2 (19.79-2.20) 89.3 (19.79-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.30 (at 2.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.185 , 0.255 0.199 , 0.264	Depositor DCC
R_{free} test set	1064 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	11.0	Xtriage
Anisotropy	0.787	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4057	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/1902	0.54	0/2575
1	B	0.40	0/1902	0.56	2/2575 (0.1%)
All	All	0.43	0/3804	0.55	2/5150 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	44	LEU	CA-CB-CG	6.51	130.28	115.30
1	B	110	LEU	CA-CB-CG	6.33	129.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1873	0	1894	25	0
1	B	1873	0	1894	20	0
2	A	162	0	0	4	0
2	B	149	0	0	5	0
All	All	4057	0	3788	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:CYS:HB3	1:A:239:VAL:HG11	1.77	0.63
1:A:18:ILE:HG21	1:A:239:VAL:HG13	1.80	0.61
1:A:86:GLU:HB3	1:A:143:HIS:HB2	1.86	0.56
1:B:37:SER:HA	1:B:40:LEU:HD12	1.86	0.56
1:B:79:LEU:HD12	1:B:110:LEU:HG	1.88	0.55
1:A:122:LEU:HD13	1:A:122:LEU:C	2.26	0.55
1:A:215:THR:HG22	1:B:211:ILE:HG21	1.88	0.55
1:B:43:ASP:OD2	1:B:243:ARG:NH1	2.35	0.54
1:B:154:ILE:HG23	1:B:171:ILE:HD13	1.89	0.54
1:A:154:ILE:HG23	1:A:171:ILE:HD13	1.90	0.52
1:B:86:GLU:HB3	1:B:143:HIS:HB2	1.92	0.52
1:B:61:VAL:HG12	1:B:104:LEU:HD21	1.93	0.51
1:A:133:LYS:NZ	2:A:304:HOH:O	2.44	0.50
1:B:52:TYR:CZ	1:B:54:ASN:HB2	2.45	0.50
1:A:52:TYR:CE2	1:A:54:ASN:HB2	2.48	0.49
1:B:122:LEU:HD12	2:B:307:HOH:O	2.11	0.49
1:A:211:ILE:HG21	1:B:215:THR:HG22	1.95	0.48
1:A:61:VAL:HG12	1:A:104:LEU:HD21	1.97	0.47
1:A:170:LYS:HA	1:A:201:VAL:O	2.14	0.47
1:B:115:ILE:O	1:B:140:LEU:HA	2.14	0.47
1:A:82:ARG:CZ	1:A:143:HIS:CE1	2.98	0.47
1:B:74:GLN:HA	2:B:443:HOH:O	2.14	0.46
1:A:52:TYR:CZ	1:A:54:ASN:HB2	2.51	0.46
1:A:79:LEU:HD12	1:A:110:LEU:HD22	1.98	0.45
1:B:148:THR:HG23	1:B:173:VAL:HG21	1.99	0.45
1:B:128:ILE:HG23	1:B:138:ILE:HG13	1.99	0.45
1:A:211:ILE:HG12	2:A:418:HOH:O	2.16	0.44
1:A:66:GLN:NE2	1:A:109:ALA:HB2	2.33	0.44
1:B:116:GLU:HG3	2:B:349:HOH:O	2.17	0.43
1:B:243:ARG:O	1:B:247:LYS:HG3	2.18	0.43
1:A:122:LEU:HD13	1:A:122:LEU:O	2.18	0.43
1:A:122:LEU:C	1:A:122:LEU:CD1	2.87	0.43
1:A:152:GLU:HG3	2:A:323:HOH:O	2.18	0.42
1:A:213:ARG:HD3	2:A:434:HOH:O	2.20	0.42
1:B:120:ASN:HA	1:B:121:PRO:HD3	1.81	0.42
1:A:141:CYS:HA	1:A:170:LYS:O	2.20	0.42
1:A:124:ALA:O	1:A:128:ILE:HG13	2.20	0.41
1:A:115:ILE:O	1:A:140:LEU:HA	2.21	0.41
1:B:170:LYS:HA	1:B:201:VAL:O	2.20	0.41
1:B:23:VAL:HG22	2:B:355:HOH:O	2.20	0.41
1:A:120:ASN:HA	1:A:121:PRO:HD3	1.69	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:LEU:HD11	1:A:136:ILE:HD13	2.04	0.40
1:B:106:LYS:HE3	2:B:379:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/256 (92%)	233 (99%)	3 (1%)	0	100	100
1	B	236/256 (92%)	234 (99%)	2 (1%)	0	100	100
All	All	472/512 (92%)	467 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/218 (95%)	201 (97%)	6 (3%)	42	54
1	B	207/218 (95%)	205 (99%)	2 (1%)	76	86
All	All	414/436 (95%)	406 (98%)	8 (2%)	57	71

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	LYS
1	A	92	PHE
1	A	110	LEU
1	A	122	LEU
1	A	203	MET
1	A	247	LYS
1	B	44	LEU
1	B	110	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/256 (93%)	-0.14	0 100 100	3, 14, 27, 35	0
1	B	240/256 (93%)	-0.11	1 (0%) 92 91	6, 14, 26, 39	0
All	All	480/512 (93%)	-0.12	1 (0%) 95 94	3, 14, 26, 39	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	240	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.