



## wwPDB EM Validation Summary Report ⓘ

Jan 11, 2024 – 07:36 pm GMT

PDB ID : 7PNT  
EMDB ID : EMD-13551  
Title : Assembly intermediate of mouse mitochondrial ribosome small subunit without mS37 in complex with RbfA and Tfb1m  
Authors : Itoh, Y.; Khawaja, A.; Laptev, I.; Sergiev, P.; Rorbach, J.; Amunts, A.  
Deposited on : 2021-09-08  
Resolution : 3.19 Å(reported)  
Based on initial model : 6RW4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	956	 7% 76% 18% 5%
2	B	291	 72% 23%
3	C	167	 75% 21%
4	D	432	 8% 74% 22%
5	E	125	 93% 5% 5%
6	F	242	 8% 82% 14%
7	G	390	 7% 79% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	H	160	
9	I	191	
10	J	139	
11	K	128	
12	L	258	
13	M	135	
14	N	120	
15	O	254	
16	P	143	
17	Q	86	
18	R	359	
19	S	177	
20	T	171	
21	U	200	
22	V	415	
23	W	186	
24	X	391	
25	Y	384	
26	Z	106	
27	0	218	
28	1	320	
29	3	200	
30	4	685	
31	a	350	
32	c	345	

## 2 Entry composition [i](#)

There are 38 unique types of molecules in this entry. The entry contains 128436 atoms, of which 59523 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 12S mitochondrial rRNA.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
1	A	909	29056	8669	9755	3459	6264	909	0	0

- Molecule 2 is a protein called 28S ribosomal protein S2, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	B	223	3590	1142	1799	326	315	8	0	0

- Molecule 3 is a protein called 28S ribosomal protein S24, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	C	132	2163	690	1091	197	180	5	0	0

- Molecule 4 is a protein called 28S ribosomal protein S5, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	D	337	5414	1686	2732	513	473	10	0	0

- Molecule 5 is a protein called 28S ribosomal protein S6, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
5	E	122	1986	617	1007	181	178	3	0	0

- Molecule 6 is a protein called 28S ribosomal protein S7, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
6	F	208	3472	1096	1750	316	299	11	0	0

- Molecule 7 is a protein called 28S ribosomal protein S9, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
7	G	325	5304	1689	2630	480	491	14	0	0

- Molecule 8 is a protein called 28S ribosomal protein S10, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
8	H	140	2346	742	1193	200	207	4	0	0

- Molecule 9 is a protein called 28S ribosomal protein S11, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
9	I	137	2047	629	1038	191	184	5	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	181	5F0	ASN	conflict	UNP Q9DCA2

- Molecule 10 is a protein called 28S ribosomal protein S12, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
10	J	108	1749	528	903	172	141	5	0	0

- Molecule 11 is a protein called 28S ribosomal protein S14, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
11	K	101	1743	534	888	175	140	6	0	0

- Molecule 12 is a protein called 28S ribosomal protein S15, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
12	L	176	3041	930	1576	274	255	6	0	0

- Molecule 13 is a protein called 28S ribosomal protein S16, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
13	M	126	2004	623	1009	194	172	6	0	0

- Molecule 14 is a protein called 28S ribosomal protein S17, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
14	N	113	1839	575	951	160	150	3	0	0

- Molecule 15 is a protein called 28S ribosomal protein S18b, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
15	O	197	3146	1014	1548	289	286	9	0	0

- Molecule 16 is a protein called 28S ribosomal protein S18c, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
16	P	97	1610	505	818	140	139	8	0	0

- Molecule 17 is a protein called 28S ribosomal protein S21, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
17	Q	86	1482	453	750	146	126	7	0	0

- Molecule 18 is a protein called 28S ribosomal protein S22, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
18	R	294	4816	1526	2416	418	449	7	0	0

- Molecule 19 is a protein called 28S ribosomal protein S23, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
19	S	136	2257	722	1133	199	201	2	0	0

- Molecule 20 is a protein called 28S ribosomal protein S25, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
20	T	170	2801	892	1413	238	246	12	0	0

- Molecule 21 is a protein called 28S ribosomal protein S26, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
21	U	174	2908	894	1459	283	270	2	0	0

- Molecule 22 is a protein called 28S ribosomal protein S27, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
22	V	365	5970	1911	2972	506	570	11	0	0

- Molecule 23 is a protein called 28S ribosomal protein S28, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
23	W	100	1606	503	813	141	146	3	0	0

- Molecule 24 is a protein called 28S ribosomal protein S29, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
24	X	357	5762	1834	2881	515	522	10	0	0

- Molecule 25 is a protein called 28S ribosomal protein S31, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
25	Y	149	2439	809	1193	201	233	3	0	0

- Molecule 26 is a protein called 28S ribosomal protein S33, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
26	Z	101	1682	526	848	157	148	3	0	0

- Molecule 27 is a protein called 28S ribosomal protein S34, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
27	0	216	3649	1139	1838	355	313	4	0	0

- Molecule 28 is a protein called 28S ribosomal protein S35, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
28	1	278	4464	1403	2241	384	424	12	0	0

- Molecule 29 is a protein called Aurora kinase A-interacting protein.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
29	3	50	929	286	489	86	68		0	0

- Molecule 30 is a protein called Pentatricopeptide repeat domain-containing protein 3, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
30	4	589	9558	3064	4800	800	872	22	0	0

- Molecule 31 is a protein called Putative ribosome-binding factor A, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
31	a	134	2169	680	1099	190	197	3	0	0

- Molecule 32 is a protein called Dimethyladenosine transferase 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
32	c	299	4871	1543	2478	419	421	10	0	0

- Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
33	A	32	Total	Mg	0
			32	32	
33	B	1	Total	Mg	0
			1	1	

*Continued on next page...*



Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
33	X	1	Total	Mg	0
			1	1	


- Molecule 34 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		AltConf
34	A	7	Total	K	0
			7	7	

- Molecule 35 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
35	O	1	Total	Zn	0
			1	1	

- Molecule 36 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
36	P	1	Total	Fe	S	0
			4	2	2	
36	T	1	Total	Fe	S	0
			4	2	2	

- Molecule 37 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
37	X	1	43	10	12	5	13	3	0

- Molecule 38 is water.

Mol	Chain	Residues	Atoms		AltConf
38	A	268	Total	O	0
			268	268	
38	B	7	Total	O	0
			7	7	
38	C	23	Total	O	0
			23	23	
38	D	14	Total	O	0
			14	14	
38	F	5	Total	O	0
			5	5	
38	G	9	Total	O	0
			9	9	
38	H	11	Total	O	0
			11	11	
38	I	3	Total	O	0
			3	3	
38	J	4	Total	O	0
			4	4	
38	K	15	Total	O	0
			15	15	
38	L	1	Total	O	0
			1	1	

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		AltConf
38	M	17	Total 17	O 17	0
38	N	2	Total 2	O 2	0
38	O	20	Total 20	O 20	0
38	P	6	Total 6	O 6	0
38	Q	2	Total 2	O 2	0
38	R	5	Total 5	O 5	0
38	S	2	Total 2	O 2	0
38	T	6	Total 6	O 6	0
38	U	3	Total 3	O 3	0
38	X	4	Total 4	O 4	0
38	Y	3	Total 3	O 3	0
38	Z	17	Total 17	O 17	0
38	0	5	Total 5	O 5	0
38	1	11	Total 11	O 11	0
38	4	7	Total 7	O 7	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 12S mitochondrial rRNA





- Molecule 2: 28S ribosomal protein S2, mitochondrial







• Molecule 3: 28S ribosomal protein S24, mitochondrial






• Molecule 4: 28S ribosomal protein S5, mitochondrial





• Molecule 5: 28S ribosomal protein S6, mitochondrial




• Molecule 6: 28S ribosomal protein S7, mitochondrial




• Molecule 7: 28S ribosomal protein S9, mitochondrial







- Molecule 8: 28S ribosomal protein S10, mitochondrial





- Molecule 9: 28S ribosomal protein S11, mitochondrial





- Molecule 10: 28S ribosomal protein S12, mitochondrial




- Molecule 11: 28S ribosomal protein S14, mitochondrial




- Molecule 12: 28S ribosomal protein S15, mitochondrial




- Molecule 13: 28S ribosomal protein S16, mitochondrial




- Molecule 14: 28S ribosomal protein S17, mitochondrial




- Molecule 15: 28S ribosomal protein S18b, mitochondrial




- Molecule 16: 28S ribosomal protein S18c, mitochondrial




- Molecule 17: 28S ribosomal protein S21, mitochondrial




- Molecule 18: 28S ribosomal protein S22, mitochondrial




• Molecule 19: 28S ribosomal protein S23, mitochondrial




• Molecule 20: 28S ribosomal protein S25, mitochondrial




• Molecule 21: 28S ribosomal protein S26, mitochondrial




• Molecule 22: 28S ribosomal protein S27, mitochondrial




• Molecule 23: 28S ribosomal protein S28, mitochondrial





• Molecule 24: 28S ribosomal protein S29, mitochondrial









• Molecule 25: 28S ribosomal protein S31, mitochondrial




• Molecule 26: 28S ribosomal protein S33, mitochondrial



• Molecule 27: 28S ribosomal protein S34, mitochondrial



• Molecule 28: 28S ribosomal protein S35, mitochondrial







## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	52361	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	31	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3600	Depositor
Magnification	165000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.622	Depositor
Minimum map value	-0.805	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.030	Depositor
Recommended contour level	0.12	Depositor
Map size (Å)	398.40192, 398.40192, 398.40192	wwPDB
Map dimensions	576, 576, 576	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.69167, 0.69167, 0.69167	Depositor

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, K, 5F0, ZN, MG, AYA, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.24	1/21600 (0.0%)	0.67	1/33606 (0.0%)
2	B	0.26	0/1832	0.41	0/2479
3	C	0.26	0/1100	0.43	0/1485
4	D	0.25	0/2737	0.42	0/3671
5	E	0.25	0/997	0.41	0/1347
6	F	0.24	0/1763	0.37	0/2368
7	G	0.25	0/2734	0.39	0/3669
8	H	0.25	0/1179	0.41	0/1597
9	I	0.25	0/1018	0.45	0/1374
10	J	0.26	0/862	0.46	0/1155
11	K	0.24	0/871	0.40	0/1167
12	L	0.23	0/1485	0.36	0/1980
13	M	0.26	0/1017	0.42	0/1366
14	N	0.26	0/907	0.45	0/1228
15	O	0.25	0/1653	0.39	0/2254
16	P	0.26	0/809	0.40	0/1085
17	Q	0.24	0/735	0.40	0/980
18	R	0.25	0/2449	0.38	0/3311
19	S	0.26	0/1148	0.39	0/1541
20	T	0.25	0/1420	0.39	0/1903
21	U	0.24	0/1470	0.37	0/1976
22	V	0.23	0/3059	0.35	0/4135
23	W	0.25	0/805	0.43	0/1084
24	X	0.24	0/2952	0.38	0/3995
25	Y	0.25	0/1283	0.36	0/1730
26	Z	0.26	0/851	0.37	0/1133
27	0	0.24	0/1856	0.42	0/2511
28	1	0.24	0/2271	0.38	0/3078
29	3	0.24	0/448	0.38	0/591
30	4	0.24	0/4868	0.36	0/6597
31	a	0.23	0/1087	0.36	0/1470
32	c	0.24	0/2447	0.38	0/3309

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
All	All	0.24	1/71713 (0.0%)	0.50	1/101175 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
9	I	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	A	OP3-P	-10.80	1.48	1.61

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	824	A	C5-C6-N1	-5.29	115.05	117.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
9	I	181	5F0	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	19301	9755	9761	40	0
2	B	1791	1799	1797	9	0
3	C	1072	1091	1087	5	0
4	D	2682	2732	2727	12	0
5	E	979	1007	1007	3	0
6	F	1722	1750	1748	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	G	2674	2630	2626	11	0
8	H	1153	1193	1190	9	0
9	I	1009	1038	1029	4	0
10	J	846	903	901	3	0
11	K	855	888	887	2	0
12	L	1465	1576	1574	3	0
13	M	995	1009	1006	4	0
14	N	888	951	947	2	0
15	O	1598	1548	1547	2	0
16	P	792	818	817	3	0
17	Q	732	750	750	3	0
18	R	2400	2416	2415	3	0
19	S	1124	1133	1132	3	0
20	T	1388	1413	1413	6	0
21	U	1449	1459	1456	2	0
22	V	2998	2972	2967	10	0
23	W	793	813	811	3	0
24	X	2881	2881	2879	10	0
25	Y	1246	1193	1191	1	0
26	Z	834	848	847	1	0
27	0	1811	1838	1834	5	0
28	1	2223	2241	2238	4	0
29	3	440	489	488	1	0
30	4	4758	4800	4793	13	0
31	a	1070	1099	1096	0	0
32	c	2393	2478	2478	0	0
33	A	32	0	0	0	0
33	B	1	0	0	0	0
33	X	1	0	0	0	0
34	A	7	0	0	0	0
35	O	1	0	0	0	0
36	P	4	0	0	0	0
36	T	4	0	0	0	0
37	X	31	12	12	0	0
38	0	5	0	0	1	0
38	1	11	0	0	0	0
38	4	7	0	0	0	0
38	A	268	0	0	4	0
38	B	7	0	0	0	0
38	C	23	0	0	1	0
38	D	14	0	0	0	0
38	F	5	0	0	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
38	G	9	0	0	1	0
38	H	11	0	0	0	0
38	I	3	0	0	0	0
38	J	4	0	0	1	0
38	K	15	0	0	0	0
38	L	1	0	0	0	0
38	M	17	0	0	0	0
38	N	2	0	0	0	0
38	O	20	0	0	1	0
38	P	6	0	0	0	0
38	Q	2	0	0	0	0
38	R	5	0	0	0	0
38	S	2	0	0	0	0
38	T	6	0	0	0	0
38	U	3	0	0	0	0
38	X	4	0	0	0	0
38	Y	3	0	0	0	0
38	Z	17	0	0	0	0
All	All	68913	59523	59451	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:567:G:N2	1:A:576:C:O2	2.20	0.74
10:J:111:GLU:OE1	38:J:201:HOH:O	2.05	0.74
1:A:217:U:OP1	15:O:146:LYS:NZ	2.22	0.73
1:A:192:A:N7	38:A:1105:HOH:O	2.21	0.72
6:F:84:SER:OG	24:X:372:GLU:OE2	2.07	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	221/291 (76%)	219 (99%)	2 (1%)	0	100	100
3	C	130/167 (78%)	124 (95%)	6 (5%)	0	100	100
4	D	333/432 (77%)	328 (98%)	5 (2%)	0	100	100
5	E	120/125 (96%)	118 (98%)	2 (2%)	0	100	100
6	F	206/242 (85%)	200 (97%)	6 (3%)	0	100	100
7	G	321/390 (82%)	315 (98%)	5 (2%)	1 (0%)	41	74
8	H	138/160 (86%)	135 (98%)	2 (1%)	1 (1%)	22	61
9	I	134/191 (70%)	130 (97%)	4 (3%)	0	100	100
10	J	106/139 (76%)	105 (99%)	1 (1%)	0	100	100
11	K	99/128 (77%)	98 (99%)	1 (1%)	0	100	100
12	L	174/258 (67%)	173 (99%)	1 (1%)	0	100	100
13	M	124/135 (92%)	124 (100%)	0	0	100	100
14	N	111/120 (92%)	108 (97%)	3 (3%)	0	100	100
15	O	195/254 (77%)	192 (98%)	3 (2%)	0	100	100
16	P	95/143 (66%)	94 (99%)	1 (1%)	0	100	100
17	Q	84/86 (98%)	83 (99%)	1 (1%)	0	100	100
18	R	292/359 (81%)	284 (97%)	8 (3%)	0	100	100
19	S	134/177 (76%)	132 (98%)	2 (2%)	0	100	100
20	T	168/171 (98%)	167 (99%)	1 (1%)	0	100	100
21	U	172/200 (86%)	170 (99%)	2 (1%)	0	100	100
22	V	361/415 (87%)	357 (99%)	4 (1%)	0	100	100
23	W	98/186 (53%)	96 (98%)	2 (2%)	0	100	100
24	X	355/391 (91%)	350 (99%)	5 (1%)	0	100	100
25	Y	147/384 (38%)	145 (99%)	2 (1%)	0	100	100
26	Z	99/106 (93%)	98 (99%)	1 (1%)	0	100	100
27	0	214/218 (98%)	211 (99%)	3 (1%)	0	100	100
28	1	276/320 (86%)	274 (99%)	2 (1%)	0	100	100
29	3	48/200 (24%)	48 (100%)	0	0	100	100
30	4	585/685 (85%)	579 (99%)	6 (1%)	0	100	100
31	a	128/350 (37%)	126 (98%)	2 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
32	c	297/345 (86%)	292 (98%)	5 (2%)	0	100	100
All	All	5965/7768 (77%)	5875 (98%)	88 (2%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	H	85	ILE
7	G	204	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	195/247 (79%)	195 (100%)	0	100	100
3	C	113/139 (81%)	113 (100%)	0	100	100
4	D	277/354 (78%)	277 (100%)	0	100	100
5	E	107/110 (97%)	107 (100%)	0	100	100
6	F	180/204 (88%)	180 (100%)	0	100	100
7	G	282/335 (84%)	281 (100%)	1 (0%)	91	95
8	H	130/150 (87%)	130 (100%)	0	100	100
9	I	102/143 (71%)	102 (100%)	0	100	100
10	J	93/117 (80%)	93 (100%)	0	100	100
11	K	90/110 (82%)	90 (100%)	0	100	100
12	L	161/224 (72%)	161 (100%)	0	100	100
13	M	103/112 (92%)	103 (100%)	0	100	100
14	N	97/104 (93%)	97 (100%)	0	100	100
15	O	176/225 (78%)	176 (100%)	0	100	100
16	P	89/125 (71%)	89 (100%)	0	100	100
17	Q	77/77 (100%)	77 (100%)	0	100	100
18	R	261/313 (83%)	261 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
19	S	117/152 (77%)	117 (100%)	0	100	100
20	T	153/154 (99%)	153 (100%)	0	100	100
21	U	149/171 (87%)	149 (100%)	0	100	100
22	V	326/362 (90%)	325 (100%)	1 (0%)	92	96
23	W	87/156 (56%)	87 (100%)	0	100	100
24	X	314/346 (91%)	312 (99%)	2 (1%)	86	94
25	Y	133/341 (39%)	133 (100%)	0	100	100
26	Z	88/93 (95%)	88 (100%)	0	100	100
27	0	190/191 (100%)	188 (99%)	2 (1%)	73	88
28	1	248/279 (89%)	248 (100%)	0	100	100
29	3	46/176 (26%)	46 (100%)	0	100	100
30	4	519/599 (87%)	519 (100%)	0	100	100
31	a	122/299 (41%)	122 (100%)	0	100	100
32	c	266/304 (88%)	265 (100%)	1 (0%)	91	95
All	All	5291/6712 (79%)	5284 (100%)	7 (0%)	93	98

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
24	X	112	HIS
27	0	30	ASP
32	c	202	TYR
27	0	48	ARG
24	X	74	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
18	R	246	HIS
30	4	372	HIS
22	V	325	GLN
30	4	376	HIS
26	Z	82	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	902/956 (94%)	138 (15%)	0

5 of 138 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	3	A
1	A	7	U
1	A	16	G
1	A	23	A
1	A	32	G

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
9	5F0	I	181	9	8,8,9	0.60	0	7,9,11	1.05	1 (14%)
17	AYA	Q	2	17	6,7,8	0.76	0	5,8,10	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	5F0	I	181	9	-	0/9/9/10	-
17	AYA	Q	2	17	-	0/4/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	I	181	5F0	O-C-CB	-2.43	118.34	125.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 45 ligands modelled in this entry, 42 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
37	ATP	X	401	33	26,33,33	0.75	0	31,52,52	0.67	0
36	FES	T	201	13,20	0,4,4	-	-	-		
36	FES	P	201	16,5	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
36	FES	T	201	13,20	-	-	0/1/1/1
36	FES	P	201	16,5	-	-	0/1/1/1
37	ATP	X	401	33	-	0/18/38/38	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13551. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.




## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 288




Y Index: 288




Z Index: 288


### 6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 348




Y Index: 265




Z Index: 241


### 6.3.2 Raw map



X Index: 262



Y Index: 232




Z Index: 231


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.12. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


### 6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_13551\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 453 nm<sup>3</sup>; this corresponds to an approximate mass of 410 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.313 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.313  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.19	-	-
Author-provided FSC curve	3.18	4.18	3.30
Unmasked-calculated*	4.13	8.66	4.25

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.13 differs from the reported value 3.19 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-13551 and PDB model 7PNT. Per-residue inclusion information can be found in section 3 on page 12.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.12 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.12).



































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 79% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.12) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8020	 0.5160
0	 0.8830	 0.5530
1	 0.8110	 0.5240
3	 0.2940	 0.3520
4	 0.5530	 0.4090
A	 0.8700	 0.5360
B	 0.9280	 0.6050
C	 0.9350	 0.5990
D	 0.8100	 0.5270
E	 0.8750	 0.5470
F	 0.7720	 0.4890
G	 0.7910	 0.5150
H	 0.8730	 0.5570
I	 0.9060	 0.5570
J	 0.8870	 0.5300
K	 0.9370	 0.5950
L	 0.7720	 0.5040
M	 0.8940	 0.6000
N	 0.8970	 0.5800
O	 0.9250	 0.5970
P	 0.9140	 0.5750
Q	 0.9090	 0.5620
R	 0.8540	 0.5550
S	 0.8020	 0.5040
T	 0.9020	 0.5750
U	 0.7710	 0.5010
V	 0.7410	 0.4880
W	 0.8710	 0.5640
X	 0.8140	 0.5200
Y	 0.6550	 0.4620
Z	 0.7990	 0.5180
a	 0.5390	 0.4140
c	 0.2460	 0.2710

