



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2024 – 02:57 AM EST

PDB ID : 4PQP
Title : Crystal structure of human SNX14 PX domain in space group P43212
Authors : Mas, C.; Norwood, S.; Bugarcic, A.; Kinna, G.; Leneva, N.; Kovtun, O.;
Teasdale, R.; Collins, B.
Deposited on : 2014-03-03
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 7% 52% 20% 8% • 20%
1	B	128	 5% 54% 21% 5% 20%
1	C	128	 5% 53% 21% 5% • 20%
1	D	128	 2% 53% 20% • 24%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3475 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorting nexin-14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	102	874	580	137	156	1	0	1	0
1	B	103	875	579	141	154	1	0	0	0
1	C	102	865	572	137	155	1	0	0	0
1	D	97	830	548	130	151	1	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	559	GLY	-	expression tag	UNP Q9Y5W7
A	560	SER	-	expression tag	UNP Q9Y5W7
B	559	GLY	-	expression tag	UNP Q9Y5W7
B	560	SER	-	expression tag	UNP Q9Y5W7
C	559	GLY	-	expression tag	UNP Q9Y5W7
C	560	SER	-	expression tag	UNP Q9Y5W7
D	559	GLY	-	expression tag	UNP Q9Y5W7
D	560	SER	-	expression tag	UNP Q9Y5W7


- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total O 4 4	0	0
3	B	6	Total O 6 6	0	0
3	C	6	Total O 6 6	0	0
3	D	3	Total O 3 3	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.67Å 121.67Å 82.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.95 – 3.00 68.24 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.95-3.00) 100.0 (68.24-3.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.76 (at 3.01Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.234 , 0.286 0.242 , 0.298	Depositor DCC
R_{free} test set	629 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	79.3	Xtrriage
Anisotropy	0.006	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 75.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3475	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/905	0.75	3/1224 (0.2%)
1	B	0.53	0/902	0.70	1/1219 (0.1%)
1	C	0.54	0/892	0.69	1/1207 (0.1%)
1	D	0.52	0/855	0.62	0/1156
All	All	0.54	0/3554	0.69	5/4806 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	648	LYS	C-N-CA	-7.86	102.06	121.70
1	B	644	ILE	CB-CA-C	-6.64	98.32	111.60
1	C	644	ILE	CB-CA-C	-6.62	98.37	111.60
1	A	586	ILE	N-CA-C	-6.29	94.02	111.00
1	A	649	ASN	N-CA-C	5.28	125.25	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	874	0	848	37	1
1	B	875	0	857	16	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	865	0	838	27	1
1	D	830	0	790	19	1
2	A	6	0	8	2	0
2	C	6	0	8	0	0
3	A	4	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	1	0
3	D	3	0	0	0	0
All	All	3475	0	3349	86	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:644:ILE:HG12	1:C:644:ILE:HD11	1.43	0.99
1:A:643:ARG:HH12	1:D:652:PHE:HD1	1.13	0.96
1:A:671:PRO:O	1:A:673:LEU:N	2.05	0.90
1:A:643:ARG:HG3	1:A:645:ILE:H	1.36	0.90
1:B:671:PRO:O	1:B:673:LEU:N	2.05	0.90

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:576:GLU:OE2	1:D:629:GLU:O[8_667]	2.05	0.15
1:B:612:SER:OG	1:C:609:GLU:OE1[5_756]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/128 (76%)	89 (92%)	5 (5%)	3 (3%)	4	23
1	B	97/128 (76%)	88 (91%)	6 (6%)	3 (3%)	4	23
1	C	96/128 (75%)	85 (88%)	7 (7%)	4 (4%)	3	16
1	D	89/128 (70%)	82 (92%)	5 (6%)	2 (2%)	6	31
All	All	379/512 (74%)	344 (91%)	23 (6%)	12 (3%)	4	22

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	672	GLU
1	B	671	PRO
1	B	672	GLU
1	C	649	ASN
1	C	672	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/118 (81%)	86 (90%)	10 (10%)	7	27
1	B	96/118 (81%)	83 (86%)	13 (14%)	4	17
1	C	95/118 (80%)	87 (92%)	8 (8%)	11	38
1	D	91/118 (77%)	86 (94%)	5 (6%)	21	57
All	All	378/472 (80%)	342 (90%)	36 (10%)	8	32

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	643	ARG
1	D	672	GLU
1	C	644	ILE
1	D	595	ARG
1	B	572	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	701	-	5,5,5	0.35	0	5,5,5	0.53	0
2	GOL	C	701	-	5,5,5	0.36	0	5,5,5	1.26	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	701	-	-	2/4/4/4	-
2	GOL	C	701	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	701	GOL	C3-C2-C1	-2.37	102.48	111.70

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	GOL	C1-C2-C3-O3
2	C	701	GOL	O1-C1-C2-C3
2	C	701	GOL	C1-C2-C3-O3
2	C	701	GOL	O1-C1-C2-O2
2	C	701	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	102/128 (79%)	0.51	9 (8%) 10 3	51, 86, 140, 198	0
1	B	103/128 (80%)	0.33	7 (6%) 17 5	55, 86, 123, 154	0
1	C	102/128 (79%)	0.45	6 (5%) 22 7	58, 87, 130, 152	0
1	D	97/128 (75%)	0.31	3 (3%) 49 21	66, 97, 127, 174	0
All	All	404/512 (78%)	0.40	25 (6%) 20 7	51, 90, 131, 198	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	643	ARG	5.2
1	A	644	ILE	3.5
1	A	576	GLU	3.3
1	C	673	LEU	3.3
1	C	644	ILE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	701	6/6	0.82	0.28	82,85,87,88	0
2	GOL	C	701	6/6	0.89	0.22	42,82,90,93	0

6.5 Other polymers [i](#)

There are no such residues in this entry.