

Full wwPDB X-ray Structure Validation Report (i)

Mar 5, 2024 - 03:09 pm GMT

PDB ID	:	8PQE
Title	:	c-KIT kinase domain in complex with a vapritinib derivative 11
Authors	:	Teuber, A.; Niggenaber, J.; Mueller, M.P.; Rauh, D.
Deposited on		
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4 02b-467
·		
Mogul	:	1.8.4, CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	327	% 8 2%	•	15%
1	В	327	2% 81%	6%	13%



8PQE

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms		ZeroOcc	AltConf	Trace	
1	Δ	277	Total	С	Ν	0	\mathbf{S}	0	F	0
	211	2173	1403	355	399	16	0	5	0	
1	D	283	Total	С	Ν	0	S	0	2	0
	ГБ	283	2209	1425	363	405	16	0		U

• Molecule 1 is a protein called Mast/stem cell growth factor receptor Kit.

Chain	Residue	Modelled	Actual	Comment	Reference
А	548	GLY	-	expression tag	UNP P10721
А	549	SER	-	expression tag	UNP P10721
А	550	MET	-	expression tag	UNP P10721
А	563	SER	ILE	engineered mutation	UNP P10721
А	569	SER	VAL	engineered mutation	UNP P10721
А	609	GLN	TYR	engineered mutation	UNP P10721
А	631	SER	LEU	engineered mutation	UNP P10721
А	651	GLU	MET	engineered mutation	UNP P10721
А	662	HIS	ILE	engineered mutation	UNP P10721
А	688	GLU	-	linker	UNP P10721
А	689	PHE	-	linker	UNP P10721
А	690	VAL	-	linker	UNP P10721
А	691	PRO	-	linker	UNP P10721
А	692	TYR	-	linker	UNP P10721
А	693	LYS	-	linker	UNP P10721
А	754	VAL	-	linker	UNP P10721
А	755	ALA	-	linker	UNP P10721
А	756	PRO	-	linker	UNP P10721
А	757	GLU	-	linker	UNP P10721
А	758	ASP	-	linker	UNP P10721
А	759	LEU	-	linker	UNP P10721
А	760	TYR	-	linker	UNP P10721
А	761	LYS	-	linker	UNP P10721
А	762	ASP	-	linker	UNP P10721
А	763	PHE	-	linker	UNP P10721

There are 70 discrepancies between the modelled and reference sequences:

Continued on next page...



8PQE	
------	--

Continued from previous page							
Chain	Residue	Modelled	Actual	Comment	Reference		
А	764	LEU	-	linker	UNP P10721		
А	765	THR	-	linker	UNP P10721		
А	768	HIS	ASP	engineered mutation	UNP P10721		
А	804	ASN	ARG	engineered mutation	UNP P10721		
А	825	ASP	VAL	engineered mutation	UNP P10721		
А	844	SER	CYS	engineered mutation	UNP P10721		
А	890	SER	LEU	engineered mutation	UNP P10721		
А	894	TYR	HIS	engineered mutation	UNP P10721		
А	912	ASP	LEU	engineered mutation	UNP P10721		
А	923	ASP	LEU	engineered mutation	UNP P10721		
В	548	GLY	-	expression tag	UNP P10721		
В	549	SER	-	expression tag	UNP P10721		
В	550	MET	-	expression tag	UNP P10721		
В	563	SER	ILE	engineered mutation	UNP P10721		
В	569	SER	VAL	engineered mutation	UNP P10721		
В	609	GLN	TYR	engineered mutation	UNP P10721		
В	631	SER	LEU	engineered mutation	UNP P10721		
В	651	GLU	MET	engineered mutation	UNP P10721		
В	662	HIS	ILE	engineered mutation	UNP P10721		
В	688	GLU	-	linker	UNP P10721		
В	689	PHE	-	linker	UNP P10721		
В	690	VAL	-	linker	UNP P10721		
В	691	PRO	-	linker	UNP P10721		
В	692	TYR	-	linker	UNP P10721		
В	753	LYS	-	linker	UNP P10721		
В	754	VAL	-	linker	UNP P10721		
В	755	ALA	-	linker	UNP P10721		
В	756	PRO	-	linker	UNP P10721		
В	757	GLU	-	linker	UNP P10721		
В	758	ASP	-	linker	UNP P10721		
В	759	LEU	_	linker	UNP P10721		
В	760	TYR	-	linker	UNP P10721		
В	761	LYS	_	linker	UNP P10721		
В	762	ASP	-	linker	UNP P10721		
В	763	PHE	-	linker	UNP P10721		
В	764	LEU	-	linker	UNP P10721		
В	765	THR	-	linker	UNP P10721		
В	768	HIS	ASP	engineered mutation	UNP P10721		
В	804	ASN	ARG	engineered mutation	UNP P10721		
В	825	ASP	VAL	engineered mutation	UNP P10721		
В	844	SER	CYS	engineered mutation	UNP P10721		
В	890	SER	LEU	engineered mutation	UNP P10721		

Continued from previous page...


Continued on next page...



Commu	Continued from previous page										
Chain	Residue	Modelled	Actual	Comment	Reference						
В	894	TYR	HIS	engineered mutation	UNP P10721						
В	912	ASP	LEU	engineered mutation	UNP P10721						
В	923	ASP	LEU	engineered mutation	UNP P10721						

Continued from previous page...


• Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	TotalCO422	0	0

• Molecule 3 is $\{N\}-[(1 \{S\})-1-(4-fluorophenyl)-1-[2-[4-[6-(1-methylpyrazol-4-yl)pyrrolo[2,1 -f][1,2,4]triazin-4-yl]piperazin-1-yl]pyrimidin-5-yl]ethyl]prop-2-enamide (three-letter code: 9WU) (formula: C₂₉H₂₉FN₁₀O) (labeled as "Ligand of Interest" by depositor).$





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	Λ	1	Total	С	F	Ν	Ο	0	0
0	D A	1	41	29	1	10	1	0	
2	р	1	Total	С	F	Ν	Ο	0	0
0	3 B	1	41	29	1	10	1	0	0

• Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Br 1 1	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	68	Total O 68 68	0	0
6	В	69	Total O 69 69	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Mast/stem cell growth factor receptor Kit



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	58.97Å 62.52Å 191.68Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.69 - 2.00	Depositor
Resolution (A)	44.69 - 2.00	EDS
% Data completeness	99.9 (44.69-2.00)	Depositor
(in resolution range)	$100.0 \ (44.69-2.00)$	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.36 (at 2.00 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.213 , 0.256	Depositor
R, R_{free}	0.210 , 0.252	DCC
R_{free} test set	2443 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	49.2	Xtriage
Anisotropy	0.558	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 55.8	EDS
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4607	wwPDB-VP
Average B, all atoms $(Å^2)$	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, 9WU, EDO, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.29	0/2243	0.51	0/3042	
1	В	0.29	0/2272	0.51	0/3082	
All	All	0.29	0/4515	0.51	0/6124	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2173	0	2095	5	0
1	В	2209	0	2128	11	0
2	А	4	0	6	0	0
3	А	41	0	0	1	0
3	В	41	0	0	1	0
4	А	1	0	0	0	0
5	В	1	0	0	0	0
6	А	68	0	0	0	0
6	В	69	0	0	1	0
All	All	4607	0	4229	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:892:PRO:HG2	1:B:895:ALA:HB2	1.74	0.69
1:B:789[A]:ILE:HG22	1:B:791:ARG:HG2	1.90	0.53
1:B:680[A]:ASN:OD1	1:B:684:ARG:NH1	2.41	0.53
1:B:625:LEU:HG	1:B:637:LEU:HD22	1.91	0.52
1:A:599:ALA:H	3:A:1002:9WU:C27	2.22	0.52
1:A:789[A]:ILE:HG22	1:A:791:ARG:HG2	1.94	0.50
1:B:634:ARG:NH1	6:B:1106:HOH:O	2.45	0.49
1:B:860:TRP:CD1	1:B:892:PRO:HD3	2.48	0.48
1:A:881:LYS:NZ	1:A:885:GLU:OE2	2.44	0.45
1:B:690:VAL:O	1:B:765:THR:HA	2.17	0.45
1:B:910:ASP:HB3	1:B:913:LYS:HD2	1.98	0.44
1:B:677:ASP:OD1	1:B:680[B]:ASN:ND2	2.52	0.43
3:B:1001:9WU:C41	3:B:1001:9WU:C15	2.97	0.43
1:A:585:PRO:HB2	1:A:588:ARG:HG2	2.01	0.42
1:A:774:TYR:CE1	1:A:925:GLU:HG3	2.54	0.42
1:B:601:GLY:HA2	1:B:626:LYS:HG3	2.02	0.41
1:B:618:MET:HB3	1:B:618:MET:HE2	1.89	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allo		Allowed	Outliers	Perce	ntiles
1	А	274/327~(84%)	273 (100%)	1 (0%)	0	100	100
1	В	279/327~(85%)	276~(99%)	3~(1%)	0	100	100
All	All	553/654~(85%)	549 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	229/285~(80%)	229 (100%)	0	100 100
1	В	232/285~(81%)	232 (100%)	0	100 100
All	All	461/570 (81%)	461 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Turne	Chain	Res	Link	Bo	ond leng	\mathbf{ths}	E	ond ang	gles
10101	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	EDO	А	1001	-	$3,\!3,\!3$	0.44	0	2,2,2	0.34	0
3	9WU	В	1001	-	39,46,46	2.03	7 (17%)	48,67,67	3.92	23 (47%)
3	9WU	А	1002	-	39,46,46	2.09	7 (17%)	48,67,67	3.79	22 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	А	1001	-	-	0/1/1/1	-
3	9WU	В	1001	-	-	8/31/41/41	0/6/6/6
3	9WU	А	1002	-	-	4/31/41/41	0/6/6/6

All (14) bond length outliers are listed below:	
---	--

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	А	1002	9WU	C40-N09	-8.09	1.30	1.40
3	В	1001	9WU	C40-N09	-7.68	1.31	1.40
3	А	1002	9WU	C18-N17	5.13	1.45	1.35
3	В	1001	9WU	C18-N17	4.87	1.44	1.35
3	В	1001	9WU	C05-N02	-4.39	1.31	1.35
3	А	1002	9WU	C05-N02	-4.00	1.31	1.35
3	А	1002	9WU	C26-C25	3.69	1.54	1.48
3	В	1001	9WU	C25-N24	3.36	1.47	1.35
3	В	1001	9WU	C26-C25	3.29	1.53	1.48
3	А	1002	9WU	C25-N24	3.17	1.46	1.35
3	В	1001	9WU	O28-C25	-3.01	1.18	1.24
3	А	1002	9WU	O28-C25	-2.86	1.18	1.24
3	В	1001	9WU	C13-N14	2.66	1.45	1.37
3	А	1002	9WU	C13-N14	2.27	1.43	1.37

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	В	1001	9WU	C23-C22-N24	-12.92	86.61	109.69
3	А	1002	9WU	C23-C22-N24	-12.21	87.89	109.69
3	А	1002	9WU	C29-C22-N24	9.34	126.71	109.17
3	В	1001	9WU	C29-C22-N24	8.04	124.27	109.17
3	В	1001	9WU	C23-C22-C29	-7.27	92.67	111.61
3	В	1001	9WU	C23-C22-C21	-7.18	92.91	111.61

Continued on next page...



Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
А	1002	9WU	C23-C22-C21	-7.06	93.23	111.61
А	1002	9WU	C23-C22-C29	-7.03	93.30	111.61
В	1001	9WU	C36-N37-C18	6.33	121.25	115.64
В	1001	9WU	C20-N19-C18	6.26	121.19	115.64
В	1001	9WU	N10-C11-N12	-6.24	118.78	128.59
А	1002	9WU	C36-N37-C18	6.23	121.16	115.64
А	1002	9WU	C20-N19-C18	6.21	121.14	115.64
В	1001	9WU	C21-C22-N24	5.91	120.26	109.17
А	1002	9WU	N10-C11-N12	-5.86	119.38	128.59
В	1001	9WU	C21-C20-N19	-5.40	118.18	123.34
А	1002	9WU	C16-N17-C18	-5.30	112.89	121.69
А	1002	9WU	C21-C36-N37	-5.10	118.48	123.34
В	1001	9WU	N19-C18-N17	4.98	122.28	116.90
В	1001	9WU	C38-N17-C18	-4.89	113.58	121.69
А	1002	9WU	C21-C22-N24	4.83	118.23	109.17
В	1001	9WU	C38-N17-C16	4.73	121.95	111.52
А	1002	9WU	C21-C20-N19	-4.38	119.16	123.34
А	1002	9WU	C38-N17-C16	4.27	120.94	111.52
В	1001	9WU	C04-N03-N02	4.10	108.50	104.23
В	1001	9WU	C21-C36-N37	-4.10	119.42	123.34
А	1002	9WU	C04-N03-N02	4.01	108.40	104.23
В	1001	9WU	C39-N14-C13	-3.60	108.47	118.73
А	1002	9WU	C15-C16-N17	3.57	117.63	110.70
В	1001	9WU	C01-N02-N03	3.56	124.69	120.50
А	1002	9WU	N37-C18-N17	3.50	120.69	116.90
В	1001	9WU	C16-N17-C18	-3.42	116.02	121.69
А	1002	9WU	N37-C18-N19	-3.30	121.49	127.06
В	1001	9WU	N37-C18-N19	-3.28	121.53	127.06
В	1001	9WU	O28-C25-C26	-3.20	117.61	122.72
А	1002	9WU	C39-N14-C13	-3.19	109.62	118.73
А	1002	9WU	C08-C07-C06	-3.15	123.61	127.74
В	1001	9WU	C27-C26-C25	-3.09	118.74	122.27
А	1002	9WU	C15-N14-C13	-3.02	110.10	118.73
А	1002	9WU	C01-N02-N03	2.95	123.97	120.50
А	1002	9WU	C41-C07-C06	2.68	128.79	126.32
	1001	OTTTT			110.00	100.07

ContinMol

There are no chirality outliers.

В

А

В

В

3

3

3

All (12) torsion outliers are listed below:

1001

1002

1001

1001

9WU

9WU

9WU

9WU

C30-C29-C22

C38-N17-C18

C39-C38-N17

C26-C25-N24



-2.57

-2.29

2.26

2.18

116.03

117.90

115.09

120.37

120.85

121.69

110.70

114.77

Mol	Chain	Res	Type	Atoms
3	А	1002	9WU	C29-C22-N24-C25
3	А	1002	9WU	N24-C25-C26-C27
3	А	1002	9WU	O28-C25-C26-C27
3	В	1001	9WU	N12-C13-N14-C15
3	В	1001	9WU	N12-C13-N14-C39
3	В	1001	9WU	C40-C13-N14-C15
3	В	1001	9WU	C40-C13-N14-C39
3	В	1001	9WU	C29-C22-N24-C25
3	В	1001	9WU	N24-C25-C26-C27
3	В	1001	9WU	O28-C25-C26-C27
3	В	1001	9WU	N19-C18-N17-C38
3	А	1002	9WU	N19-C18-N17-C38


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	\mathbf{Res}	Type	Clashes	Symm-Clashes
3	В	1001	9WU	1	0
3	А	1002	9WU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RS	SRZ:	>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	277/327~(84%)	0.22	3 (1%)	80	79	44, 58, 87, 115	0
1	В	283/327~(86%)	0.28	6 (2%)	63	62	48, 61, 93, 127	0
All	All	560/654~(85%)	0.25	9 (1%)	72	70	44, 60, 91, 127	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	692	TYR	3.6
1	В	928	ILE	3.4
1	А	628	SER	2.9
1	А	646	TYR	2.9
1	В	687	ASP	2.6
1	А	928	ILE	2.5
1	В	689	PHE	2.3
1	В	801	THR	2.2
1	В	630	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	$\mathbf{Q} < 0.9$
2	EDO	А	1001	4/4	0.79	0.16	71,73,75,79	0
3	9WU	А	1002	41/41	0.94	0.15	48,66,78,92	0
3	9WU	В	1001	41/41	0.95	0.13	47,59,76,86	0
4	BR	А	1003	1/1	0.99	0.26	47,47,47,47	0
5	CL	В	1002	1/1	0.99	0.32	72,72,72,72	0

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

