



Full wwPDB EM Validation Report ⓘ

Jan 2, 2024 – 05:40 pm GMT

PDB ID : 8PSZ
EMDB ID : EMD-17865
Title : Tilapia Lake Virus polymerase in vRNA elongation state with additional mode B promoter (transcriptase conformation)
Authors : Arragain, B.; Cusack, S.
Deposited on : 2023-07-13
Resolution : 2.42 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 92%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 100%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
2	B	519	<div style="display: flex; align-items: center;"> <div style="width: 1%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 90%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: grey; margin-right: 5px;"></div> <div style="width: 100%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
3	C	478	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey; margin-right: 5px;"></div> <div style="width: 100%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
4	D	40	<div style="display: flex; align-items: center;"> <div style="width: 1%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 32%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 65%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
4	S	40	<div style="display: flex; align-items: center;"> <div style="width: 1%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 28%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 68%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
4	V	40	<div style="display: flex; align-items: center;"> <div style="width: 25%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 68%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>
5	P	21	<div style="display: flex; align-items: center;"> <div style="width: 19%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 57%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 38%; height: 10px; background-color: grey; margin-left: 5px;"></div> </div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 23111 atoms, of which 11208 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase acidic protein (PA-like).

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	A	417	6653	2105	3315	597	617	19	0	0

- Molecule 2 is a protein called Putative PB1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	B	513	7958	2491	3987	688	762	30	2	0

- Molecule 3 is a protein called RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	C	425	6652	2111	3326	592	609	14	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	391	LYS	ARG	conflict	UNP A0A7G3S745
C	458	GLY	-	expression tag	UNP A0A7G3S745
C	459	SER	-	expression tag	UNP A0A7G3S745
C	460	GLY	-	expression tag	UNP A0A7G3S745
C	461	SER	-	expression tag	UNP A0A7G3S745
C	462	GLU	-	expression tag	UNP A0A7G3S745
C	463	ASN	-	expression tag	UNP A0A7G3S745
C	464	LEU	-	expression tag	UNP A0A7G3S745
C	465	TYR	-	expression tag	UNP A0A7G3S745
C	466	PHE	-	expression tag	UNP A0A7G3S745
C	467	GLN	-	expression tag	UNP A0A7G3S745
C	468	GLY	-	expression tag	UNP A0A7G3S745
C	469	HIS	-	expression tag	UNP A0A7G3S745
C	470	HIS	-	expression tag	UNP A0A7G3S745
C	471	HIS	-	expression tag	UNP A0A7G3S745

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	472	HIS	-	expression tag	UNP A0A7G3S745
C	473	HIS	-	expression tag	UNP A0A7G3S745
C	474	HIS	-	expression tag	UNP A0A7G3S745
C	475	HIS	-	expression tag	UNP A0A7G3S745
C	476	HIS	-	expression tag	UNP A0A7G3S745
C	477	HIS	-	expression tag	UNP A0A7G3S745
C	478	HIS	-	expression tag	UNP A0A7G3S745

- Molecule 4 is a RNA chain called 5' vRNA end - vRNA loop (40-mer).

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
4	V	13	Total	C	H	N	O	P	0	0
			406	121	138	42	92	13		
4	S	13	Total	C	H	N	O	P	0	0
			422	126	140	53	90	13		
4	D	14	Total	C	H	N	O	P	0	0
			453	135	150	55	99	14		


- Molecule 5 is a RNA chain called Transcription-like product.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
5	P	13	Total	C	H	N	O	P	0	0
			411	123	139	46	90	13		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
6	A	1	Total	Zn	0
			1	1	
6	C	2	Total	Zn	0
			2	2	

- Molecule 7 is [(2 {R},3 {S},4 {R},5 {R})-5-(4-azanyl-2-oxidanylidene-pyrimidin-1-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy- {N}-[oxidanyl(phosphonoxy)phosphoryl]phosphonamic acid (three-letter code: A0I) (formula: C₉H₁₇N₄O₁₃P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
7	B	1	42	9	13	4	13	3	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
8	B	2	2	2	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
9	A	42	42	42	0
9	B	40	40	40	0
9	C	7	7	7	0
9	V	14	14	14	0
9	S	2	2	2	0
9	D	4	4	4	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Polymerase acidic protein (PA-like)




- Molecule 2: Putative PB1



- Molecule 3: RNA-dependent RNA polymerase






- Molecule 4: 5' vRNA end - vRNA loop (40-mer)




- Molecule 4: 5' vRNA end - vRNA loop (40-mer)



- Molecule 4: 5' vRNA end - vRNA loop (40-mer)



- Molecule 5: Transcription-like product



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	103012	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.080	Depositor
Minimum map value	-0.031	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.012	Depositor
Map size (Å)	251.99998, 251.99998, 251.99998	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84, 0.84, 0.84	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, A0I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3413	0.51	0/4611
2	B	0.25	0/4043	0.50	0/5472
3	C	0.26	0/3401	0.50	0/4594
4	D	0.18	0/338	0.70	0/523
4	S	0.22	0/316	0.74	0/491
4	V	0.21	0/297	0.76	0/458
5	P	0.20	0/302	0.70	0/464
All	All	0.25	0/12110	0.53	0/16613

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	176	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3338	3315	3313	13	0
2	B	3971	3987	3984	23	0
3	C	3326	3326	3322	19	0
4	D	303	150	151	1	0
4	S	282	140	140	2	0
4	V	268	138	139	1	0
5	P	272	139	140	0	0
6	A	1	0	0	0	0
6	C	2	0	0	0	0
7	B	29	13	0	0	0
8	B	2	0	0	0	0
9	A	42	0	0	1	0
9	B	40	0	0	0	0
9	C	7	0	0	0	0
9	D	4	0	0	0	0
9	S	2	0	0	0	0
9	V	14	0	0	1	0
All	All	11903	11208	11189	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:74:SER:OG	4:S:29:A:OP1	1.87	0.91
4:V:9:U:OP2	9:V:101:HOH:O	2.05	0.74
1:A:128:SER:O	2:B:44:ASN:ND2	2.24	0.70
1:A:309:GLY:O	9:A:601:HOH:O	2.12	0.68
3:C:89:GLU:N	3:C:89:GLU:OE1	2.27	0.67
1:A:322:LEU:HD11	2:B:383:LEU:HD11	1.77	0.67
3:C:353:ALA:O	3:C:359:ARG:NH1	2.29	0.66
2:B:199:ALA:HB1	2:B:292:ILE:HG21	1.80	0.64
2:B:418:ILE:HD11	2:B:428:CYS:HB3	1.79	0.64
2:B:103:THR:OG1	2:B:125:ILE:O	2.09	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:30:SER:OG	2:B:219:GLU:OE2	2.19	0.60
1:A:303:LYS:NZ	2:B:19:SER:O	2.35	0.59
2:B:103:THR:O	2:B:103:THR:HG23	2.04	0.58
3:C:81:SER:OG	3:C:113:VAL:HG23	2.04	0.57
2:B:103:THR:HG21	2:B:111:LEU:HD13	1.85	0.57
2:B:199:ALA:CB	2:B:292:ILE:HG21	2.35	0.56
1:A:267:THR:O	4:D:12:A:N6	2.37	0.56
3:C:247:PRO:O	3:C:248:THR:HG23	2.07	0.55
1:A:8:LEU:HD22	1:A:100:ILE:HD11	1.89	0.53
1:A:77:ALA:O	1:A:78:THR:HG23	2.10	0.51
3:C:144:ASP:OD1	3:C:144:ASP:N	2.44	0.51
1:A:43:ASP:OD2	1:A:62:ARG:NH1	2.45	0.50
2:B:479:SER:OG	2:B:482:ASP:OD1	2.30	0.49
3:C:334:THR:HG21	3:C:342:CYS:HB3	1.95	0.48
3:C:83:ARG:NH1	3:C:83:ARG:HB3	2.29	0.48
1:A:239:GLY:N	1:A:251:ALA:O	2.36	0.47
3:C:151:VAL:HG12	3:C:250:LEU:HD12	1.98	0.46
1:A:320:TYR:CZ	2:B:13:LEU:HD12	2.50	0.46
2:B:59:GLU:HA	2:B:59:GLU:OE1	2.16	0.46
2:B:357:VAL:HG13	2:B:444:ASP:HB2	1.97	0.45
2:B:513:VAL:HG12	2:B:513:VAL:O	2.16	0.45
2:B:114:LYS:O	2:B:118:VAL:HG13	2.16	0.45
3:C:228:VAL:O	3:C:228:VAL:HG13	2.16	0.45
2:B:189:THR:O	2:B:189:THR:CG2	2.65	0.44
2:B:292:ILE:HD13	2:B:328:GLU:O	2.19	0.43
2:B:157:ILE:HG21	4:S:30:G:C5	2.54	0.43
1:A:101:ASP:N	1:A:101:ASP:OD1	2.52	0.43
3:C:94:LEU:HD23	3:C:99:TYR:CD2	2.55	0.42
3:C:187:ASP:O	3:C:187:ASP:OD1	2.38	0.42
2:B:292:ILE:O	2:B:292:ILE:HG23	2.19	0.42
3:C:361:ILE:HD12	3:C:363:ILE:HD11	2.01	0.42
2:B:160:SER:OG	2:B:165:ARG:NH1	2.50	0.42
3:C:371:ASP:O	3:C:375:ARG:HG2	2.20	0.42
1:A:323:ASP:CG	1:A:323:ASP:O	2.58	0.41
2:B:103:THR:HG21	2:B:111:LEU:CD1	2.51	0.41
1:A:160:GLU:H	1:A:160:GLU:CD	2.24	0.41
3:C:258:PHE:CD1	3:C:264:VAL:HG23	2.56	0.41
3:C:377:PHE:O	3:C:381:LYS:O	2.39	0.41
3:C:80:LEU:HD11	3:C:112:VAL:HG13	2.02	0.41
3:C:177:ILE:HD12	3:C:259:CYS:SG	2.61	0.40
3:C:251:ASP:HB2	3:C:252:PRO:HD2	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:80:LEU:CD1	3:C:112:VAL:HG13	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/419 (99%)	403 (97%)	12 (3%)	0	100	100
2	B	511/519 (98%)	489 (96%)	22 (4%)	0	100	100
3	C	421/478 (88%)	397 (94%)	24 (6%)	0	100	100
All	All	1347/1416 (95%)	1289 (96%)	58 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/365 (100%)	351 (96%)	13 (4%)	35	52
2	B	445/447 (100%)	427 (96%)	18 (4%)	31	48
3	C	357/405 (88%)	340 (95%)	17 (5%)	25	40
All	All	1166/1217 (96%)	1118 (96%)	48 (4%)	34	47

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	7	GLN
1	A	13	CYS
1	A	17	THR
1	A	18	TYR
1	A	31	THR
1	A	133	ARG
1	A	189	ARG
1	A	208	LYS
1	A	278	SER
1	A	332	ARG
1	A	376	SER
1	A	399	SER
2	B	21	LYS
2	B	30	SER
2	B	42	SER
2	B	122	SER
2	B	145	ARG
2	B	154	SER
2	B	176	ARG
2	B	180	SER
2	B	182	VAL
2	B	255	ASP
2	B	327	LEU
2	B	337	ASP
2	B	415	LYS
2	B	449	SER
2	B	461	ARG
2	B	475	ARG
2	B	484	SER
2	B	487	ARG
3	C	40	LYS
3	C	45	ARG
3	C	77	ARG
3	C	88	GLN
3	C	111	ARG
3	C	120	SER
3	C	199	LYS
3	C	207	SER
3	C	291	ASN
3	C	301	SER
3	C	317	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	345	ASP
3	C	356	SER
3	C	381	LYS
3	C	388	SER
3	C	393	MET
3	C	401	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	D	12/40 (30%)	1 (8%)	0
4	S	12/40 (30%)	0	0
4	V	12/40 (30%)	2 (16%)	0
5	P	11/21 (52%)	1 (9%)	0
All	All	47/141 (33%)	4 (8%)	0

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	V	7	C
4	V	10	U
5	P	14	A
4	D	12	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	A0I	B	601	8	29,30,30	1.03	4 (13%)	42,47,47	1.09	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	A0I	B	601	8	-	3/19/38/38	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	601	A0I	P2-O5	2.71	1.50	1.46
7	B	601	A0I	P1-O11	2.68	1.50	1.46
7	B	601	A0I	P1-O4	-2.37	1.50	1.56
7	B	601	A0I	P2-O6	-2.34	1.50	1.56

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	601	A0I	O6-P2-O5	4.20	118.73	109.92
7	B	601	A0I	O4-P1-O11	4.04	118.39	109.92
7	B	601	A0I	O3-P1-O11	-2.41	104.95	114.24

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	601	A0I	P2-N4-P1-O3

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	B	601	A0I	P1-N4-P2-O5
7	B	601	A0I	C6-C5-N1-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
5	P	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	P	6:A	O3'	12:C	P	14.95


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17865. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 154




Y Index: 138




Z Index: 157


6.3.2 Raw map



X Index: 154



Y Index: 138




Z Index: 157


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.012. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_17865_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 48 nm³; this corresponds to an approximate mass of 43 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.413 Å⁻¹

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.413 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.42	-	-
Author-provided FSC curve	2.41	2.87	2.46
Unmasked-calculated*	3.05	3.74	3.12

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.05 differs from the reported value 2.42 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-17865 and PDB model 8PSZ. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.012 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.012).

















9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.012) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8200	 0.6470
A	 0.8510	 0.6590
B	 0.8790	 0.6670
C	 0.7570	 0.6250
D	 0.8020	 0.5780
P	 0.6990	 0.5890
S	 0.8440	 0.6130
V	 0.8950	 0.6380

