



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 10:43 am BST

PDB ID : 1PV4
Title : X-ray crystal structure of the Rho transcription termination factor in complex with single stranded DNA
Authors : Skordalakes, E.; Berger, J.M.
Deposited on : 2003-06-26
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	2	
1	H	2	
1	J	2	
1	K	2	
1	L	2	
2	A	419	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	419	<p>7% 37% 39% 8% 15%</p>
2	C	419	<p>2% 40% 45% 12% ..</p>
2	D	419	<p>5% 41% 44% 11% ..</p>
2	E	419	<p>3% 38% 47% 11% ..</p>
2	F	419	<p>7% 39% 49% 8% ..</p>

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 19077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	G	2	Total 38	C 18	N 6	O 12	P 2	0	0	0
1	H	2	Total 38	C 18	N 6	O 12	P 2	0	0	0
1	J	2	Total 38	C 18	N 6	O 12	P 2	0	0	0
1	K	2	Total 38	C 18	N 6	O 12	P 2	0	0	0
1	L	2	Total 38	C 18	N 6	O 12	P 2	0	0	0

- Molecule 2 is a protein called Transcription termination factor rho.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	A	408	Total 3208	C 2025	N 562	O 604	S 1	Se 16	0	0	0
2	B	358	Total 2813	C 1776	N 494	O 529	S 1	Se 13	0	0	0
2	C	408	Total 3208	C 2025	N 562	O 604	S 1	Se 16	0	0	0
2	D	408	Total 3208	C 2025	N 562	O 604	S 1	Se 16	0	0	0
2	E	407	Total 3201	C 2020	N 561	O 603	S 1	Se 16	0	0	0
2	F	408	Total 3208	C 2025	N 562	O 604	S 1	Se 16	0	0	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
A	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
B	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
C	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
D	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
E	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	1	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	21	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	29	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	147	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	186	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	205	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	219	MSE	MET	MODIFIED RESIDUE	UNP P0AG30

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	245	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	327	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	341	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	380	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	390	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	396	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	405	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	415	MSE	MET	MODIFIED RESIDUE	UNP P0AG30
F	416	MSE	MET	MODIFIED RESIDUE	UNP P0AG30

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total O 1 1	0	0
3	J	1	Total O 1 1	0	0
3	A	3	Total O 3 3	0	0
3	B	2	Total O 2 2	0	0
3	C	7	Total O 7 7	0	0
3	D	7	Total O 7 7	0	0
3	E	14	Total O 14 14	0	0
3	F	6	Total O 6 6	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(P*CP*C)-3'



- Molecule 1: 5'-D(P*CP*C)-3'




- Molecule 1: 5'-D(P*CP*C)-3'



- Molecule 1: 5'-D(P*CP*C)-3'




- Molecule 1: 5'-D(P*CP*C)-3'




- Molecule 2: Transcription termination factor rho






• Molecule 2: Transcription termination factor rho




• Molecule 2: Transcription termination factor rho






• Molecule 2: Transcription termination factor rho




• Molecule 2: Transcription termination factor rho





• Molecule 2: Transcription termination factor rho



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.56Å 204.33Å 147.37Å 90.00° 95.86° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00 29.89 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.9 (20.00-3.00) 96.6 (29.89-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.270 , 0.296 0.244 , 0.269	Depositor DCC
R_{free} test set	3735 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	86.7	Xtrriage
Anisotropy	0.162	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 102.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	19077	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.67	0/41	1.16	0/60
1	H	0.89	0/41	1.34	1/60 (1.7%)
1	J	0.99	0/41	1.58	1/60 (1.7%)
1	K	0.95	0/41	1.59	1/60 (1.7%)
1	L	0.93	0/41	1.79	3/60 (5.0%)
2	A	0.57	0/3238	0.77	9/4334 (0.2%)
2	B	0.57	0/2841	0.80	13/3805 (0.3%)
2	C	0.76	0/3238	0.88	11/4334 (0.3%)
2	D	0.74	0/3238	0.90	11/4334 (0.3%)
2	E	0.70	0/3231	0.89	11/4324 (0.3%)
2	F	0.60	0/3238	0.80	11/4334 (0.3%)
All	All	0.67	0/19229	0.85	72/25765 (0.3%)

There are no bond length outliers.

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	233	ASP	CB-CG-OD2	10.35	127.61	118.30
2	E	163	ASP	CB-CG-OD2	7.97	125.47	118.30
2	D	233	ASP	CB-CG-OD1	-7.79	111.29	118.30
2	D	77	ASP	CB-CG-OD2	7.54	125.08	118.30
2	A	265	ASP	CB-CG-OD2	7.30	124.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	38	0	23	2	0
1	H	38	0	23	3	0
1	J	38	0	23	5	0
1	K	38	0	23	5	0
1	L	38	0	23	8	0
2	A	3208	0	3284	278	0
2	B	2813	0	2870	229	0
2	C	3208	0	3284	308	0
2	D	3208	0	3284	273	0
2	E	3201	0	3275	308	0
2	F	3208	0	3284	259	0
3	A	3	0	0	2	0
3	B	2	0	0	2	0
3	C	7	0	0	1	0
3	D	7	0	0	3	0
3	E	14	0	0	6	0
3	F	6	0	0	3	0
3	G	1	0	0	0	0
3	J	1	0	0	0	0
All	All	19077	0	19396	1597	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 1597 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:355:PHE:HB3	2:F:356:PRO:CD	1.59	1.33
2:C:379:LYS:HG2	2:C:412:PHE:CD2	1.72	1.23
2:C:177:VAL:CG1	2:C:321:ILE:HD12	1.69	1.22
2:A:265:ASP:O	2:A:318:THR:HB	1.36	1.22
2:C:141:ALA:CB	2:C:370:LEU:HB2	1.72	1.19

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	400/419 (96%)	363 (91%)	30 (8%)	7 (2%)	8	37
2	B	350/419 (84%)	306 (87%)	31 (9%)	13 (4%)	3	19
2	C	400/419 (96%)	354 (88%)	37 (9%)	9 (2%)	6	30
2	D	400/419 (96%)	352 (88%)	33 (8%)	15 (4%)	3	18
2	E	399/419 (95%)	346 (87%)	39 (10%)	14 (4%)	3	20
2	F	400/419 (96%)	352 (88%)	35 (9%)	13 (3%)	4	21
All	All	2349/2514 (93%)	2073 (88%)	205 (9%)	71 (3%)	4	24

5 of 71 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	357	ALA
2	B	140	HIS
2	B	355	PHE
2	C	140	HIS
2	C	408	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	350/343 (102%)	308 (88%)	42 (12%)	5	22
2	B	306/343 (89%)	273 (89%)	33 (11%)	6	26
2	C	350/343 (102%)	288 (82%)	62 (18%)	2	9
2	D	350/343 (102%)	291 (83%)	59 (17%)	2	11
2	E	349/343 (102%)	291 (83%)	58 (17%)	2	11
2	F	350/343 (102%)	309 (88%)	41 (12%)	5	22
All	All	2055/2058 (100%)	1760 (86%)	295 (14%)	3	15

5 of 295 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	407	LYS
2	D	245	MSE
2	F	194	SER
2	D	7	LYS
2	D	114	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	41	GLN
2	D	129	ASN
2	E	135	ASN
2	C	306	ASN
2	E	275	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	2/2 (100%)	0.90	0 100 100	150, 150, 150, 150	0
1	H	2/2 (100%)	2.21	1 (50%) 0 0	150, 150, 150, 150	0
1	J	2/2 (100%)	3.94	2 (100%) 0 0	150, 150, 150, 150	0
1	K	2/2 (100%)	2.99	1 (50%) 0 0	150, 150, 150, 150	0
1	L	2/2 (100%)	2.87	2 (100%) 0 0	150, 150, 150, 150	0
2	A	392/419 (93%)	0.44	37 (9%) 8 3	35, 69, 121, 127	0
2	B	345/419 (82%)	0.39	28 (8%) 12 3	30, 69, 114, 127	0
2	C	392/419 (93%)	0.10	8 (2%) 65 36	18, 44, 96, 113	0
2	D	392/419 (93%)	0.02	19 (4%) 30 11	14, 45, 77, 126	0
2	E	391/419 (93%)	0.04	11 (2%) 53 25	18, 47, 87, 126	0
2	F	392/419 (93%)	0.36	30 (7%) 13 4	36, 67, 114, 124	0
All	All	2314/2524 (91%)	0.23	139 (6%) 21 7	14, 57, 110, 150	0

The worst 5 of 139 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	51	GLY	7.8
2	F	47	GLU	6.8
2	F	286	THR	6.1
2	B	141	ALA	5.9
2	E	414	GLU	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.