

Full wwPDB X-ray Structure Validation Report (i)

Jan 20, 2024 – 03:56 pm GMT

PDB ID : 7PVI

Title : dTDP-sugar epimerase

Authors : Cross, A.R.; Harmer, N.J.; Isupov, M.N.

Deposited on : 2021-10-04

Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	2021 (1.46-1.42)
Clashscore	141614	2086 (1.46-1.42)
Ramachandran outliers	138981	2047 (1.46-1.42)
Sidechain outliers	138945	2047 (1.46-1.42)
RSRZ outliers	127900	1993 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	AAA	211	79%	14% • 6%
1	BBB	211	76%	14% • 9%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called dTDP-4-dehydrorhamnose 3,5-epimerase.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace
1	AAA	199	Total	С	N	О	S	0	29	0
1	AAA	199	1794	1180	297	308	9	0	49	
1	BBB	191	Total	С	N	О	S	0	29	0
1	ррр	191	1746	1148	285	304	9	0	32	0

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-21	MET	-	initiating methionine	UNP Q83AP3
AAA	-20	HIS	-	expression tag	UNP Q83AP3
AAA	-19	HIS	-	expression tag	UNP Q83AP3
AAA	-18	HIS	-	expression tag	UNP Q83AP3
AAA	-17	HIS	-	expression tag	UNP Q83AP3
AAA	-16	HIS	-	expression tag	UNP Q83AP3
AAA	-15	HIS	-	expression tag	UNP Q83AP3
AAA	-14	SER	-	expression tag	UNP Q83AP3
AAA	-13	SER	-	expression tag	UNP Q83AP3
AAA	-12	GLY	-	expression tag	UNP Q83AP3
AAA	-11	VAL	-	expression tag	UNP Q83AP3
AAA	-10	ASP	-	expression tag	UNP Q83AP3
AAA	-9	LEU	-	expression tag	UNP Q83AP3
AAA	-8	GLY	-	expression tag	UNP Q83AP3
AAA	-7	THR	-	expression tag	UNP Q83AP3
AAA	-6	GLU	-	expression tag	UNP Q83AP3
AAA	-5	ASN	-	expression tag	UNP Q83AP3
AAA	-4	LEU	-	expression tag	UNP Q83AP3
AAA	-3	TYR	-	expression tag	UNP Q83AP3
AAA	-2	PHE	-	expression tag	UNP Q83AP3
AAA	-1	GLN	_	expression tag	UNP Q83AP3
AAA	0	SER	-	expression tag	UNP Q83AP3
BBB	-21	MET	-	initiating methionine	UNP Q83AP3
BBB	-20	HIS	-	expression tag	UNP Q83AP3
BBB	-19	HIS	-	expression tag	UNP Q83AP3



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	-18	HIS	-	expression tag	UNP Q83AP3
BBB	-17	HIS	-	expression tag	UNP Q83AP3
BBB	-16	HIS	-	expression tag	UNP Q83AP3
BBB	-15	HIS	-	expression tag	UNP Q83AP3
BBB	-14	SER	-	expression tag	UNP Q83AP3
BBB	-13	SER	-	expression tag	UNP Q83AP3
BBB	-12	GLY	-	expression tag	UNP Q83AP3
BBB	-11	VAL	-	expression tag	UNP Q83AP3
BBB	-10	ASP	-	expression tag	UNP Q83AP3
BBB	-9	LEU	-	expression tag	UNP Q83AP3
BBB	-8	GLY	-	expression tag	UNP Q83AP3
BBB	-7	THR	-	expression tag	UNP Q83AP3
BBB	-6	GLU	-	expression tag	UNP Q83AP3
BBB	-5	ASN	-	expression tag	UNP Q83AP3
BBB	-4	LEU	-	expression tag	UNP Q83AP3
BBB	-3	TYR	-	expression tag	UNP Q83AP3
BBB	-2	PHE	-	expression tag	UNP Q83AP3
BBB	-1	GLN	-	expression tag	UNP Q83AP3
BBB	0	SER	-	expression tag	UNP Q83AP3

• Molecule 2 is beta-D-xylopyranose (three-letter code: XYP) (formula: $C_5H_{10}O_5$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total C O 10 5 5	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	BBB	1	Total C O 10 5 5	0	0

• Molecule 3 is alpha-D-xylopyranose (three-letter code: XYS) (formula: $C_5H_{10}O_5$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total C O 10 5 5	0	0
3	BBB	1	Total C O 10 5 5	0	0

• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	3	Total Na 3 3	0	0
4	BBB	3	Total Na 3 3	0	0

• Molecule 5 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
5	BBB	1	Total 13	C 6	O 7	0	0

• Molecule 6 is water.


\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	226	Total O 226 226	0	0
6	BBB	227	Total O 227 227	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: dTDP-4-dehydrorhamnose 3,5-epimerase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	77.03Å 84.54Å 77.67Å	Depositor
a, b, c, α , β , γ	90.00° 113.59° 90.00°	Depositor
Resolution (Å)	51.39 - 1.43	Depositor
rtesolution (A)	71.18 - 1.43	EDS
% Data completeness	65.7 (51.39-1.43)	Depositor
(in resolution range)	65.7 (71.18-1.43)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.13 (at 1.43Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.167 , 0.199	Depositor
It, It free	0.172 , 0.207	DCC
R_{free} test set	2739 reflections (4.99%)	wwPDB-VP
Wilson B-factor (\mathring{A}^2)	18.5	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 50.5	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4052	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, XYS, NA, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	AAA	0.58	0/1914	0.79	0/2581	
1	BBB	0.59	0/1880	0.75	0/2534	
All	All	0.58	0/3794	0.77	0/5115	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1794	0	1889	34	1
1	BBB	1746	0	1828	36	0
2	AAA	10	0	0	0	0
2	BBB	10	0	0	0	0
3	AAA	10	0	10	0	0
3	BBB	10	0	10	0	0
4	AAA	3	0	0	0	0
4	BBB	3	0	0	0	0
5	BBB	13	0	5	0	0
6	AAA	226	0	0	13	0
6	BBB	227	0	0	15	0
All	All	4052	0	3742	68	1



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:57[B]:LYS:HE2	6:AAA:427:HOH:O	1.24	1.31
1:BBB:66[B]:GLN:NE2	6:BBB:303:HOH:O	1.58	1.29
1:BBB:80[B]:LEU:HD12	6:BBB:444:HOH:O	1.19	1.28
1:BBB:80[B]:LEU:CD1	6:BBB:444:HOH:O	1.71	1.24
1:AAA:112[B]:MET:HE1	6:AAA:464:HOH:O	0.99	1.16
1:AAA:112[A]:MET:SD	6:AAA:449:HOH:O	2.14	1.05
1:BBB:80[B]:LEU:CG	6:BBB:378:HOH:O	2.11	0.99
1:BBB:1[B]:MET:HG2	6:BBB:310:HOH:O	1.61	0.99
1:BBB:80[B]:LEU:CD2	6:BBB:378:HOH:O	2.10	0.98
1:BBB:80[B]:LEU:HG	6:BBB:378:HOH:O	1.64	0.98
1:AAA:112[A]:MET:CG	6:AAA:449:HOH:O	2.12	0.97
1:BBB:66[A]:GLN:OE1	1:BBB:140[A]:TYR:CD1	2.19	0.96
1:BBB:80[B]:LEU:HD21	6:BBB:378:HOH:O	1.68	0.94
1:AAA:80[B]:LEU:HD12	6:AAA:465:HOH:O	1.72	0.88
1:AAA:112[A]:MET:HG2	6:AAA:449:HOH:O	1.74	0.87
1:AAA:99[B]:LYS:NZ	6:AAA:302:HOH:O	2.06	0.86
1:AAA:1[B]:MET:HG2	1:AAA:2:PRO:HD2	1.59	0.84
1:BBB:145[B]:GLU:OE1	6:BBB:304:HOH:O	2.00	0.80
1:BBB:66[A]:GLN:OE1	1:BBB:140[A]:TYR:CE1	2.36	0.79
1:BBB:45[B]:GLU:OE2	6:BBB:305:HOH:O	2.01	0.79
1:AAA:83:VAL:HG11	1:AAA:132[B]:LEU:HD11	1.66	0.78
1:BBB:140[B]:TYR:CZ	1:BBB:142:PRO:HG3	2.20	0.77
1:BBB:93:GLU:OE1	6:BBB:302:HOH:O	2.05	0.73
1:AAA:83:VAL:CG1	1:AAA:132[B]:LEU:HD11	2.19	0.72
1:BBB:83:VAL:HG11	1:BBB:132[A]:LEU:HD21	1.76	0.68
1:AAA:80[B]:LEU:CD1	6:AAA:465:HOH:O	2.35	0.68
1:AAA:161[B]:ILE:HA	6:AAA:359:HOH:O	1.94	0.68
1:AAA:1[B]:MET:HG2	1:AAA:2:PRO:CD	2.26	0.64
1:BBB:66[A]:GLN:OE1	1:BBB:140[A]:TYR:HD1	1.77	0.63
1:BBB:169[B]:GLU:HG3	6:BBB:348:HOH:O	1.99	0.63
1:AAA:160[B]:PRO:O	1:AAA:161[B]:ILE:HB	1.98	0.62
1:BBB:165:LEU:HD23	1:BBB:167:LEU:HG	1.80	0.62
1:BBB:165:LEU:CD2	1:BBB:167:LEU:HG	2.30	0.62
1:BBB:1[B]:MET:SD	1:BBB:4[B]:GLU:HG2	2.42	0.60
1:BBB:80[B]:LEU:HD11	6:BBB:444:HOH:O	1.63	0.58
1:AAA:161[A]:ILE:HA	6:AAA:359:HOH:O	2.03	0.57
1:BBB:140[B]:TYR:CE2	1:BBB:142:PRO:HG3	2.39	0.56



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:82[B]:LYS:HE2	1:AAA:127:GLU:HG3	1.88	0.55
1:BBB:149:ARG:HG2	1:BBB:174:GLY:O	2.10	0.51
1:BBB:66[B]:GLN:HE21	1:BBB:72:GLN:H	1.58	0.50
1:BBB:188:GLY:HA3	6:BBB:376:HOH:O	2.11	0.50
1:BBB:66[B]:GLN:NE2	1:BBB:72:GLN:H	2.09	0.50
1:BBB:161:ILE:HG12	1:BBB:165:LEU:HD12	1.95	0.48
1:AAA:76:VAL:CG1	1:AAA:132[A]:LEU:HD11	2.44	0.48
1:BBB:82[B]:LYS:NZ	6:BBB:306:HOH:O	2.32	0.47
1:BBB:79[B]:VAL:HG11	1:BBB:133:LEU:HD11	1.95	0.47
1:AAA:1[B]:MET:CG	1:AAA:2:PRO:HD2	2.39	0.47
1:AAA:132[A]:LEU:HD23	1:AAA:134:TYR:CZ	2.50	0.47
1:AAA:161[B]:ILE:CG2	1:AAA:165:LEU:HD21	2.46	0.45
1:AAA:53[A]:LEU:HD23	1:AAA:131:HIS:CD2	2.50	0.45
1:AAA:132[A]:LEU:HD23	1:AAA:134:TYR:OH	2.16	0.45
1:AAA:79[B]:VAL:HG11	1:AAA:133:LEU:HD11	1.98	0.45
1:BBB:83:VAL:CG1	1:BBB:132[A]:LEU:HD21	2.46	0.45
1:AAA:112[B]:MET:CE	6:AAA:464:HOH:O	1.87	0.44
1:BBB:147:TYR:CE2	1:BBB:170[A]:LYS:HG2	2.52	0.44
1:AAA:4[B]:GLU:OE1	1:AAA:17[B]:LYS:HE2	2.18	0.43
1:AAA:79[A]:VAL:HG23	1:BBB:51:HIS:CE1	2.53	0.43
1:AAA:83:VAL:HG13	1:AAA:132[B]:LEU:HD11	1.98	0.42
1:BBB:33[B]:LYS:HD2	1:BBB:36[B]:ASP:OD1	2.20	0.42
1:BBB:53[B]:LEU:HD23	1:BBB:131:HIS:CD2	2.54	0.41
1:AAA:149:ARG:HG3	1:AAA:151:ASP:OD1	2.20	0.41
1:AAA:51:HIS:CE1	1:BBB:79[A]:VAL:HG23	2.55	0.41
1:AAA:123:MET:HE2	1:AAA:132[B]:LEU:HD13	2.03	0.41
1:AAA:130:THR:HG22	1:AAA:132[B]:LEU:HD12	2.02	0.41
1:BBB:80[B]:LEU:HA	1:BBB:107:SER:HB3	2.01	0.41
1:AAA:16[B]:ILE:HD13	1:AAA:37:PHE:CE2	2.56	0.40
1:AAA:179:ARG:NH2	6:AAA:320:HOH:O	2.52	0.40
1:AAA:39[A]:ARG:NH1	6:AAA:326:HOH:O	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:AAA:-2:PHE:CZ	1:AAA:6[A]:GLN:NE2[2 555]	2.18	0.02



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	AAA	226/211 (107%)	224 (99%)	0	2 (1%)	17	3
1	BBB	$221/211 \ (105\%)$	218 (99%)	3 (1%)	0	100	100
All	All	447/422 (106%)	442 (99%)	3 (1%)	2 (0%)	41	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	161[A]	ILE
1	AAA	161[B]	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	AAA	206/189 (109%)	202 (98%)	4 (2%)	57	22	
1	BBB	202/189 (107%)	197 (98%)	5 (2%)	47	13	
All	All	408/378 (108%)	399 (98%)	9 (2%)	60	17	

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	25	ARG
1	AAA	39[A]	ARG
1	AAA	39[B]	ARG
1	AAA	123	MET



Continued from previous page...

Mol	Chain	Res	Type
1	BBB	123[A]	MET
1	BBB	123[B]	MET
1	BBB	165	LEU
1	BBB	170[A]	LYS
1	BBB	170[B]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain Res		Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XYP	AAA	201	-	10,10,10	0.09	0	14,14,14	0.56	0
2	XYP	BBB	201	4	10,10,10	0.64	0	14,14,14	0.52	0
3	XYS	BBB	202	-	10,10,10	0.66	0	14,14,14	0.93	1 (7%)
3	XYS	AAA	202	-	10,10,10	0.47	0	14,14,14	0.58	0
5	FLC	BBB	203	-	12,12,12	1.05	1 (8%)	17,17,17	1.36	4 (23%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	AAA	201	-	-	-	0/1/1/1
2	XYP	BBB	201	4	-	-	0/1/1/1
3	XYS	BBB	202	-	-	-	0/1/1/1
3	XYS	AAA	202	-	-	-	0/1/1/1
5	FLC	BBB	203	-	-	0/16/16/16	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
5	BBB	203	FLC	CB-CBC	2.28	1.55	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
5	BBB	203	FLC	OB1-CBC-CB	-3.13	117.82	122.25
5	BBB	203	FLC	OG1-CGC-CG	-2.33	116.13	122.94
3	BBB	202	XYS	C1-C2-C3	2.32	115.12	110.31
5	BBB	203	FLC	OB2-CBC-CB	2.29	117.03	113.05
5	BBB	203	FLC	OA1-CAC-CA	-2.06	116.92	122.94

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	AAA	199/211 (94%)	0.05	8 (4%) 38 39	12, 20, 39, 64	0
1	BBB	191/211 (90%)	-0.07	5 (2%) 56 56	11, 18, 44, 77	0
All	All	390/422 (92%)	-0.01	13 (3%) 46 47	11, 18, 41, 77	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	-2	PHE	4.8
1	BBB	1[A]	MET	4.8
1	AAA	-9[A]	LEU	4.1
1	AAA	163[A]	ASN	3.8
1	AAA	161[A]	ILE	3.8
1	AAA	162	LYS	3.6
1	BBB	0	SER	3.5
1	BBB	24[A]	ASP	3.2
1	AAA	-3	TYR	3.1
1	AAA	164	ASN	2.8
1	BBB	162	LYS	2.8
1	AAA	165	LEU	2.6
1	AAA	187	HIS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	XYS	AAA	202	10/10	0.76	0.18	50,52,53,54	10
4	NA	AAA	204	1/1	0.86	0.47	53,53,53,53	0
4	NA	AAA	205	1/1	0.89	0.10	40,40,40,40	0
4	NA	BBB	204	1/1	0.89	0.15	44,44,44,44	0
2	XYP	BBB	201	10/10	0.93	0.12	20,24,27,29	0
3	XYS	BBB	202	10/10	0.94	0.11	27,30,34,35	0
2	XYP	AAA	201	10/10	0.95	0.08	20,22,23,24	10
5	FLC	BBB	203	13/13	0.96	0.10	20,21,23,25	0
4	NA	BBB	206	1/1	0.97	0.12	21,21,21,21	1
4	NA	AAA	203	1/1	0.99	0.22	38,38,38,38	0
4	NA	BBB	205	1/1	0.99	0.09	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

