



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2023 – 04:31 PM EDT

PDB ID : 6PXN  
Title : Human Casein Kinase 1 delta Tau mutant (R178C)  
Authors : Philpott, J.M.; Tripathi, S.M.; Partch, C.L.  
Deposited on : 2019-07-26  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	 5% (poor fit), 65% (0-1 outliers), 5% (2 outliers), 30% (3+ outliers or not modelled)
1	B	415	 4% (poor fit), 63% (0-1 outliers), 6% (2 outliers), 30% (3+ outliers or not modelled)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9771 atoms, of which 4714 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Casein kinase I isoform delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	290	4735	1519	2363	414	424	15	0	1	0
1	B	290	4714	1515	2351	410	423	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	CYS	ARG	engineered mutation	UNP P48730
B	178	CYS	ARG	engineered mutation	UNP P48730

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	152	Total	O	0	0
			152	152		
3	B	145	Total	O	0	0
			145	145		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.08Å 129.64Å 51.86Å 90.00° 113.41° 90.00°	Depositor
Resolution (Å)	44.68 – 1.55 45.96 – 1.55	Depositor EDS
% Data completeness (in resolution range)	97.6 (44.68-1.55) 97.7 (45.96-1.55)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 1.55Å)	Xtrriage
Refinement program	PHENIX (1.15.2_3472)	Depositor
R, $R_{free}$	0.177 , 0.205 0.179 , 0.206	Depositor DCC
$R_{free}$ test set	4363 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtrriage
Anisotropy	0.790	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 50.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9771	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2427	0.71	0/3263
1	B	0.52	0/2417	0.72	0/3248
All	All	0.52	0/4844	0.72	0/6511

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2372	2363	2364	20	0
1	B	2363	2351	2352	19	0
2	A	10	0	0	0	0
2	B	15	0	0	0	0
3	A	152	0	0	3	0
3	B	145	0	0	1	0
All	All	5057	4714	4716	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:VAL:HG11	1:B:79:VAL:HG11	1.61	0.80
1:B:63:VAL:CG1	1:B:144:LEU:HD11	2.10	0.80
1:B:63:VAL:HG12	1:B:144:LEU:HD11	1.66	0.77
1:A:28:ASP:OD2	1:A:31:ALA:HB3	1.90	0.71
1:A:216:LEU:HD13	1:A:224:LYS:CD	2.21	0.70
1:B:2:GLU:N	3:B:602:HOH:O	2.31	0.62
1:A:220:THR:HG23	1:A:223:GLN:OE1	2.00	0.61
1:A:216:LEU:HD13	1:A:224:LYS:HD3	1.83	0.61
1:A:216:LEU:HD13	1:A:224:LYS:HE2	1.84	0.59
1:A:28:ASP:OD2	1:A:31:ALA:CB	2.52	0.55
1:A:216:LEU:HD13	1:A:224:LYS:CE	2.37	0.54
1:B:223:GLN:HA	1:B:223:GLN:OE1	2.08	0.54
1:B:118:TYR:CZ	1:B:122:LYS:HE2	2.43	0.54
1:B:5:VAL:HG13	1:B:74:GLU:HG3	1.93	0.50
1:A:219:ALA:HB1	1:A:223:GLN:HB2	1.96	0.47
1:B:279:ARG:HH11	1:B:279:ARG:HG3	1.80	0.47
1:A:215:GLY:O	1:A:227:ARG:NE	2.48	0.46
1:A:216:LEU:HA	1:A:227:ARG:HE	1.80	0.46
1:B:130:LYS:HD3	1:B:176:THR:CG2	2.45	0.46
1:A:168:ARG:NE	3:A:607:HOH:O	2.49	0.45
1:B:130:LYS:HD3	1:B:176:THR:HG21	1.99	0.45
1:A:220:THR:O	1:A:221:LYS:HB2	2.16	0.45
1:A:28:ASP:CG	1:A:31:ALA:HB3	2.38	0.44
3:A:601:HOH:O	1:B:294:LYS:CG	2.66	0.44
1:B:152:LEU:HD22	1:B:174:THR:HG23	1.99	0.44
1:A:128:ASP:OD1	1:A:176:THR:HG23	2.18	0.43
1:A:231:LYS:HE3	3:A:647:HOH:O	2.19	0.43
1:A:227:ARG:HH11	1:A:227:ARG:HG2	1.84	0.42
1:A:68:ILE:C	1:A:68:ILE:HD12	2.40	0.42
1:A:215:GLY:O	1:A:216:LEU:HD23	2.20	0.42
1:A:25:LEU:HD11	1:A:34:GLU:OE1	2.20	0.42
1:B:186:LEU:HB2	1:B:188:ILE:HD12	2.01	0.42
1:B:128:ASP:OD1	1:B:176:THR:HG23	2.20	0.42
1:B:263:LYS:HE2	1:B:263:LYS:HB3	1.87	0.42
1:A:4:ARG:HB2	1:B:294:LYS:HZ3	1.84	0.41
1:B:230:GLU:O	1:B:234:SER:HB3	2.20	0.41
1:B:97:SER:O	1:B:99:LYS:HD2	2.21	0.40
1:B:182:ILE:HD12	1:B:256:ARG:CZ	2.51	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/415 (70%)	280 (97%)	9 (3%)	0	100	100
1	B	286/415 (69%)	279 (98%)	7 (2%)	0	100	100
All	All	575/830 (69%)	559 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/360 (71%)	250 (98%)	5 (2%)	55	26
1	B	254/360 (71%)	249 (98%)	5 (2%)	55	26
All	All	509/720 (71%)	499 (98%)	10 (2%)	55	26

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	THR
1	A	140	LYS
1	A	222	ARG
1	A	224	LYS
1	A	227	ARG
1	B	2	GLU
1	B	69	ARG
1	B	173	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	234	SER
1	B	279	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	60	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	503	-	4,4,4	0.15	0	6,6,6	0.20	0
2	SO4	A	502	-	4,4,4	0.19	0	6,6,6	0.18	0
2	SO4	A	501	-	4,4,4	0.24	0	6,6,6	0.50	0
2	SO4	B	502	-	4,4,4	0.32	0	6,6,6	0.47	0
2	SO4	B	501	-	4,4,4	0.17	0	6,6,6	0.23	0

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/415 (69%)	0.53	21 (7%) <b>15</b> <b>18</b>	16, 29, 73, 108	0
1	B	290/415 (69%)	0.27	17 (5%) <b>22</b> <b>26</b>	16, 28, 68, 93	0
All	All	580/830 (69%)	0.40	38 (6%) <b>18</b> <b>21</b>	16, 29, 71, 108	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	ALA	20.9
1	A	216	LEU	15.2
1	A	225	TYR	9.5
1	A	219	ALA	8.5
1	A	220	THR	8.4
1	B	225	TYR	7.6
1	B	219	ALA	7.1
1	B	161	THR	6.5
1	A	30	ALA	6.4
1	A	217	LYS	6.3
1	B	221	LYS	5.7
1	B	159	ALA	5.4
1	A	221	LYS	5.3
1	A	224	LYS	5.2
1	B	29	ILE	4.8
1	B	223	GLN	4.7
1	A	222	ARG	4.4
1	A	215	GLY	4.3
1	A	32	GLY	4.2
1	A	173	LEU	4.1
1	A	171	LYS	4.1
1	B	173	LEU	3.9
1	A	31	ALA	3.9
1	A	223	GLN	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	220	THR	3.7
1	A	6	GLY	3.6
1	A	227	ARG	3.3
1	A	29	ILE	3.3
1	B	222	ARG	3.3
1	A	228	ILE	2.9
1	B	224	LYS	2.8
1	B	5	VAL	2.6
1	B	73	ALA	2.5
1	A	140	LYS	2.5
1	B	172	ASN	2.4
1	B	71	CYS	2.2
1	B	7	ASN	2.2
1	B	174	THR	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	503	5/5	0.93	0.25	28,62,81,90	0
2	SO4	B	502	5/5	0.94	0.13	26,26,28,28	0
2	SO4	A	501	5/5	0.95	0.11	28,33,48,67	0
2	SO4	B	501	5/5	0.96	0.11	44,47,52,53	0
2	SO4	A	502	5/5	0.97	0.08	35,37,57,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SO4 B 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 A 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SO4 B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.