



wwPDB EM Validation Summary Report ⓘ

Nov 19, 2022 – 09:26 PM EST

PDB ID : 1Q5A
EMDB ID : EMD-1052
Title : S-shaped trans interactions of cadherins model based on fitting C-cadherin (1L3W) to 3D map of desmosomes obtained by electron tomography
Authors : He, W.; Cowin, P.; Stokes, D.L.
Deposited on : 2003-08-06
Resolution : 30.00 Å(reported)
Based on initial model : 1L3W

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 30.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	880	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: center;"> <p>61%</p> <p>17% 30% 10% . 39%</p> </div> </div>
1	B	880	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: center;"> <p>61%</p> <p>16% 30% 11% . 39%</p> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	801	-	-	X	-
2	NAG	A	805	X	-	X	-
2	NAG	A	806	X	-	X	-
2	NAG	A	807	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	809	-	-	X	-
2	NAG	A	810	-	-	X	-
2	NAG	A	902	X	-	X	-
2	NAG	A	903	X	-	-	-
2	NAG	A	904	-	-	X	-
2	NAG	B	801	-	-	X	-
2	NAG	B	805	X	-	X	-
2	NAG	B	806	X	-	X	-
2	NAG	B	807	-	-	X	-
2	NAG	B	809	-	-	X	-
2	NAG	B	810	-	-	X	-
2	NAG	B	902	X	-	X	-
2	NAG	B	903	X	-	-	-
2	NAG	B	904	-	-	X	-
3	NDG	A	811	-	-	X	-
3	NDG	B	811	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EP-cadherin.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	540	Total	C	N	O	S	0	0
			4191	2635	695	850	11		
1	B	540	Total	C	N	O	S	0	0
			4191	2635	695	850	11		

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
2	A	1	Total	C	N	O	0
			182	104	13	65	
2	A	1	Total	C	N	O	0
			182	104	13	65	
2	A	1	Total	C	N	O	0
			182	104	13	65	
2	A	1	Total	C	N	O	0
			182	104	13	65	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	A	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0
2	B	1	Total 182	C 104	N 13	O 65	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
2	B	1	182	104	13	65	0

- Molecule 3 is 2-acetamido-2-deoxy-alpha-D-glucopyranose (three-letter code: NDG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
3	A	1	28	16	2	10	0
3	A	1	28	16	2	10	0
3	B	1	28	16	2	10	0
3	B	1	28	16	2	10	0


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	AltConf
4	A	12	Total Ca 12 12	0
4	B	12	Total Ca 12 12	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EP-cadherin



G386	T446	D506	PHE	ARG	SER
I387	M447	Y507	LEU	ASN	LEU
V388	C448	S508	LEU	ASP	SER
T389	D449	I509	LEU	VAL	LEU
G390	Q450	Y510	LEU	VAL	LEU
N391	M451	V511	PHE	THR	SER
G392	P452	L512	LEU	LEU	SER
N393	E453	L513	LYS	MET	ASN
L394	P454	S514	ARG	ASN	ASN
D395	Q455	D515	LYS	ALA	ASN
R396	V456	A516	VAL	HIS	ASP
E397	L457	A517	VAL	THR	HIS
S398	T458	Q617	LYS	ARG	ASP
E399	I459	N618	GLU	PRO	TYR
Y400	S460	N619	PRO	ARG	ASN
V401	D461	P520	LEU	PRO	LEU
K402	A462	Q521	LEU	ASN	SER
M403	D463	L522	PRO	ASP	ASP
Y404	I464	T523	GLU	ASP	TRP
T405	P465	V524	ASP	ILE	SER
Y406	P466	V525	THR	GLY	ARG
T407	M467	N526	ARG	ASN	PHE
V408	T468	A527	PHE	ARG	ARG
I409	Y469	T528	ILE	ILE	ARG
M410	P470	V529	GLU	ASP	LYS
L411	Y471	C530	TYR	ASN	LEU
V412	K472	S531	GLY	LEU	ASP
T413	V473	C532	GLU	ALA	ALA
D414	S474	E533	GLY	ALA	GLY
D415	L475	G534	GLY	ASP	ASP
G416	S476	K535	GLY	ASP	ASP
V417	H477	A536	GLU	THR	GLU
S418	G478	I537	GLU	THR	THR
V419	S479	K538	ALA	ALA	ALA
G420	D480	C539	GLN	PRO	PRO
T421	L481	Q540	ASP	TYR	TYR
G422	T482	GLU	ASP	ASP	ASP
T423	W483	LYS	LEU	SER	SER
G424	K484	LEU	LEU	LEU	LEU
T425	A485	GLN	VAL	VAL	VAL
L426	E486	ASP	PHE	PHE	ASP
I427	L487	LEU	GLY	TYR	TYR
L428	D488	PRO	LEU	GLY	GLY
H429	S489	ILE	ARG	GLY	SER
V430	K490	ILE	ARG	SER	GLY
L431	G491	LEU	PRO	SER	GLY
D432	T492	VAL	ASP	GLU	GLU
V433	S493	ILE	ILE	ALA	ALA
M434	M494	LEU	MET	ALA	ALA
D435	N495	GLY			
M436	L496	LEU			
G437	S497	LEU			
P438	P498	LEU			
V439	T499	ILE			
P440	Q500	LEU			
S441	Q501	LEU			
P442	L502	LEU			
R443	K503	LEU			
V444	K504	LEU			
F445	G505	LEU			

4 Experimental information

Property	Value	Source
EM reconstruction method	TOMOGRAPHY	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of tilted images used	Not provided	
Resolution determination method	Not provided	
CTF correction method	no CTF correction. Imaging at underfocus 0.4 micron with CM200FEG microscope at 50,000 magnification	Depositor
Microscope	FEI/PHILIPS CM200FEG	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1200	Depositor
Minimum defocus (nm)	300	Depositor
Maximum defocus (nm)	500	Depositor
Magnification	68276	Depositor
Image detector	GENERIC GATAN	Depositor
Maximum voxel value	2603.000	Depositor
Minimum voxel value	-1866.000	Depositor
Average voxel value	1323.980	Depositor
Voxel value standard deviation	218.755	Depositor
Recommended contour level	1760.0	Depositor
Tomogram size (Å)	3720.19, 3720.19, 617.61	wwPDB
Tomogram dimensions	512, 512, 85	wwPDB
Tomogram angles (°)	90, 90, 90	wwPDB
Grid spacing (Å)	7.266, 7.266, 7.266	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	8/4276 (0.2%)	1.44	81/5839 (1.4%)
1	B	0.70	8/4276 (0.2%)	1.39	79/5839 (1.4%)
All	All	0.70	16/8552 (0.2%)	1.42	160/11678 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	4
1	B	0	4
All	All	1	8

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	335	ALA	CA-CB	-8.37	1.34	1.52
1	B	335	ALA	CA-CB	-8.33	1.34	1.52
1	A	539	CYS	CB-SG	8.17	1.96	1.82
1	B	539	CYS	CB-SG	8.15	1.96	1.82
1	B	223	PRO	CG-CD	7.02	1.73	1.50

The worst 5 of 160 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	490	LYS	N-CA-CB	-28.44	59.41	110.60
1	A	520	PRO	CA-C-N	-13.30	87.95	117.20
1	B	520	PRO	CA-C-N	-13.28	87.99	117.20
1	B	290	PHE	N-CA-C	12.73	145.38	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	ILE	N-CA-C	12.72	145.34	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	490	LYS	CA

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	17	PHE	Sidechain
1	A	18	PRO	Mainchain
1	A	222	ASP	Mainchain
1	A	520	PRO	Mainchain
1	B	17	PHE	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4191	0	4085	708	0
1	B	4191	0	4090	720	0
2	A	182	0	169	93	0
2	B	182	0	169	92	0
3	A	28	0	24	9	0
3	B	28	0	24	9	0
4	A	12	0	0	0	0
4	B	12	0	0	0	0
All	All	8826	0	8561	1393	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 80.

The worst 5 of 1393 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:464:ILE:HD12	1:B:465:PRO:CD	1.30	1.58
1:A:464:ILE:HD12	1:A:465:PRO:CD	1.30	1.55
1:A:87:PRO:CG	1:B:89:GLU:HB3	1.16	1.54
1:B:464:ILE:CD1	1:B:465:PRO:HD2	1.50	1.41
1:A:464:ILE:CD1	1:A:465:PRO:HD2	1.50	1.37

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/880 (61%)	401 (74%)	92 (17%)	45 (8%)	1	12
1	B	538/880 (61%)	401 (74%)	92 (17%)	45 (8%)	1	12
All	All	1076/1760 (61%)	802 (74%)	184 (17%)	90 (8%)	2	12

5 of 90 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	PRO
1	A	155	PRO
1	A	235	ILE
1	A	347	ARG
1	A	363	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	480/779 (62%)	381 (79%)	99 (21%)	1	7
1	B	480/779 (62%)	381 (79%)	99 (21%)	1	7
All	All	960/1558 (62%)	762 (79%)	198 (21%)	3	7

5 of 198 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	156	GLU
1	B	310	VAL
1	B	195	ASP
1	B	250	PRO
1	B	345	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	104	ASN
1	B	264	ASN
1	B	110	GLN
1	B	217	ASN
1	B	299	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 54 ligands modelled in this entry, 24 are monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	802	1	14,14,15	0.73	0	17,19,21	0.82	0
3	NDG	A	811	1	14,14,15	0.86	0	17,19,21	1.86	1 (5%)
3	NDG	B	804	1	14,14,15	0.63	0	17,19,21	0.78	0
2	NAG	A	904	1	14,14,15	0.72	1 (7%)	17,19,21	0.68	0
2	NAG	A	806	1	14,14,15	0.56	0	17,19,21	1.34	2 (11%)
2	NAG	B	812	1	14,14,15	0.81	1 (7%)	17,19,21	0.73	1 (5%)
2	NAG	B	801	1	14,14,15	0.68	0	17,19,21	0.95	1 (5%)
2	NAG	B	803	1	14,14,15	0.96	1 (7%)	17,19,21	1.12	2 (11%)
2	NAG	A	805	1	14,14,15	0.71	0	17,19,21	1.04	1 (5%)
2	NAG	A	810	1	14,14,15	0.66	0	17,19,21	1.34	4 (23%)
2	NAG	B	902	1	14,14,15	1.06	1 (7%)	17,19,21	1.08	1 (5%)
2	NAG	B	806	1	14,14,15	0.56	0	17,19,21	1.34	2 (11%)
2	NAG	A	801	1	14,14,15	0.68	0	17,19,21	0.94	1 (5%)
2	NAG	B	807	1	14,14,15	0.63	0	17,19,21	1.15	2 (11%)
2	NAG	A	903	1	14,14,15	0.53	0	17,19,21	0.76	0
2	NAG	B	808	1	14,14,15	0.66	0	17,19,21	0.69	0
2	NAG	B	904	1	14,14,15	0.72	1 (7%)	17,19,21	0.68	0
2	NAG	B	802	1	14,14,15	0.73	0	17,19,21	0.82	0
3	NDG	A	804	1	14,14,15	0.64	0	17,19,21	0.77	0
2	NAG	B	903	1	14,14,15	0.53	0	17,19,21	0.77	0
2	NAG	A	812	1	14,14,15	0.80	1 (7%)	17,19,21	0.73	1 (5%)
2	NAG	A	807	1	14,14,15	0.64	0	17,19,21	1.15	2 (11%)
2	NAG	A	803	1	14,14,15	0.96	1 (7%)	17,19,21	1.12	2 (11%)
2	NAG	B	805	1	14,14,15	0.71	0	17,19,21	1.05	1 (5%)
2	NAG	B	809	1	14,14,15	0.74	0	17,19,21	0.93	0
2	NAG	A	809	1	14,14,15	0.72	0	17,19,21	0.92	0
3	NDG	B	811	1	14,14,15	0.85	0	17,19,21	1.87	1 (5%)
2	NAG	B	810	1	14,14,15	0.65	0	17,19,21	1.34	4 (23%)
2	NAG	A	808	1	14,14,15	0.64	0	17,19,21	0.69	0
2	NAG	A	902	1	14,14,15	1.07	1 (7%)	17,19,21	1.08	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	802	1	-	2/6/23/26	0/1/1/1
3	NDG	A	811	1	-	2/6/23/26	0/1/1/1
3	NDG	B	804	1	-	0/6/23/26	0/1/1/1
2	NAG	A	904	1	-	3/6/23/26	0/1/1/1
2	NAG	A	806	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	B	812	1	-	4/6/23/26	0/1/1/1
2	NAG	B	801	1	-	3/6/23/26	0/1/1/1
2	NAG	B	803	1	-	2/6/23/26	0/1/1/1
2	NAG	A	805	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	A	810	1	-	3/6/23/26	0/1/1/1
2	NAG	B	902	1	1/1/5/7	1/6/23/26	0/1/1/1
2	NAG	B	806	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	A	801	1	-	3/6/23/26	0/1/1/1
2	NAG	B	807	1	-	5/6/23/26	0/1/1/1
2	NAG	A	903	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	B	808	1	-	3/6/23/26	0/1/1/1
2	NAG	B	904	1	-	3/6/23/26	0/1/1/1
2	NAG	B	802	1	-	2/6/23/26	0/1/1/1
3	NDG	A	804	1	-	0/6/23/26	0/1/1/1
2	NAG	B	903	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	A	812	1	-	4/6/23/26	0/1/1/1
2	NAG	A	807	1	-	5/6/23/26	0/1/1/1
2	NAG	B	805	1	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	A	803	1	-	2/6/23/26	0/1/1/1
2	NAG	B	809	1	-	2/6/23/26	0/1/1/1
2	NAG	A	809	1	-	2/6/23/26	0/1/1/1
3	NDG	B	811	1	-	2/6/23/26	0/1/1/1
2	NAG	B	810	1	-	3/6/23/26	0/1/1/1
2	NAG	A	808	1	-	3/6/23/26	0/1/1/1
2	NAG	A	902	1	1/1/5/7	1/6/23/26	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	902	NAG	C1-C2	3.09	1.57	1.52
2	B	902	NAG	C1-C2	3.02	1.56	1.52
2	B	803	NAG	O5-C5	2.51	1.48	1.43
2	A	803	NAG	O5-C5	2.51	1.48	1.43
2	B	904	NAG	C1-C2	-2.19	1.49	1.52

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	811	NDG	C2-N2-C7	-7.01	112.92	122.90
3	A	811	NDG	C2-N2-C7	-6.99	112.95	122.90
2	A	806	NAG	C2-N2-C7	-3.42	118.04	122.90
2	B	806	NAG	C2-N2-C7	-3.41	118.05	122.90
2	B	805	NAG	C2-N2-C7	-3.01	118.61	122.90

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	805	NAG	C1
2	A	806	NAG	C1
2	A	902	NAG	C1
2	A	903	NAG	C1
2	B	805	NAG	C1

5 of 72 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	807	NAG	C3-C2-N2-C7
2	A	902	NAG	C3-C2-N2-C7
2	A	904	NAG	C3-C2-N2-C7
2	B	807	NAG	C3-C2-N2-C7
2	B	902	NAG	C3-C2-N2-C7

There are no ring outliers.

26 monomers are involved in 203 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	811	NDG	7	0
3	B	804	NDG	2	0
2	A	904	NAG	8	0
2	A	806	NAG	12	0
2	B	812	NAG	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	NAG	21	0
2	B	803	NAG	4	0
2	A	805	NAG	7	0
2	A	810	NAG	13	0
2	B	902	NAG	8	0
2	B	806	NAG	12	0
2	A	801	NAG	21	0
2	B	807	NAG	16	0
2	B	808	NAG	2	0
2	B	904	NAG	8	0
3	A	804	NDG	2	0
2	A	812	NAG	3	0
2	A	807	NAG	17	0
2	A	803	NAG	4	0
2	B	805	NAG	7	0
2	B	809	NAG	8	0
2	A	809	NAG	8	0
3	B	811	NDG	7	0
2	B	810	NAG	13	0
2	A	808	NAG	2	0
2	A	902	NAG	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.


5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Tomogram visualisation [i](#)


This section contains visualisations of the EMDB entry EMD-1052. These allow visual inspection of the internal detail of the tomogram and identification of artifacts.

6.1 Orthogonal projections [i](#)



The images above show the tomogram projected in three orthogonal directions.

6.2 Central slices [i](#)





The images above show central slices of the tomogram in three orthogonal directions.

6.3 Largest variance slices [i](#)



X Index: 243

Y Index:
315

Z Index: 39

The images above show the largest variance slices of the tomogram in three orthogonal directions.


6.4 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Tomogram analysis [i](#)

This section contains the results of statistical analysis of the tomogram.

7.1 Voxel-value distribution [i](#)



The voxel-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-1052 and PDB model 1Q5A. Per-residue inclusion information can be found in section 3 on page 7.

8.1 Map-model overlay [i](#)

This section was not generated.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.







8.4 Atom inclusion [i](#)



At the recommended contour level, 0% of all backbone atoms, 0% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (1760.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.0008	 0.0060
A	 0.0016	 0.0080
B	 0.0000	 0.0030

