



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:24 pm BST

PDB ID : 6QEQ  
Title : PcfF from Enterococcus faecalis pCF10  
Authors : Rehman, S.; Berntsson, R.P.A.  
Deposited on : 2019-01-08  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
1	B	118	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 92%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
1	C	118	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 78%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey;"></div> </div>
1	D	118	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7569 atoms, of which 3724 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PcfF.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	Se			
1	A	111	1881	581	953	173	173	1	0	2	0
1	B	111	1866	578	944	170	173	1	0	0	0
1	C	105	1758	548	888	158	163	1	0	0	0
1	D	109	1852	575	939	167	170	1	0	1	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	51	Total	O	0	0
			51	51		
2	B	60	Total	O	0	0
			60	60		
2	C	39	Total	O	0	0
			39	39		
2	D	62	Total	O	0	0
			62	62		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PcfF




- Molecule 1: PcfF



- Molecule 1: PcfF



- Molecule 1: PcfF



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.85Å 61.48Å 89.12Å 90.00° 100.97° 90.00°	Depositor
Resolution (Å)	37.81 – 1.90 47.96 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.0 (37.81-1.90) 98.2 (47.96-1.76)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 1.76Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.202 , 0.248 0.207 , 0.253	Depositor DCC
$R_{free}$ test set	1076 reflections (2.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.6	Xtrriage
Anisotropy	0.162	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7569	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6098e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	0/946	0.78	0/1268
1	B	0.82	0/932	0.78	0/1248
1	C	0.74	0/880	0.76	0/1181
1	D	0.75	0/923	0.78	0/1237
All	All	0.78	0/3681	0.77	0/4934

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	13	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	928	953	938	14	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	922	944	944	2	0
1	C	870	888	888	17	0
1	D	913	939	938	4	0
2	A	51	0	0	1	0
2	B	60	0	0	0	0
2	C	39	0	0	3	0
2	D	62	0	0	2	0
All	All	3845	3724	3708	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:MSE:SE	1:C:93:LEU:HD11	2.28	0.84
1:A:75:MSE:HE1	1:C:75:MSE:CB	2.09	0.83
1:A:75:MSE:HE1	1:C:75:MSE:HB2	1.69	0.74
1:A:75:MSE:HE1	1:C:75:MSE:HB3	1.71	0.72
1:C:87:SER:OG	2:C:201:HOH:O	2.16	0.62
1:D:60:ARG:N	2:D:202:HOH:O	2.35	0.60
1:B:114:ARG:NH2	1:D:111:LEU:O	2.35	0.58
1:A:75:MSE:CE	1:C:75:MSE:HB3	2.34	0.57
1:A:104:VAL:O	1:A:108:LEU:HB2	2.05	0.56
1:A:75:MSE:CE	1:C:75:MSE:CB	2.83	0.54
1:C:13:ARG:NH2	2:C:203:HOH:O	2.41	0.54
1:A:91:LYS:HA	1:A:94:THR:HG22	1.92	0.52
1:C:57:GLU:HG3	1:C:111:LEU:HD13	1.91	0.52
1:A:72:ILE:HG21	1:A:97:VAL:HG21	1.98	0.46
1:A:75:MSE:SE	1:C:93:LEU:HD21	2.66	0.45
1:C:64:GLU:HA	1:C:67:LYS:HE2	1.99	0.44
1:A:75:MSE:SE	1:C:93:LEU:CD1	3.09	0.44
1:A:14:ILE:HD11	1:C:14:ILE:HG12	1.98	0.44
1:C:60:ARG:HG3	2:C:222:HOH:O	2.19	0.43
1:B:35:PRO:HG3	1:C:82:PHE:HA	2.01	0.43
1:D:107:GLU:O	1:D:110:LYS:HB3	2.19	0.42
1:A:91:LYS:O	1:A:94:THR:HG22	2.20	0.42
1:A:75:MSE:HE3	1:C:76:ALA:N	2.34	0.42
1:D:60:ARG:HB2	2:D:202:HOH:O	2.19	0.42
2:A:244:HOH:O	1:C:26:LYS:HG3	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/118 (94%)	109 (98%)	2 (2%)	0	100	100
1	B	109/118 (92%)	108 (99%)	1 (1%)	0	100	100
1	C	103/118 (87%)	103 (100%)	0	0	100	100
1	D	108/118 (92%)	108 (100%)	0	0	100	100
All	All	431/472 (91%)	428 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/110 (96%)	105 (99%)	1 (1%)	78	79
1	B	105/110 (96%)	104 (99%)	1 (1%)	76	76
1	C	99/110 (90%)	99 (100%)	0	100	100
1	D	104/110 (94%)	104 (100%)	0	100	100
All	All	414/440 (94%)	412 (100%)	2 (0%)	88	89

All (2) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	98	GLN
1	B	106	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	52	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	110/118 (93%)	0.72	11 (10%) <b>7</b> <b>8</b>	34, 65, 138, 179	0
1	B	110/118 (93%)	0.56	5 (4%) <b>33</b> <b>36</b>	33, 54, 112, 191	0
1	C	104/118 (88%)	0.86	9 (8%) <b>10</b> <b>11</b>	34, 80, 126, 192	0
1	D	108/118 (91%)	0.81	8 (7%) <b>14</b> <b>16</b>	32, 65, 130, 247	0
All	All	432/472 (91%)	0.74	33 (7%) <b>13</b> <b>15</b>	32, 66, 130, 247	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	111	LEU	8.1
1	D	5	GLN	7.7
1	A	7	LEU	5.8
1	C	60	ARG	4.9
1	B	117	GLN	4.8
1	A	83	HIS	4.6
1	B	8	LYS	4.6
1	D	70	ILE	4.5
1	A	117	GLN	4.2
1	D	7	LEU	4.2
1	D	4	LYS	4.0
1	C	93	LEU	3.8
1	C	85	ILE	3.8
1	A	88	GLU	3.7
1	B	113	LYS	3.5
1	D	112	ILE	3.3
1	C	50	ILE	3.1
1	C	49	GLU	2.8
1	A	92	ASP	2.8
1	D	67	LYS	2.7
1	C	97	VAL	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	114	ARG	2.5
1	A	100	LEU	2.4
1	C	70	ILE	2.4
1	C	110	LYS	2.4
1	A	99	SER	2.3
1	B	116	ASP	2.2
1	A	87	SER	2.2
1	D	60	ARG	2.2
1	A	104	VAL	2.2
1	C	67	LYS	2.1
1	A	95	ASP	2.1
1	B	105	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.