

wwPDB EM Validation Summary Report (i)

Jul 8, 2024 – 11:03 am BST

PDB ID : 7QEN

EMDB ID : EMD-13934

Title : S.c. Condensin core in DNA- and ATP-bound state Authors : Lecomte, L.; Hassler, M.; Haering, C.; Eustermann, S.

Deposited on : 2021-12-03

Resolution : 3.46 Å(reported)

Based on initial model : 6YVU

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $\mathrm{MapQ} \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$		
Clashscore	158937	4297		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length		Quality	of chain	
1	F	50	6%	66%	•	30%
2	G	50		66%		30%
3	В	1478	24%	·	72%	
4	A	1170	35%	5%	60%	
5	С	811	32%	·	64%	
6	D	1176	<u>-</u>	79%		10% 11%



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 19205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a DNA chain called DNA (35-MER).

Mol	Chain	Residues		A	toms			AltConf	Trace
1	F	35	Total 735	C 350	N 175	O 175	P 35	0	0

• Molecule 2 is a DNA chain called DNA (35-MER).

Mol	Chain	Residues		At	oms			AltConf	Trace
2	G	35	Total 700	C 350	N 70	O 245	P 35	0	0

• Molecule 3 is a protein called Structural maintenance of chromosomes protein 4.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			AltConf	Trace	
3	В	419	Total 3318	C 2100	N 570	O 634	S 14	0	0	Ì

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	14	ALA	SER	conflict	UNP Q12267
В	30	GLU	ASP	conflict	UNP Q12267
В	143	SER	ARG	conflict	UNP Q12267
В	425	ARG	LYS	conflict	UNP Q12267
В	546	ASP	ASN	conflict	UNP Q12267
В	697	ALA	VAL	conflict	UNP Q12267
В	704	ILE VAL		conflict	UNP Q12267
В	1028	ASN AS		conflict	UNP Q12267
В	1052 THR		ASN	conflict	UNP Q12267
В	B 1165 ASP		ALA	conflict	UNP Q12267
В	1177	VAL	ILE	conflict	UNP Q12267
В	1352	GLN	GLU	engineered mutation	UNP Q12267
В	1419	ARG	-	expression tag	UNP Q12267
В	1420	ILE	-	expression tag	UNP Q12267
В	1421	PRO	-	expression tag	UNP Q12267



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	1422	GLY	-	expression tag	UNP Q12267
В	1423	LEU	-	expression tag	UNP Q12267
В	1424	ILE	-	expression tag	UNP Q12267
В	1425	ASN	-	expression tag	UNP Q12267
В	1426	GLY	-	expression tag	UNP Q12267
В	1427	ALA	-	expression tag	UNP Q12267
В	1428	THR	-	expression tag	UNP Q12267
В	1429	GLY	-	expression tag	UNP Q12267
В	1430	TRP	-	expression tag	UNP Q12267
В	1431	SER	-	expression tag	UNP Q12267
В	1432	HIS	-	expression tag	UNP Q12267
В	1433	PRO	-	expression tag	UNP Q12267
В	1434	GLN	-	expression tag	UNP Q12267
В	1435	PHE	-	expression tag	UNP Q12267
В	1436	GLU	-	expression tag	UNP Q12267
В	1437	LYS	-	expression tag	UNP Q12267
В	1438	ALA	-	expression tag	UNP Q12267
В	1439	GLY	-	expression tag	UNP Q12267
В	1440	GLY	-	expression tag	UNP Q12267
В	1441	GLY	-	expression tag	UNP Q12267
В	1442	SER	-	expression tag	UNP Q12267
В	1443	GLY	-	expression tag	UNP Q12267
В	1444	GLY	-	expression tag	UNP Q12267
В	1445	GLY	-	expression tag	UNP Q12267
В	1446	SER	-	expression tag	UNP Q12267
В	1447	GLY	-	expression tag	UNP Q12267
В	1448	GLY	-	expression tag	UNP Q12267
В	1449	GLY	-	expression tag	UNP Q12267
В	1450	SER	-	expression tag	UNP Q12267
В	1451	TRP	-	expression tag	UNP Q12267
В	1452	SER	-	expression tag	UNP Q12267
В	1453	HIS	-	expression tag	UNP Q12267
В	1454	PRO	-	expression tag	UNP Q12267
В	1455	GLN	-	expression tag	UNP Q12267
В	1456	PHE	-	expression tag	UNP Q12267
В	1457	GLU	-	expression tag	UNP Q12267
В	1458	LYS	-	expression tag	UNP Q12267
В	1459	GLY	-	expression tag	UNP Q12267
В	1460	GLY	-	expression tag	UNP Q12267
В	1461	GLY	-	expression tag	UNP Q12267
В	1462	SER	-	expression tag	UNP Q12267
В	1463	GLY	-	expression tag	UNP Q12267



Continued from previous page						
	\mathcal{C}	lontinued	l fr	rom	previous	page

Chain	Residue	Modelled	Actual	Comment	Reference
В	1464	GLY	-	expression tag	UNP Q12267
В	1465	GLY	-	expression tag	UNP Q12267
В	1466	SER	-	expression tag	UNP Q12267
В	1467	GLY	-	expression tag	UNP Q12267
В	1468	GLY	-	expression tag	UNP Q12267
В	1469	GLY	-	expression tag	UNP Q12267
В	1470	SER	-	expression tag	UNP Q12267
В	1471	TRP	-	expression tag	UNP Q12267
В	1472	SER	-	expression tag	UNP Q12267
В	1473	HIS	-	expression tag	UNP Q12267
В	1474	PRO	-	expression tag	UNP Q12267
В	1475	GLN	-	expression tag	UNP Q12267
В	1476	PHE	-	expression tag	UNP Q12267
В	1477	GLU	-	expression tag	UNP Q12267
В	1478	LYS	-	expression tag	UNP Q12267

• Molecule 4 is a protein called Structural maintenance of chromosomes protein 2.

Mol	Chain	Residues		At	oms			AltConf	Trace
4	A	465	Total 3709	C 2351	N 639	O 702	S 17	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1113	GLN	GLU	engineered mutation	UNP P38989

• Molecule 5 is a protein called Condensin complex subunit 2.

Mol	Chain	Residues		At	oms			AltConf	Trace
5	С	291	Total 2345	C 1477	N 394	O 463	S 11	0	0

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	517	ALA	GLY	$\operatorname{conflict}$	UNP P38170
С	755	LEU	-	expression tag	UNP P38170
С	756	VAL	-	expression tag	UNP P38170
С	757	GLY	-	expression tag	UNP P38170
С	758	GLY	-	expression tag	UNP P38170
С	759	GLY	-	expression tag	UNP P38170



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	760	HIS	-	expression tag	UNP P38170
С	761	HIS	-	expression tag	UNP P38170
С	762	ARG	-	expression tag	UNP P38170
С	763	PRO	-	expression tag	UNP P38170
С	764	HIS	-	expression tag	UNP P38170
С	765	HIS	-	expression tag	UNP P38170
С	766	GLY	-	expression tag	UNP P38170
С	767	GLY	-	expression tag	UNP P38170
С	768	HIS	_	expression tag	UNP P38170
С	769	HIS	-	expression tag	UNP P38170
С	770	HIS	_	expression tag	UNP P38170
С	771	HIS	-	expression tag	UNP P38170
С	772	HIS	-	expression tag	UNP P38170
С	773	HIS	-	expression tag	UNP P38170
С	774	GLY	-	expression tag	UNP P38170
С	775	GLY	-	expression tag	UNP P38170
С	776	ARG	-	expression tag	UNP P38170
С	777	ILE	-	expression tag	UNP P38170
С	778	PHE	_	expression tag	UNP P38170
С	779	TYR	-	expression tag	UNP P38170
С	780	PRO	-	expression tag	UNP P38170
С	781	TYR	-	expression tag	UNP P38170
С	782	ASP	-	expression tag	UNP P38170
С	783	VAL	-	expression tag	UNP P38170
С	784	PRO	-	expression tag	UNP P38170
С	785	ASP	-	expression tag	UNP P38170
С	786	TYR	-	expression tag	UNP P38170
С	787	ALA	-	expression tag	UNP P38170
С	788	GLY	-	expression tag	UNP P38170
С	789	TYR	-	expression tag	UNP P38170
С	790	PRO	-	expression tag	UNP P38170
С	791	TYR	-	expression tag	UNP P38170
С	792	ASP	-	expression tag	UNP P38170
С	793	VAL	-	expression tag	UNP P38170
С	794	PRO	-	expression tag	UNP P38170
С	795	ASP	-	expression tag	UNP P38170
С	796	TYR	-	expression tag	UNP P38170
C	797	ALA	-	expression tag	UNP P38170
С	798	GLY	-	expression tag	UNP P38170
С	799	SER	-	expression tag	UNP P38170
С	800	TYR	-	expression tag	UNP P38170
С	801	PRO	-	expression tag	UNP P38170



~ · · 1	c		
Continued	trom	nremous	naae.

Chain	Residue	Modelled	Actual	Comment	Reference
С	802	TYR	-	expression tag	UNP P38170
С	803	ASP	-	expression tag	UNP P38170
С	804	VAL	-	expression tag	UNP P38170
С	805	PRO	_	expression tag	UNP P38170
С	806	ASN	-	expression tag	UNP P38170
С	807	TYR	_	expression tag	UNP P38170
С	808	ALA	-	expression tag	UNP P38170
С	809	ALA	-	expression tag	UNP P38170
С	810	GLY	-	expression tag	UNP P38170
С	811	HIS	-	expression tag	UNP P38170

• Molecule 6 is a protein called Condensin complex subunit 1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
6	D	1050	Total	С	N	О	S	0	0
0	ט	1050	8334	5317	1387	1581	49	U	U

• Molecule 7 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).

N	Iol	Chain	Residues		Ato	oms			AltConf
	7	B	1	Total	С	N	О	Р	O
	'	D			10	5	13	3	0
	7	Λ	1	Total	С	N	О	Р	0
	1	A	1	31	10	5	13	3	U




• Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
8	В	1	Total Mg 1 1	0
8	A	1	Total Mg 1 1	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.














• Molecule 5: Condensin complex subunit 2







 \bullet Molecule 6: Condensin complex subunit 1









4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	66205	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	1.0	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	2.048	Depositor
Minimum map value	-0.015	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.025	Depositor
Recommended contour level	0.05	Depositor
Map size (Å)	332.8, 332.8, 332.8	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.04, 1.04, 1.04	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	F	0.84	0/839	1.10	1/1291~(0.1%)	
2	G	0.87	0/769	1.53	0/1186	
3	В	0.32	0/3288	0.56	0/4416	
4	A	0.31	0/3758	0.53	0/5051	
5	С	0.29	0/2371	0.51	0/3184	
6	D	0.31	0/8456	0.55	1/11397~(0.0%)	
All	All	0.38	0/19481	0.66	$2/26525 \ (0.0\%)$	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
6	D	58	LEU	CA-CB-CG	5.46	127.86	115.30
1	F	16	DA	O4'-C1'-N9	-5.01	104.49	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	H(added)	Clashes	Symm-Clashes
1	F	735	0	386	1	0
2	G	700	0	421	2	0
3	В	3318	0	3292	34	0



, ,	omtomorod	trom	mmonia	maaa
	011.1.111111.111		previous	THEFT

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	3709	0	3839	35	0
5	С	2345	0	2340	27	0
6	D	8334	0	8562	69	0
7	A	31	0	12	1	0
7	В	31	0	12	0	0
8	A	1	0	0	0	0
8	В	1	0	0	0	0
All	All	19205	0	18864	149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
6:D:1025:ASP:H	6:D:1031:GLN:HE21	1.35	0.72
4:A:7:ILE:HB	4:A:80:THR:HB	1.77	0.65
3:B:242:GLU:HG2	3:B:244:SER:H	1.64	0.63
3:B:187:ASN:HD21	4:A:1118:LEU:HA	1.63	0.63
3:B:172:VAL:HG12	3:B:174:GLY:H	1.65	0.61

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percei	ntiles
3	В	399/1478 (27%)	388 (97%)	11 (3%)	0	100	100
4	A	461/1170 (39%)	450 (98%)	11 (2%)	0	100	100
5	С	279/811 (34%)	261 (94%)	18 (6%)	0	100	100
6	D	1038/1176 (88%)	999 (96%)	39 (4%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2177/4635 (47%)	2098 (96%)	79 (4%)	0	100 100	П

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
3	В	362/1321~(27%)	362 (100%)	0	100	100	
4	A	418/1071~(39%)	416 (100%)	2 (0%)	88	95	
5	С	267/731~(36%)	265 (99%)	2 (1%)	84	93	
6	D	955/1072~(89%)	953 (100%)	2 (0%)	93	98	
All	All	$2002/4195\ (48\%)$	1996 (100%)	6 (0%)	92	98	

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	С	211	ASP
6	D	720	LYS
6	D	1032	ARG
4	A	980	LYS
4	A	218	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
5	С	314	GLN
6	D	14	GLN
6	D	1031	GLN
5	С	738	ASN
6	D	70	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
7	ATP	В	2000	8	26,33,33	0.92	1 (3%)	31,52,52	1.58	5 (16%)	
7	ATP	A	2000	8	26,33,33	0.90	1 (3%)	31,52,52	1.52	5 (16%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ATP	В	2000	8	-	3/18/38/38	0/3/3/3
7	ATP	A	2000	8	-	7/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
7	В	2000	ATP	C5-C4	2.30	1.47	1.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
7	A	2000	ATP	C5-C4	2.22	1.46	1.40

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
7	В	2000	ATP	PB-O3B-PG	-3.47	120.93	132.83
7	В	2000	ATP	PA-O3A-PB	-3.45	120.98	132.83
7	В	2000	ATP	N3-C2-N1	-3.43	123.32	128.68
7	A	2000	ATP	N3-C2-N1	-3.34	123.46	128.68
7	A	2000	ATP	PA-O3A-PB	-3.26	121.64	132.83

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	2000	ATP	C5'-O5'-PA-O1A
7	A	2000	ATP	C5'-O5'-PA-O2A
7	A	2000	ATP	C3'-C4'-C5'-O5'
7	В	2000	ATP	O4'-C4'-C5'-O5'
7	В	2000	ATP	C3'-C4'-C5'-O5'


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	2000	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-13934. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map



6.2.2 Raw map




The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




6.6 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_13934_msk_1.map (i)








7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $135~\mathrm{nm}^3$; this corresponds to an approximate mass of $122~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.289 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.289 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.46	-	-
Author-provided FSC curve	3.44	4.00	3.55
Unmasked-calculated*	4.09	7.12	4.20


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.09 differs from the reported value 3.46 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-13934 and PDB model 7QEN. Per-residue inclusion information can be found in section 3 on page 9.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).



9.4 Atom inclusion (i)



At the recommended contour level, 92% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8280	0.4110
A	0.8290	0.4260
В	0.8390	0.4500
С	0.7940	0.4130
D	0.8320	0.4030
F	0.8160	0.3130
G	0.8530	0.3460



