



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2023 – 03:20 PM EDT

PDB ID : 2QG4  
Title : Crystal structure of human UDP-glucose dehydrogenase product complex with UDP-glucuronate  
Authors : Kavanagh, K.L.; Guo, K.; Bunkoczi, G.; Savitsky, P.; Pilka, E.; Bhatia, C.; Niesen, F.; Smee, C.; Berridge, G.; von Delft, F.; Weigelt, J.; Arrowsmith, C.H.; Sundstrom, M.; Edwards, A.; Oppermann, U.; Structural Genomics Consortium (SGC)  
Deposited on : 2007-06-28  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)






The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	96%
1	B	467	95%
1	C	467	94%  5%

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.35

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	D	467	 94% . .
1	E	467	 93% 6% .
1	F	467	 96% . .
1	G	467	 94% . .
1	H	467	 96% . .

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 32149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-glucose 6-dehydrogenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	460	3559	2254	612	674	19	0	0	0
1	B	461	3595	2280	614	682	19	0	3	0
1	C	459	3528	2240	598	671	19	0	0	0
1	D	461	3590	2273	613	685	19	0	2	0
1	E	459	3541	2245	607	670	19	0	0	0
1	F	461	3592	2277	614	682	19	0	2	0
1	G	460	3547	2249	604	675	19	0	0	0
1	H	461	3594	2275	615	685	19	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	cloning artifact	UNP O60701
B	0	SER	-	cloning artifact	UNP O60701
C	0	SER	-	cloning artifact	UNP O60701
D	0	SER	-	cloning artifact	UNP O60701
E	0	SER	-	cloning artifact	UNP O60701
F	0	SER	-	cloning artifact	UNP O60701
G	0	SER	-	cloning artifact	UNP O60701
H	0	SER	-	cloning artifact	UNP O60701

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	B	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	C	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	D	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	E	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	F	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	G	1	Total 37	C 15	N 2	O 18	P 2	0	0
3	H	1	Total 37	C 15	N 2	O 18	P 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total 2 Cl 2	0	0
4	D	2	Total 2 Cl 2	0	0
4	F	2	Total 2 Cl 2	0	0
4	H	2	Total 2 Cl 2	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0
5	G	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	388	Total O 388 388	0	0
6	B	428	Total O 428 428	0	0
6	C	292	Total O 292 292	0	0
6	D	410	Total O 410 410	0	0
6	E	285	Total O 285 285	0	0
6	F	410	Total O 410 410	0	0
6	G	341	Total O 341 341	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	H	445	Total 445	O 445	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-glucose 6-dehydrogenase

Chain A:  96%



- Molecule 1: UDP-glucose 6-dehydrogenase

Chain B:  95%



- Molecule 1: UDP-glucose 6-dehydrogenase

Chain C:  94% 5%




- Molecule 1: UDP-glucose 6-dehydrogenase

Chain D:  94%




- Molecule 1: UDP-glucose 6-dehydrogenase

Chain E:  93% 6%



- Molecule 1: UDP-glucose 6-dehydrogenase

Chain F:  96% ..




- Molecule 1: UDP-glucose 6-dehydrogenase

Chain G:  94% ..



- Molecule 1: UDP-glucose 6-dehydrogenase

Chain H:  96% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	193.87Å 193.87Å 352.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.66 – 2.10 59.70 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (59.66-2.10) 100.0 (59.70-2.10)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.3.0037	Depositor
R, $R_{free}$	0.176 , 0.225 0.178 , 0.225	Depositor DCC
$R_{free}$ test set	14848 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtrriage
Anisotropy	0.056	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 30.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.478 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	32149	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3730e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UGA, NAD, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/3623	0.58	0/4910
1	B	0.50	0/3668	0.62	0/4967
1	C	0.45	0/3592	0.58	0/4872
1	D	0.49	0/3660	0.59	0/4957
1	E	0.47	0/3605	0.58	0/4887
1	F	0.52	0/3662	0.61	1/4959 (0.0%)
1	G	0.44	0/3611	0.58	1/4896 (0.0%)
1	H	0.51	0/3661	0.61	0/4957
All	All	0.48	0/29082	0.59	2/39405 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	298	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	F	41	ARG	NE-CZ-NH2	5.19	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3559	0	3530	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3595	0	3587	10	0
1	C	3528	0	3478	11	0
1	D	3590	0	3570	11	0
1	E	3541	0	3513	15	0
1	F	3592	0	3583	5	0
1	G	3547	0	3504	10	0
1	H	3594	0	3580	6	0
2	A	27	0	12	0	0
2	B	44	0	26	0	0
2	C	27	0	12	0	0
2	D	44	0	26	0	0
2	E	27	0	12	0	0
2	F	44	0	26	0	0
2	G	27	0	12	0	0
2	H	44	0	26	0	0
3	A	37	0	19	0	0
3	B	37	0	19	0	0
3	C	37	0	19	0	0
3	D	37	0	19	0	0
3	E	37	0	19	0	0
3	F	37	0	19	0	0
3	G	37	0	19	0	0
3	H	37	0	19	0	0
4	B	2	0	0	0	0
4	D	2	0	0	0	0
4	F	2	0	0	0	0
4	H	2	0	0	0	0
5	B	4	0	6	0	0
5	C	4	0	6	0	0
5	F	4	0	6	0	0
5	G	4	0	6	0	0
6	A	388	0	0	0	0
6	B	428	0	0	2	0
6	C	292	0	0	0	0
6	D	410	0	0	0	0
6	E	285	0	0	1	0
6	F	410	0	0	0	0
6	G	341	0	0	2	0
6	H	445	0	0	0	0
All	All	32149	0	28673	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:7:ILE:HG23	1:G:84:LEU:HD23	1.82	0.61
1:G:129:LYS:NZ	6:G:5306:HOH:O	2.35	0.60
1:H:133:PRO:HB3	1:H:287:LEU:HD21	1.84	0.60
1:E:452:LEU:HB3	1:E:459:ILE:HD11	1.84	0.59
1:G:89:VAL:HG11	1:G:112:CYS:SG	2.44	0.57

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/467 (98%)	445 (98%)	11 (2%)	0	100	100
1	B	460/467 (98%)	451 (98%)	9 (2%)	0	100	100
1	C	455/467 (97%)	445 (98%)	10 (2%)	0	100	100
1	D	459/467 (98%)	450 (98%)	9 (2%)	0	100	100
1	E	455/467 (97%)	447 (98%)	8 (2%)	0	100	100
1	F	459/467 (98%)	450 (98%)	9 (2%)	0	100	100
1	G	456/467 (98%)	445 (98%)	11 (2%)	0	100	100
1	H	458/467 (98%)	450 (98%)	8 (2%)	0	100	100
All	All	3658/3736 (98%)	3583 (98%)	75 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/401 (95%)	378 (99%)	3 (1%)	81	86
1	B	388/401 (97%)	386 (100%)	2 (0%)	88	92
1	C	374/401 (93%)	372 (100%)	2 (0%)	88	92
1	D	388/401 (97%)	386 (100%)	2 (0%)	88	92
1	E	379/401 (94%)	376 (99%)	3 (1%)	81	86
1	F	388/401 (97%)	387 (100%)	1 (0%)	92	95
1	G	378/401 (94%)	376 (100%)	2 (0%)	88	92
1	H	389/401 (97%)	387 (100%)	2 (0%)	88	92
All	All	3065/3208 (96%)	3048 (99%)	17 (1%)	86	90

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	177	ARG
1	H	177	ARG
1	D	177	ARG
1	D	370	LYS
1	E	129	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	259	GLN
1	D	324	ASN
1	G	229	GLN
1	F	229	GLN
1	C	229	GLN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 8 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	C	500	-	24,29,48	1.27	2 (8%)	29,45,73	1.45	4 (13%)
3	UGA	F	501	-	32,39,39	1.27	2 (6%)	42,60,60	0.99	3 (7%)
3	UGA	G	501	-	32,39,39	1.51	3 (9%)	42,60,60	1.04	3 (7%)
5	EDO	C	5002	-	3,3,3	0.32	0	2,2,2	0.48	0
2	NAD	H	500	-	42,48,48	1.70	4 (9%)	50,73,73	1.26	4 (8%)
2	NAD	E	500	-	24,29,48	1.28	2 (8%)	29,45,73	1.50	4 (13%)
3	UGA	A	501	-	32,39,39	1.56	3 (9%)	42,60,60	1.01	3 (7%)
2	NAD	D	500	-	42,48,48	1.71	4 (9%)	50,73,73	1.22	4 (8%)
3	UGA	D	501	-	32,39,39	1.43	3 (9%)	42,60,60	1.02	2 (4%)
5	EDO	B	5001	-	3,3,3	0.23	0	2,2,2	0.63	0
2	NAD	A	500	-	24,29,48	1.33	3 (12%)	29,45,73	1.51	5 (17%)
3	UGA	E	501	-	32,39,39	1.75	3 (9%)	42,60,60	1.26	3 (7%)
2	NAD	G	500	-	24,29,48	1.28	2 (8%)	29,45,73	1.38	2 (6%)
3	UGA	H	501	-	32,39,39	1.57	3 (9%)	42,60,60	0.93	1 (2%)
5	EDO	F	5003	-	3,3,3	0.31	0	2,2,2	0.41	0
3	UGA	C	501	-	32,39,39	1.46	3 (9%)	42,60,60	1.09	5 (11%)
5	EDO	G	5004	-	3,3,3	0.37	0	2,2,2	0.46	0
3	UGA	B	501	-	32,39,39	1.48	4 (12%)	42,60,60	0.86	1 (2%)
2	NAD	F	500	-	42,48,48	1.86	4 (9%)	50,73,73	1.19	2 (4%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	B	500	-	42,48,48	1.75	4 (9%)	50,73,73	1.20	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	C	500	-	-	4/12/32/62	0/3/3/5
3	UGA	F	501	-	-	1/23/61/61	0/3/3/3
3	UGA	G	501	-	-	3/23/61/61	0/3/3/3
5	EDO	C	5002	-	-	1/1/1/1	-
2	NAD	H	500	-	-	5/26/62/62	0/5/5/5
2	NAD	E	500	-	-	1/12/32/62	0/3/3/5
3	UGA	A	501	-	-	2/23/61/61	0/3/3/3
2	NAD	D	500	-	-	5/26/62/62	0/5/5/5
3	UGA	D	501	-	-	4/23/61/61	0/3/3/3
5	EDO	B	5001	-	-	1/1/1/1	-
2	NAD	A	500	-	-	3/12/32/62	0/3/3/5
3	UGA	E	501	-	-	4/23/61/61	0/3/3/3
2	NAD	G	500	-	-	2/12/32/62	0/3/3/5
3	UGA	H	501	-	-	3/23/61/61	0/3/3/3
5	EDO	F	5003	-	-	1/1/1/1	-
3	UGA	C	501	-	-	3/23/61/61	0/3/3/3
5	EDO	G	5004	-	-	0/1/1/1	-
3	UGA	B	501	-	-	4/23/61/61	0/3/3/3
2	NAD	F	500	-	-	5/26/62/62	0/5/5/5
2	NAD	B	500	-	-	5/26/62/62	0/5/5/5

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	500	NAD	O7N-C7N	9.37	1.42	1.24
2	B	500	NAD	O7N-C7N	9.03	1.41	1.24
2	H	500	NAD	O7N-C7N	8.52	1.40	1.24
2	D	500	NAD	O7N-C7N	8.46	1.40	1.24
3	E	501	UGA	O4D-C1D	5.42	1.48	1.41

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	500	NAD	N3A-C2A-N1A	-5.63	119.88	128.68
2	A	500	NAD	N3A-C2A-N1A	-5.62	119.90	128.68
2	C	500	NAD	N3A-C2A-N1A	-5.56	119.98	128.68
2	B	500	NAD	N3A-C2A-N1A	-5.55	120.00	128.68
2	H	500	NAD	N3A-C2A-N1A	-5.53	120.04	128.68

There are no chirality outliers.



5 of 57 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	500	NAD	O4D-C1D-N1N-C2N
2	B	500	NAD	O4D-C1D-N1N-C6N
2	B	500	NAD	C2D-C1D-N1N-C2N
2	B	500	NAD	C2D-C1D-N1N-C6N
2	D	500	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	460/467 (98%)	-0.68	1 (0%) 95   95	9, 19, 34, 51	0
1	B	461/467 (98%)	-0.78	0 100   100	7, 13, 25, 46	0
1	C	459/467 (98%)	-0.67	0 100   100	8, 18, 33, 51	0
1	D	461/467 (98%)	-0.79	0 100   100	5, 13, 26, 47	0
1	E	459/467 (98%)	-0.68	0 100   100	9, 18, 34, 51	0
1	F	461/467 (98%)	-0.78	0 100   100	5, 11, 24, 43	0
1	G	460/467 (98%)	-0.69	0 100   100	9, 18, 34, 51	0
1	H	461/467 (98%)	-0.79	0 100   100	6, 12, 25, 46	0
All	All	3682/3736 (98%)	-0.73	1 (0%) 100   100	5, 15, 30, 51	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	98	MET	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAD	C	500	27/44	0.97	0.09	14,23,32,40	0
2	NAD	G	500	27/44	0.97	0.08	13,25,35,39	0
5	EDO	B	5001	4/4	0.97	0.09	6,7,11,15	0
3	UGA	A	501	37/37	0.98	0.08	13,22,32,36	0
3	UGA	C	501	37/37	0.98	0.07	9,17,25,27	0
3	UGA	E	501	37/37	0.98	0.08	10,19,28,35	0
2	NAD	E	500	27/44	0.98	0.07	7,16,23,28	0
5	EDO	C	5002	4/4	0.98	0.12	8,13,15,16	0
5	EDO	F	5003	4/4	0.98	0.10	6,11,14,14	0
5	EDO	G	5004	4/4	0.98	0.07	7,9,11,14	0
2	NAD	B	500	44/44	0.99	0.07	3,11,15,17	0
3	UGA	D	501	37/37	0.99	0.06	4,11,16,17	0
2	NAD	F	500	44/44	0.99	0.07	2,10,13,17	0
3	UGA	F	501	37/37	0.99	0.06	2,9,14,16	0
3	UGA	G	501	37/37	0.99	0.06	4,18,23,30	0
3	UGA	H	501	37/37	0.99	0.06	3,10,17,18	0
4	CL	B	4002	1/1	0.99	0.08	10,10,10,10	0
4	CL	B	4005	1/1	0.99	0.03	21,21,21,21	0
4	CL	D	4006	1/1	0.99	0.03	22,22,22,22	0
4	CL	F	4007	1/1	0.99	0.03	22,22,22,22	0
4	CL	H	4004	1/1	0.99	0.07	10,10,10,10	0
4	CL	H	4008	1/1	0.99	0.03	22,22,22,22	0
2	NAD	A	500	27/44	0.99	0.06	9,18,25,27	0
2	NAD	H	500	44/44	0.99	0.07	5,9,15,17	0
2	NAD	D	500	44/44	0.99	0.06	2,10,15,15	0
3	UGA	B	501	37/37	0.99	0.06	6,11,17,20	0
4	CL	F	4003	1/1	1.00	0.07	11,11,11,11	0
4	CL	D	4001	1/1	1.00	0.07	10,10,10,10	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NAD C 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD G 500:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around UGA A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UGA C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UGA E 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAD E 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD B 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UGA D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD F 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UGA F 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UGA G 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UGA H 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD A 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAD H 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD D 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UGA B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.