



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2024 – 02:08 pm GMT

PDB ID : 6QGT
Title : The carbon monoxide inhibition of F420-reducing [NiFe] hydrogenase complex from *Methanosarcina barkeri*
Authors : Iina, Y.; Lorent, C.; Katz, S.; Jeoung, J.H.; Shima, S.; Horch, M.; Zebger, I.; Dobbek, H.
Deposited on : 2019-01-12
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	253	 5% 91% 9%
2	A	438	 % 92% 8%
3	B	291	 % 94% 6%

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 8237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coenzyme F420 hydrogenase subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	G	252	1911	1200	322	366	23	0	0	0

- Molecule 2 is a protein called F420-reducing hydrogenase, subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	437	3421	2154	609	631	27	0	3	0

- Molecule 3 is a protein called Coenzyme F420 hydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	B	291	2271	1442	390	425	14	0	1	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	265	GLN	GLU	conflict	UNP A0A0E3QWH3

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total	Fe S	0	0
			8	4 4		
4	G	1	Total	Fe S	0	0
			8	4 4		
4	G	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		

- Molecule 5 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	Fe	S	0	0
			3	1	2		

- Molecule 6 is TRIS-HYDROXYMETHYL-METHYL-AMMONIUM (three-letter code: 144) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	G	1	Total	C	N	O	0	0
			8	4	1	3		
6	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is (R,R)-2,3-BUTANEDIOL (three-letter code: BU3) (formula: $C_4H_{10}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	G	1	Total C O 6 4 2	0	0
7	G	1	Total C O 6 4 2	0	0
7	B	1	Total C O 6 4 2	0	0

- Molecule 8 is dicyano-(oxidaniumylidynemethylnickelio)-(oxidanylidenemethylidene)iron (three-letter code: J52) (formula: $C_4FeN_2NiO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Fe	N	Ni			O
8	A	1	10	4	1	2	1	2	0	0


- Molecule 9 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Fe 1 1	0	0

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total Mg 1 1	0	0

- Molecule 11 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total C O 8 6 2	0	0

- Molecule 12 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
12	B	1	53	27	9	15	2	0	0


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	G	102	Total	O	0	2
			104	104		
13	A	257	Total	O	0	3
			260	260		
13	B	127	Total	O	0	1
			128	128		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Coenzyme F420 hydrogenase subunit gamma



- Molecule 2: F420-reducing hydrogenase, subunit alpha



- Molecule 3: Coenzyme F420 hydrogenase subunit beta



4 Data and refinement statistics i

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, α , β , γ	235.43Å 235.43Å 235.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.06 – 1.99 48.06 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.06-1.99) 99.9 (48.06-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.98Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.169 , 0.209 0.169 , 0.209	Depositor DCC
R_{free} test set	2100 reflections (2.84%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.034 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8237	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FAD, 144, BU3, FE, FES, J52, MPD, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	G	0.51	0/1943	0.64	0/2626
2	A	0.53	1/3502 (0.0%)	0.63	0/4757
3	B	0.46	0/2315	0.60	0/3125
All	All	0.50	1/7760 (0.0%)	0.63	0/10508

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	435	CYS	CB-SG	-5.81	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1911	0	1893	18	0
2	A	3421	0	3394	19	0
3	B	2271	0	2305	9	0
4	B	8	0	0	0	0
4	G	24	0	0	1	0
5	G	3	0	0	0	0
6	A	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	G	8	0	11	3	0
7	B	6	0	10	0	0
7	G	12	0	20	3	0
8	A	10	0	0	2	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
11	A	8	0	14	0	0
12	B	53	0	30	1	0
13	A	260	0	0	1	0
13	B	128	0	0	1	0
13	G	104	0	0	2	0
All	All	8237	0	7689	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:153:ARG:HH12	7:G:306:BU3:H43	1.50	0.77
2:A:348[B]:ASN:ND2	13:A:601:HOH:O	2.23	0.71
1:G:89:ASP:OD1	6:G:305:144:H13	1.93	0.69
1:G:153:ARG:HH12	7:G:306:BU3:C4	2.07	0.68
3:B:263:ALA:O	3:B:267:ILE:HG12	1.96	0.64
6:G:305:144:H42	2:A:9:PRO:HD3	1.81	0.62
2:A:148:ILE:HG23	2:A:186:LEU:HB3	1.82	0.60
2:A:148:ILE:HG21	2:A:190:CYS:HB2	1.84	0.59
2:A:301:VAL:HG13	2:A:408:PRO:HG3	1.85	0.58
3:B:137:PHE:CD2	3:B:213:GLY:HA3	2.40	0.56
3:B:199:VAL:HG22	3:B:274:VAL:HG22	1.88	0.54
3:B:146:SER:HB3	3:B:192:TYR:HB3	1.90	0.51
1:G:22:LYS:NZ	13:G:406:HOH:O	2.43	0.50
2:A:14:GLU:HB3	2:A:432:CYS:HA	1.93	0.50
3:B:39:TYR:OH	3:B:237:LYS:HG2	2.12	0.50
1:G:142:ILE:HG23	1:G:151:LEU:HD23	1.94	0.49
1:G:65:ARG:O	6:G:305:144:H41	2.12	0.49
2:A:119:ILE:HD12	2:A:124:LYS:HD3	1.94	0.49
3:B:250:GLU:H	3:B:250:GLU:CD	2.16	0.47
2:A:280:PRO:HA	2:A:281:ILE:HA	1.66	0.46
1:G:266:LYS:HB3	1:G:266:LYS:HE2	1.56	0.45
1:G:24:GLY:HA3	1:G:70:MET:SD	2.56	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:153:ARG:NH1	7:G:306:BU3:H43	2.26	0.45
4:G:301:SF4:S4	2:A:61:ARG:HG2	2.57	0.45
2:A:402:PRO:HG2	8:A:501:J52:C2	2.46	0.44
2:A:432:CYS:HB3	2:A:435:CYS:HB2	1.99	0.44
3:B:9:LYS:HB2	3:B:9:LYS:HE2	1.64	0.44
2:A:97:LEU:HD13	2:A:157:GLY:HA3	1.99	0.44
2:A:328:THR:HG23	2:A:428:ALA:HB2	2.00	0.43
1:G:27:HIS:CD2	13:G:429:HOH:O	2.71	0.43
1:G:257:PHE:HA	3:B:118:LEU:HD11	2.01	0.43
3:B:250:GLU:HA	13:B:489:HOH:O	2.19	0.43
1:G:80:CYS:HB2	1:G:129:TYR:CE2	2.54	0.42
1:G:206:MET:CE	1:G:245:PRO:HG2	2.49	0.42
1:G:252:ASP:OD1	1:G:252:ASP:N	2.53	0.42
1:G:34:CYS:SG	1:G:104:GLY:HA3	2.59	0.42
2:A:317:LEU:HB3	2:A:323:PHE:HB2	2.02	0.42
2:A:430:ASP:N	2:A:431:PRO:HD3	2.35	0.42
12:B:302:FAD:H9	12:B:302:FAD:H1'1	1.85	0.42
2:A:308:PRO:HB3	2:A:408:PRO:HG2	2.02	0.41
1:G:236:LYS:HB3	1:G:236:LYS:HE3	1.88	0.41
2:A:380:ARG:CZ	8:A:501:J52:O	2.70	0.40
2:A:34:TRP:CH2	2:A:431:PRO:HB3	2.56	0.40
1:G:60:THR:O	2:A:12:ARG:HA	2.21	0.40
1:G:235:ILE:H	1:G:235:ILE:HG13	1.73	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	G	250/253 (99%)	238 (95%)	11 (4%)	1 (0%)	34 22
2	A	437/438 (100%)	430 (98%)	7 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	290/291 (100%)	280 (97%)	10 (3%)	0	100	100
All	All	977/982 (100%)	948 (97%)	28 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	58	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	214/215 (100%)	212 (99%)	2 (1%)	78	77
2	A	370/368 (100%)	362 (98%)	8 (2%)	52	46
3	B	247/246 (100%)	241 (98%)	6 (2%)	49	41
All	All	831/829 (100%)	815 (98%)	16 (2%)	57	50

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	69	GLU
1	G	206	MET
2	A	13	LEU
2	A	35	LEU
2	A	104	HIS
2	A	225	LYS
2	A	264	ASP
2	A	309	VAL
2	A	351	ASP
2	A	402	PRO
3	B	42	GLU
3	B	79	ARG
3	B	140	GLU
3	B	199	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	272	LYS
3	B	275	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	BU3	G	306	-	4,5,5	0.79	0	6,6,6	0.89	0
6	144	A	504	-	1,7,7	4.91	1 (100%)	3,9,9	0.03	0
12	FAD	B	302	-	53,58,58	3.57	23 (43%)	68,89,89	2.58	17 (25%)
4	SF4	G	302	1	0,12,12	-	-	-	-	-
4	SF4	B	303	3	0,12,12	-	-	-	-	-
4	SF4	G	303	1	0,12,12	-	-	-	-	-
7	BU3	G	307	-	4,5,5	0.76	0	6,6,6	0.50	0
7	BU3	B	301	-	4,5,5	0.98	0	6,6,6	0.53	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	J52	A	501	2	1,9,9	0.14	0	-		
4	SF4	G	301	1	0,12,12	-	-	-		
5	FES	G	304	1	0,2,4	-	-	-		
11	MPD	A	505	-	7,7,7	0.29	0	9,10,10	0.76	0
6	144	G	305	-	1,7,7	1.25	0	3,9,9	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BU3	G	306	-	-	0/4/4/4	-
6	144	A	504	-	-	0/0/9/9	-
4	SF4	G	302	1	-	-	0/6/5/5
4	SF4	B	303	3	-	-	0/6/5/5
4	SF4	G	303	1	-	-	0/6/5/5
7	BU3	G	307	-	-	4/4/4/4	-
7	BU3	B	301	-	-	4/4/4/4	-
12	FAD	B	302	-	-	4/30/50/50	0/6/6/6
4	SF4	G	301	1	-	-	0/6/5/5
11	MPD	A	505	-	-	0/5/5/5	-
6	144	G	305	-	-	0/0/9/9	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	B	302	FAD	O4B-C1B	13.30	1.59	1.41
12	B	302	FAD	C2B-C1B	-9.89	1.38	1.53
12	B	302	FAD	C4X-N5	7.79	1.45	1.30
12	B	302	FAD	C8A-N7A	6.77	1.46	1.34
12	B	302	FAD	C4A-N3A	5.93	1.43	1.35
12	B	302	FAD	C10-N1	5.26	1.44	1.33
6	A	504	144	C1-N	-4.91	1.30	1.50
12	B	302	FAD	C2A-N3A	4.45	1.39	1.32
12	B	302	FAD	O4B-C4B	-4.40	1.35	1.45
12	B	302	FAD	O4-C4	-4.40	1.15	1.23
12	B	302	FAD	C7M-C7	4.16	1.59	1.51
12	B	302	FAD	C9-C9A	4.04	1.46	1.39
12	B	302	FAD	O2B-C2B	3.69	1.51	1.43
12	B	302	FAD	C6A-N6A	3.63	1.47	1.34
12	B	302	FAD	C9-C8	-3.37	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	B	302	FAD	C2-N1	3.25	1.44	1.36
12	B	302	FAD	C5X-N5	3.09	1.45	1.39
12	B	302	FAD	C8M-C8	2.75	1.56	1.51
12	B	302	FAD	O3B-C3B	-2.71	1.36	1.43
12	B	302	FAD	C5'-C4'	2.68	1.55	1.51
12	B	302	FAD	C4-N3	2.61	1.43	1.38
12	B	302	FAD	O2'-C2'	-2.53	1.38	1.43
12	B	302	FAD	C2A-N1A	2.28	1.38	1.33
12	B	302	FAD	C10-N10	2.28	1.42	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	B	302	FAD	C7M-C7-C6	-9.61	101.72	119.49
12	B	302	FAD	C5A-C6A-N6A	8.20	132.82	120.35
12	B	302	FAD	C7M-C7-C8	7.91	136.95	120.74
12	B	302	FAD	N6A-C6A-N1A	-6.60	104.88	118.57
12	B	302	FAD	N3A-C2A-N1A	-6.54	118.46	128.68
12	B	302	FAD	C9-C9A-N10	-3.93	116.53	121.84
12	B	302	FAD	C4X-C10-N10	3.66	121.83	116.48
12	B	302	FAD	O4-C4-C4X	-2.93	118.82	126.60
12	B	302	FAD	C5X-N5-C4X	-2.87	113.30	118.07
12	B	302	FAD	C1B-N9A-C4A	-2.82	121.69	126.64
12	B	302	FAD	C4X-C4-N3	2.65	119.92	113.19
12	B	302	FAD	C5X-C9A-N10	2.54	120.58	117.95
12	B	302	FAD	C8M-C8-C7	-2.24	116.14	120.74
12	B	302	FAD	C9A-N10-C10	-2.16	117.40	120.77
12	B	302	FAD	C10-N1-C2	2.10	121.09	116.90
12	B	302	FAD	C4-N3-C2	-2.07	121.81	125.64
12	B	302	FAD	O3B-C3B-C2B	-2.02	105.30	111.82

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	G	307	BU3	O5-C2-C3-O6
7	G	307	BU3	C1-C2-C3-O6
7	G	307	BU3	O5-C2-C3-C4
7	G	307	BU3	C1-C2-C3-C4
7	B	301	BU3	O5-C2-C3-O6
7	B	301	BU3	C1-C2-C3-O6
12	B	302	FAD	O4B-C4B-C5B-O5B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	B	301	BU3	O5-C2-C3-C4
12	B	302	FAD	PA-O3P-P-O2P
7	B	301	BU3	C1-C2-C3-C4
12	B	302	FAD	C3B-C4B-C5B-O5B
12	B	302	FAD	PA-O3P-P-O1P

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	G	306	BU3	3	0
12	B	302	FAD	1	0
8	A	501	J52	2	0
4	G	301	SF4	1	0
6	G	305	144	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	252/253 (99%)	0.29	13 (5%) 27 29	31, 41, 60, 94	0
2	A	437/438 (99%)	-0.14	3 (0%) 87 88	29, 36, 49, 61	0
3	B	291/291 (100%)	-0.09	4 (1%) 75 77	34, 44, 62, 74	0
All	All	980/982 (99%)	-0.01	20 (2%) 65 66	29, 39, 58, 94	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	270	ASP	3.2
1	G	235	ILE	3.0
1	G	234	CYS	2.9
2	A	58	ILE	2.9
1	G	167	GLU	2.8
1	G	193	CYS	2.7
2	A	63	CYS	2.7
1	G	192	PHE	2.5
1	G	191	CYS	2.4
1	G	237	CYS	2.4
1	G	195	LEU	2.4
2	A	43	ILE	2.4
1	G	166	ASN	2.4
3	B	5	PRO	2.2
1	G	115	PHE	2.2
3	B	257	ASP	2.1
1	G	217	VAL	2.1
1	G	236	LYS	2.1
3	B	252	VAL	2.0
3	B	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	BU3	G	307	6/6	0.86	0.15	52,56,64,70	0
7	BU3	B	301	6/6	0.86	0.25	43,50,53,54	6
7	BU3	G	306	6/6	0.88	0.19	42,51,54,55	6
6	144	A	504	8/8	0.90	0.17	46,46,47,50	8
11	MPD	A	505	8/8	0.92	0.21	44,48,52,54	8
6	144	G	305	8/8	0.94	0.10	39,46,50,56	0
12	FAD	B	302	53/53	0.97	0.08	31,38,43,45	0
10	MG	A	503	1/1	0.98	0.12	37,37,37,37	1
4	SF4	B	303	8/8	0.99	0.06	38,40,41,42	0
5	FES	G	304	3/4	0.99	0.12	35,35,38,43	2
8	J52	A	501	10/10	0.99	0.14	31,33,37,37	2
4	SF4	G	301	8/8	0.99	0.10	35,35,36,37	0
4	SF4	G	302	8/8	0.99	0.12	32,35,36,38	0
4	SF4	G	303	8/8	0.99	0.09	36,37,39,39	0
9	FE	A	502	1/1	1.00	0.09	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.