



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 27, 2022 – 01:15 pm BST

PDB ID : 7QHL
Title : Crystal structure of Cyclin-dependent kinase 2/cyclin A in complex with 3,5,7-Substituted pyrazolo[4,3-d]pyrimidine inhibitor 24
Authors : Djukic, S.; Skerlova, J.; Rezacova, P.
Deposited on : 2021-12-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.29
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	 91% 5%
1	C	299	 90% 6%
2	B	258	 91% 9%
2	D	258	 92% 7%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	285	Total	C	N	O	P	S	0	6	0
			2328	1517	395	407	1	8			
1	C	287	Total	C	N	O	P	S	0	5	0
			2336	1521	396	410	1	8			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P24941
C	0	SER	-	expression tag	UNP P24941

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	258	Total	C	N	O	S	0	4	0
			2107	1367	342	385	13			
2	D	256	Total	C	N	O	S	0	3	0
			2085	1353	338	382	12			

- Molecule 3 is 5-(2-amino-1-ethyl)thio-3-cyclobutyl-7-[4-(pyrazol-1-yl)benzyl]amino-1(2H)-pyrazolo[4,3-d]pyrimidine (three-letter code: D5P) (formula: C₂₁H₂₄N₈S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	N	S	0	0
			30	21	8	1		
3	C	1	Total	C	N	S	0	0
			30	21	8	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			6	3 3		
4	C	1	Total	C O	0	0
			6	3 3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		
6	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is MONOTHIOGLYCEROL (three-letter code: SGM) (formula: C₃H₈O₂S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	O	S	0	0
			6	3	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	O	S	0	0
			6	3	2	1		
7	D	1	Total	C	O	S	0	0
			6	3	2	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	252	Total	O	0	5
			255	255		
8	B	269	Total	O	0	0
			269	269		
8	C	251	Total	O	0	3
			251	251		
8	D	192	Total	O	0	2
			194	194		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclin-dependent kinase 2





- Molecule 1: Cyclin-dependent kinase 2



- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.89Å 163.90Å 73.17Å 90.00° 106.61° 90.00°	Depositor
Resolution (Å)	46.96 – 1.70 46.96 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (46.96-1.70) 97.4 (46.96-1.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.200 , 0.224 0.209 , 0.231	Depositor DCC
R_{free} test set	2100 reflections (1.25%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtrriage
Anisotropy	0.237	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.009 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9960	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, GOL, SGM, EDO, PEG, D5P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/2392	0.83	0/3239
1	C	0.75	0/2399	0.84	0/3251
2	B	0.73	0/2170	0.80	0/2945
2	D	0.70	0/2144	0.79	0/2907
All	All	0.73	0/9105	0.82	0/12342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	2395	13	0
1	C	2336	0	2395	11	0
2	B	2107	0	2139	17	0
2	D	2085	0	2112	14	0
3	A	30	0	0	0	0
3	C	30	0	0	0	0
4	A	6	0	8	0	0
4	C	6	0	8	0	0
5	A	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	8	0	12	1	0
5	C	4	0	6	0	0
5	D	4	0	6	0	0
6	B	14	0	20	2	0
6	D	7	0	10	0	0
7	B	6	0	7	0	0
7	D	12	0	15	0	0
8	A	255	0	0	2	0
8	B	269	0	0	3	0
8	C	251	0	0	2	0
8	D	194	0	0	3	0
All	All	9960	0	9145	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268[B]:HIS:CE1	8:A:401[B]:HOH:O	1.94	1.14
1:A:268[B]:HIS:ND1	8:A:401[B]:HOH:O	1.77	1.12
2:D:229:ASN:HD22	2:D:334[B]:MET:CE	1.99	0.75
2:B:296[B]:HIS:HD2	8:B:801:HOH:O	1.71	0.74
1:A:35:ILE:HD12	1:A:78:LEU:HD13	1.74	0.69
2:D:229:ASN:HD22	2:D:334[B]:MET:HE3	1.59	0.68
2:D:296[B]:HIS:HE1	8:D:752:HOH:O	1.75	0.68
1:C:71[A]:HIS:ND1	8:C:401[A]:HOH:O	2.26	0.67
1:C:71[B]:HIS:HB2	1:C:76:LEU:HD13	1.78	0.64
2:B:229:ASN:HD22	2:B:334[B]:MET:CE	2.14	0.60
2:D:296[B]:HIS:CE1	8:D:752:HOH:O	2.51	0.59
2:B:184:THR:HG23	6:B:502:PEG:H11	1.84	0.59
1:A:22[B]:ARG:HH11	2:D:249:LEU:HD11	1.69	0.57
2:B:327:CYS:HB3	2:B:419:HIS:CE1	2.41	0.56
1:A:49:ILE:HG23	2:B:306:LEU:HD12	1.87	0.56
1:A:268[A]:HIS:HD2	1:A:270:ASP:H	1.53	0.55
2:B:252:LYS:NZ	5:B:504:EDO:H11	2.20	0.55
2:B:229:ASN:HD22	2:B:334[B]:MET:HE2	1.71	0.54
2:D:327:CYS:HB3	2:D:419:HIS:CE1	2.42	0.54
2:B:296[B]:HIS:HE1	8:B:805:HOH:O	1.89	0.54
2:B:296[B]:HIS:CE1	8:B:805:HOH:O	2.61	0.53
1:C:49:ILE:HG23	2:D:306:LEU:HD12	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:177:CYS:HB2	1:C:233:MET:CE	2.40	0.51
2:B:274:GLU:HG2	2:B:277:GLU:HG3	1.96	0.48
1:A:71:HIS:CD2	1:A:71:HIS:O	2.67	0.48
2:D:229:ASN:ND2	2:D:334[B]:MET:HE3	2.27	0.48
2:B:388:LYS:HB3	2:B:389:PRO:HD3	1.97	0.47
1:C:2:GLU:O	1:C:24:LYS:CE	2.63	0.47
1:A:71:HIS:O	1:A:71:HIS:CG	2.68	0.46
1:C:89:LYS:HD3	1:C:89:LYS:HA	1.65	0.46
2:B:254:GLN:HB2	6:B:501:PEG:H31	1.97	0.46
1:C:34:LYS:HE3	1:C:75:LYS:HD2	1.98	0.46
2:D:388:LYS:HB3	2:D:389:PRO:HD3	1.98	0.46
1:A:35:ILE:HD12	1:A:78:LEU:CD1	2.43	0.45
1:A:2:GLU:O	1:A:24:LYS:CE	2.65	0.44
2:D:223:GLU:OE1	8:D:601[A]:HOH:O	2.20	0.44
2:D:401:ALA:N	2:D:402:PRO:CD	2.80	0.44
2:B:275:VAL:HG21	2:B:292:LEU:HD21	2.00	0.43
2:B:364:LEU:HG	2:B:370:GLN:HB2	2.00	0.43
1:A:60:HIS:CG	1:A:61:PRO:HD2	2.52	0.43
1:C:177:CYS:HB2	1:C:233:MET:HE1	1.99	0.43
1:C:78:LEU:HD23	1:C:78:LEU:N	2.34	0.43
2:B:401:ALA:N	2:B:402:PRO:CD	2.82	0.42
2:B:345:ASP:HA	2:B:346:PRO:HA	1.82	0.42
1:C:199:ARG:HA	8:C:444:HOH:O	2.20	0.42
2:D:274:GLU:HG2	2:D:277[A]:GLU:HG3	2.02	0.42
2:D:345:ASP:HA	2:D:346:PRO:HA	1.81	0.42
1:A:78:LEU:HD12	1:A:78:LEU:N	2.36	0.41
1:A:268[A]:HIS:CD2	1:A:270:ASP:H	2.37	0.41
2:B:193:CYS:O	2:B:241:ARG:HD2	2.21	0.40
2:D:384:LEU:HD23	2:D:384:LEU:HA	1.87	0.40
1:C:60:HIS:CG	1:C:61:PRO:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/299 (94%)	277 (98%)	4 (1%)	1 (0%)	34	18
1	C	285/299 (95%)	279 (98%)	5 (2%)	1 (0%)	34	18
2	B	260/258 (101%)	258 (99%)	2 (1%)	0	100	100
2	D	255/258 (99%)	253 (99%)	2 (1%)	0	100	100
All	All	1082/1114 (97%)	1067 (99%)	13 (1%)	2 (0%)	47	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	VAL
1	C	164	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/263 (98%)	256 (100%)	1 (0%)	91	87
1	C	258/263 (98%)	255 (99%)	3 (1%)	71	59
2	B	236/232 (102%)	234 (99%)	2 (1%)	81	74
2	D	233/232 (100%)	232 (100%)	1 (0%)	91	87
All	All	984/990 (99%)	977 (99%)	7 (1%)	86	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	ARG
2	B	323[A]	GLN
2	B	323[B]	GLN
1	C	74	ASN
1	C	122	ARG
1	C	150	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	428	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	HIS
1	A	265	GLN
2	D	396	GLN
2	D	425	ASN
2	D	431	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	160	1	8,10,11	0.79	0	10,14,16	1.02	0
1	TPO	C	160	1	8,10,11	0.56	0	10,14,16	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/9/11/13	-
1	TPO	C	160	1	-	0/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PEG	D	503	-	6,6,6	0.30	0	5,5,5	0.18	0
4	GOL	A	302	-	5,5,5	0.16	0	5,5,5	0.70	0
3	D5P	C	301	-	32,34,34	1.06	2 (6%)	31,47,47	0.57	0
5	EDO	A	304	-	3,3,3	0.07	0	2,2,2	0.47	0
7	SGM	D	501	2	5,5,5	0.34	0	5,5,5	0.83	0
5	EDO	B	504	-	3,3,3	0.19	0	2,2,2	0.22	0
5	EDO	D	504	-	3,3,3	0.15	0	2,2,2	0.19	0
4	GOL	C	302	-	5,5,5	0.10	0	5,5,5	0.92	0
7	SGM	B	503	2	5,5,5	0.34	0	5,5,5	1.40	1 (20%)
6	PEG	B	501	-	6,6,6	0.61	0	5,5,5	0.49	0
7	SGM	D	502	-	5,5,5	0.16	0	5,5,5	0.34	0
5	EDO	A	303	-	3,3,3	0.32	0	2,2,2	0.38	0
5	EDO	B	505	-	3,3,3	0.63	0	2,2,2	0.54	0
6	PEG	B	502	-	6,6,6	0.24	0	5,5,5	0.21	0
5	EDO	C	303	-	3,3,3	0.22	0	2,2,2	0.35	0
3	D5P	A	301	-	32,34,34	0.84	1 (3%)	31,47,47	0.68	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	D	503	-	-	3/4/4/4	-
4	GOL	A	302	-	-	1/4/4/4	-
3	D5P	C	301	-	-	2/13/23/23	0/5/5/5
5	EDO	A	304	-	-	0/1/1/1	-
7	SGM	D	501	2	-	1/4/4/4	-
5	EDO	B	504	-	-	1/1/1/1	-
5	EDO	D	504	-	-	0/1/1/1	-
4	GOL	C	302	-	-	2/4/4/4	-
7	SGM	B	503	2	-	2/4/4/4	-
6	PEG	B	501	-	-	2/4/4/4	-
7	SGM	D	502	-	-	2/4/4/4	-
5	EDO	A	303	-	-	1/1/1/1	-
5	EDO	B	505	-	-	1/1/1/1	-
6	PEG	B	502	-	-	2/4/4/4	-
5	EDO	C	303	-	-	1/1/1/1	-
3	D5P	A	301	-	-	2/13/23/23	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	301	D5P	C16-N7	3.81	1.39	1.34
3	A	301	D5P	C16-C21	-2.77	1.38	1.43
3	C	301	D5P	C16-C17	-2.11	1.48	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	503	SGM	C2-C1-S1	2.79	122.59	114.47
3	A	301	D5P	C15-C4-N2	-2.21	118.97	120.81

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	302	GOL	C1-C2-C3-O3
7	B	503	SGM	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	D	502	SGM	S1-C1-C2-O2
7	D	502	SGM	S1-C1-C2-C3
6	B	502	PEG	C1-C2-O2-C3
7	B	503	SGM	O2-C2-C3-O3
4	A	302	GOL	C1-C2-C3-O3
4	C	302	GOL	O2-C2-C3-O3
5	A	303	EDO	O1-C1-C2-O2
5	B	504	EDO	O1-C1-C2-O2
5	B	505	EDO	O1-C1-C2-O2
6	D	503	PEG	O1-C1-C2-O2
6	B	501	PEG	C4-C3-O2-C2
6	B	502	PEG	O2-C3-C4-O4
3	C	301	D5P	C13-C9-N4-C10
5	C	303	EDO	O1-C1-C2-O2
6	D	503	PEG	C4-C3-O2-C2
6	D	503	PEG	C1-C2-O2-C3
3	A	301	D5P	C8-C9-N4-C10
3	A	301	D5P	C13-C9-N4-C10
3	C	301	D5P	C8-C9-N4-C10
6	B	501	PEG	C1-C2-O2-C3
7	D	501	SGM	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	504	EDO	1	0
6	B	501	PEG	1	0
6	B	502	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/299 (94%)	-0.36	2 (0%) 87 90	18, 28, 55, 85	0
1	C	286/299 (95%)	-0.35	6 (2%) 63 67	19, 28, 57, 95	0
2	B	258/258 (100%)	-0.30	3 (1%) 79 82	18, 27, 46, 71	0
2	D	256/258 (99%)	-0.09	5 (1%) 65 69	19, 32, 58, 83	0
All	All	1084/1114 (97%)	-0.28	16 (1%) 73 77	18, 29, 55, 95	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	432	LEU	5.3
1	A	36	ARG	4.5
1	C	36	ARG	4.4
1	C	73	GLU	3.2
2	D	384	LEU	2.7
1	A	38	ASP	2.6
1	C	250	LYS	2.5
2	D	431	ASN	2.5
1	C	17	VAL	2.5
2	B	175	VAL	2.5
1	C	74	ASN	2.4
2	D	301	VAL	2.4
1	C	71[A]	HIS	2.2
2	B	431	ASN	2.1
2	D	432	LEU	2.1
2	D	177	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	160	11/12	0.98	0.06	19,20,24,25	0
1	TPO	C	160	11/12	0.98	0.06	19,23,24,25	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SGM	D	502	6/6	0.72	0.21	75,80,90,103	0
6	PEG	B	501	7/7	0.75	0.14	45,49,53,56	0
5	EDO	A	303	4/4	0.75	0.12	48,52,53,55	0
5	EDO	B	505	4/4	0.77	0.11	40,47,47,47	0
6	PEG	D	503	7/7	0.83	0.14	45,53,60,61	0
6	PEG	B	502	7/7	0.90	0.14	34,51,57,58	0
5	EDO	C	303	4/4	0.91	0.23	52,54,56,60	0
5	EDO	A	304	4/4	0.93	0.13	52,52,53,55	0
5	EDO	B	504	4/4	0.93	0.07	43,46,48,49	0
4	GOL	A	302	6/6	0.95	0.10	34,42,47,51	0
3	D5P	A	301	30/30	0.96	0.07	22,26,37,40	0
5	EDO	D	504	4/4	0.96	0.12	52,54,57,57	0
4	GOL	C	302	6/6	0.96	0.09	29,34,37,38	0
7	SGM	B	503	6/6	0.97	0.09	49,62,64,67	0
7	SGM	D	501	6/6	0.97	0.08	45,52,57,62	0
3	D5P	C	301	30/30	0.97	0.07	21,27,42,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D5P A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around D5P C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.