



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2023 – 09:05 PM EDT

PDB ID : 1QI3  
Title : MUTANT (D193N) MALTOTETRAOSE-FORMING EXO-AMYLASE IN  
COMPLEX WITH MALTOTETRAOSE  
Authors : Hasegawa, K.; Kubota, M.; Matsuura, Y.  
Deposited on : 1999-06-01  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	 71% 21% 5%
2	B	4	 75% 25%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (EXO-MALTOTETRAOHYDROLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	418	3297	2070	597	620	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	193	ASN	ASP	engineered mutation	UNP P13507

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	4	45	24	21	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	157	Total	O	0	0
			157	157		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (EXO-MALTOTETRAHYDROLASE)



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.77Å 170.80Å 46.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 61.38 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 91.3 (61.38-1.99)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 2.00Å)	Xtrriage
Refinement program	PROFFT	Depositor
R, $R_{free}$	0.186 , (Not available) 0.165 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtrriage
Anisotropy	0.247	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 48.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3501	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	1/3402 (0.0%)	1.58	51/4631 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	219	GLU	CD-OE2	7.34	1.33	1.25

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	ARG	NE-CZ-NH1	18.29	129.45	120.30
1	A	175	ARG	CD-NE-CZ	16.90	147.25	123.60
1	A	175	ARG	NE-CZ-NH2	-13.25	113.67	120.30
1	A	191	ARG	NE-CZ-NH1	11.85	126.22	120.30
1	A	233	ARG	NE-CZ-NH1	10.82	125.71	120.30
1	A	137	ARG	NE-CZ-NH1	10.34	125.47	120.30
1	A	11	ARG	NE-CZ-NH1	10.15	125.38	120.30
1	A	82	ASP	CB-CG-OD2	10.08	127.37	118.30
1	A	137	ARG	NE-CZ-NH2	-9.81	115.39	120.30
1	A	417	ARG	CD-NE-CZ	-9.55	110.22	123.60
1	A	353	ARG	NE-CZ-NH1	-9.24	115.68	120.30
1	A	248	ARG	NE-CZ-NH1	9.11	124.86	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	358	ARG	NE-CZ-NH2	-9.05	115.78	120.30
1	A	233	ARG	NE-CZ-NH2	-8.40	116.10	120.30
1	A	279	ASP	CB-CG-OD1	7.92	125.43	118.30
1	A	347	GLN	CA-CB-CG	7.82	130.61	113.40
1	A	29	ARG	NE-CZ-NH2	7.79	124.20	120.30
1	A	390	ASP	CB-CG-OD1	7.47	125.02	118.30
1	A	96	ARG	NE-CZ-NH2	-7.39	116.60	120.30
1	A	120	ARG	NE-CZ-NH2	-7.21	116.70	120.30
1	A	417	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	A	414	ARG	NE-CZ-NH1	6.87	123.74	120.30
1	A	334	ASP	CA-CB-CG	-6.85	98.34	113.40
1	A	321	TYR	CB-CG-CD1	-6.82	116.91	121.00
1	A	212	ASP	CB-CG-OD1	6.75	124.38	118.30
1	A	346	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	A	319	TYR	CB-CG-CD1	-6.40	117.16	121.00
1	A	96	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	248	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	A	162	ASP	CB-CG-OD1	-6.00	112.89	118.30
1	A	414	ARG	CD-NE-CZ	5.89	131.84	123.60
1	A	373	VAL	CA-CB-CG1	5.84	119.66	110.90
1	A	82	ASP	CB-CG-OD1	-5.82	113.06	118.30
1	A	279	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	A	316	ARG	NE-CZ-NH1	5.67	123.14	120.30
1	A	397	VAL	N-CA-CB	-5.63	99.12	111.50
1	A	40	ARG	NE-CZ-NH1	-5.59	117.50	120.30
1	A	140	CYS	CA-CB-SG	5.53	123.96	114.00
1	A	369	TYR	CA-CB-CG	-5.53	102.89	113.40
1	A	226	GLU	OE1-CD-OE2	5.49	129.89	123.30
1	A	196	ARG	NE-CZ-NH2	5.44	123.02	120.30
1	A	92	ASP	CB-CG-OD2	5.37	123.13	118.30
1	A	373	VAL	CB-CA-C	5.32	121.50	111.40
1	A	244	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	A	231	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	372	LEU	CA-CB-CG	5.18	127.22	115.30
1	A	68	ASP	N-CA-CB	5.14	119.85	110.60
1	A	366	HIS	C-N-CA	5.13	134.52	121.70
1	A	62	ASP	CB-CG-OD1	5.10	122.89	118.30
1	A	283	ARG	NE-CZ-NH2	5.07	122.83	120.30
1	A	358	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	417	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3297	0	3015	77	0
2	B	45	0	39	4	0
3	A	2	0	0	0	0
4	A	157	0	0	6	0
All	All	3501	0	3054	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:PRO:HD3	1:A:334:ASP:OD2	1.64	0.95
1:A:237:SER:H	1:A:240:GLN:HE21	1.18	0.90
1:A:167:HIS:HD2	1:A:169:GLN:H	1.22	0.88
1:A:317:GLN:H	1:A:317:GLN:HE21	1.20	0.86
1:A:277:ASN:HD22	1:A:279:ASP:H	1.25	0.83
1:A:405:ALA:HB2	1:A:417:ARG:HD3	1.61	0.82
1:A:11:ARG:HH12	1:A:204:ASN:HD22	1.28	0.82
1:A:193:ASN:ND2	2:B:1:GLC:H1	1.98	0.78
1:A:193:ASN:HD21	2:B:1:GLC:H1	1.51	0.76
1:A:127:ILE:HG21	1:A:173:MET:HE1	1.66	0.76
1:A:216:CYS:O	1:A:252:PRO:HD2	1.88	0.73
1:A:195:VAL:HG13	1:A:245:TRP:NE1	2.06	0.70
1:A:145:ASN:ND2	1:A:155:ARG:H	1.90	0.69
1:A:145:ASN:HD21	1:A:154:ASP:HB3	1.57	0.69
1:A:237:SER:H	1:A:240:GLN:NE2	1.89	0.68
1:A:299:PRO:CD	1:A:334:ASP:OD2	2.41	0.67
1:A:146:TYR:CD1	1:A:147:PRO:HD2	2.33	0.63
1:A:81:HIS:HD2	4:A:508:HOH:O	1.82	0.62
1:A:277:ASN:ND2	1:A:279:ASP:H	1.96	0.61

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:ASN:ND2	1:A:295:THR:H	2.01	0.58
1:A:2:GLN:HE22	1:A:358:ARG:NH1	2.01	0.58
1:A:26:ASN:HD22	1:A:29:ARG:HE	1.50	0.58
1:A:145:ASN:HD21	1:A:155:ARG:H	1.52	0.58
1:A:405:ALA:CB	1:A:417:ARG:HD3	2.33	0.57
1:A:127:ILE:CG2	1:A:173:MET:HE1	2.34	0.57
1:A:2:GLN:NE2	1:A:358:ARG:HH12	2.02	0.56
1:A:86:ASN:ND2	1:A:92:ASP:H	2.04	0.55
1:A:127:ILE:HD13	1:A:173:MET:HE1	1.87	0.55
1:A:277:ASN:O	1:A:283:ARG:HD3	2.08	0.54
1:A:325:SER:HB2	1:A:326:PRO:CD	2.38	0.54
1:A:26:ASN:ND2	1:A:29:ARG:HE	2.06	0.53
1:A:167:HIS:CD2	1:A:169:GLN:H	2.13	0.53
1:A:404:GLU:HG3	4:A:765:HOH:O	2.09	0.53
1:A:11:ARG:HH12	1:A:204:ASN:ND2	2.02	0.53
1:A:11:ARG:HH22	1:A:204:ASN:ND2	2.06	0.52
1:A:325:SER:HB2	1:A:326:PRO:HD2	1.90	0.52
1:A:193:ASN:HD21	2:B:1:GLC:C1	2.22	0.51
1:A:332:TYR:HB3	1:A:335:HIS:CD2	2.45	0.51
1:A:31:ALA:N	1:A:32:PRO:CD	2.73	0.51
1:A:61:ARG:NH2	1:A:82:ASP:OD2	2.39	0.51
1:A:292:ASN:ND2	1:A:294:ASP:H	2.10	0.50
1:A:407:ASN:C	1:A:407:ASN:HD22	2.15	0.50
1:A:303:GLY:HA2	4:A:768:HOH:O	2.11	0.50
1:A:292:ASN:HD22	1:A:294:ASP:H	1.59	0.49
1:A:24:HIS:HD2	1:A:26:ASN:H	1.59	0.49
1:A:146:TYR:CG	1:A:147:PRO:HD2	2.49	0.48
1:A:317:GLN:HE21	1:A:317:GLN:N	2.01	0.48
1:A:292:ASN:HD22	1:A:292:ASN:C	2.17	0.48
1:A:224:PRO:O	1:A:234:ASN:HA	2.14	0.47
1:A:332:TYR:HB3	1:A:335:HIS:HD2	1.78	0.47
1:A:61:ARG:NE	1:A:84:ASN:HB3	2.29	0.46
4:A:725:HOH:O	2:B:1:GLC:H2	2.14	0.46
1:A:334:ASP:N	1:A:334:ASP:OD1	2.48	0.46
1:A:127:ILE:CG2	1:A:173:MET:CE	2.94	0.46
1:A:317:GLN:H	1:A:317:GLN:NE2	2.00	0.46
1:A:145:ASN:HD21	1:A:154:ASP:CB	2.29	0.45
1:A:353:ARG:HE	1:A:353:ARG:HB2	1.47	0.45
1:A:13:HIS:CE1	1:A:108:LYS:HE2	2.52	0.45
1:A:40:ARG:HH11	1:A:40:ARG:HD2	1.60	0.44
1:A:319:TYR:OH	1:A:335:HIS:CD2	2.71	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:ASP:OD2	1:A:108:LYS:NZ	2.31	0.44
1:A:334:ASP:OD1	1:A:338:ASP:OD2	2.35	0.44
1:A:182:ARG:HA	1:A:187:ALA:HB3	2.00	0.43
1:A:2:GLN:NE2	1:A:358:ARG:NH1	2.63	0.43
1:A:24:HIS:CD2	1:A:299:PRO:HD2	2.53	0.43
1:A:2:GLN:HE22	1:A:358:ARG:HH12	1.63	0.43
1:A:199:ALA:HA	1:A:200:PRO:HD3	1.90	0.43
1:A:417:ARG:HD2	4:A:526:HOH:O	2.17	0.43
1:A:392:GLY:N	1:A:396:GLN:OE1	2.50	0.43
1:A:62:ASP:O	1:A:75:GLY:HA2	2.19	0.43
1:A:319:TYR:O	1:A:323:LEU:HB2	2.19	0.43
1:A:381:GLN:OE1	1:A:417:ARG:HG2	2.19	0.42
1:A:271:LYS:HE2	4:A:705:HOH:O	2.19	0.42
1:A:50:GLY:HA2	1:A:353:ARG:HH12	1.85	0.41
1:A:332:TYR:O	1:A:333:TRP:C	2.58	0.41
1:A:127:ILE:HD13	1:A:173:MET:CE	2.50	0.41
1:A:68:ASP:OD1	1:A:68:ASP:O	2.38	0.40
1:A:218:GLY:HA3	1:A:245:TRP:CH2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/429 (97%)	402 (97%)	12 (3%)	2 (0%)	29 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	ILE
1	A	143	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	333/337 (99%)	311 (93%)	22 (7%)	16 12

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	12	TYR
1	A	65	SER
1	A	95	LEU
1	A	102	LEU
1	A	119	ASN
1	A	162	ASP
1	A	175	ARG
1	A	181	LEU
1	A	182	ARG
1	A	183	SER
1	A	292	ASN
1	A	301	GLN
1	A	308	TRP
1	A	317	GLN
1	A	323	LEU
1	A	367	SER
1	A	373	VAL
1	A	383	LEU
1	A	397	VAL
1	A	401	SER
1	A	407	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	24	HIS
1	A	26	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	81	HIS
1	A	86	ASN
1	A	97	GLN
1	A	119	ASN
1	A	128	ASN
1	A	145	ASN
1	A	167	HIS
1	A	204	ASN
1	A	240	GLN
1	A	264	ASN
1	A	277	ASN
1	A	292	ASN
1	A	311	GLN
1	A	317	GLN
1	A	335	HIS
1	A	347	GLN
1	A	366	HIS
1	A	388	ASN
1	A	407	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	B	1	2	12,12,12	1.24	1 (8%)	17,17,17	2.53	5 (29%)
2	GLC	B	2	2	11,11,12	0.81	0	15,15,17	1.35	3 (20%)
2	GLC	B	3	2	11,11,12	0.53	0	15,15,17	1.01	1 (6%)
2	GLC	B	4	2	11,11,12	0.58	0	15,15,17	1.42	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	2/2/19/22	0/1/1/1
2	GLC	B	4	2	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLC	O3-C3	2.22	1.48	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-O5-C5	6.25	125.46	113.66
2	B	1	GLC	C1-C2-C3	4.32	119.27	110.31
2	B	1	GLC	O5-C1-C2	3.94	117.32	110.28
2	B	4	GLC	C1-O5-C5	3.38	116.77	112.19
2	B	1	GLC	C4-C3-C2	3.38	116.72	110.82
2	B	1	GLC	C3-C4-C5	3.27	116.07	110.24
2	B	2	GLC	C1-O5-C5	3.27	116.62	112.19
2	B	4	GLC	O3-C3-C2	2.59	114.96	109.99
2	B	4	GLC	C1-C2-C3	2.46	112.69	109.67
2	B	2	GLC	C1-C2-C3	2.31	112.50	109.67
2	B	3	GLC	O3-C3-C4	2.16	115.34	110.35
2	B	2	GLC	O5-C1-C2	2.14	114.07	110.77

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	4	GLC	O5-C5-C6-O6
2	B	3	GLC	O5-C5-C6-O6
2	B	4	GLC	C4-C5-C6-O6
2	B	3	GLC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	GLC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	418/429 (97%)	-0.43	2 (0%) 91 90	11, 18, 32, 52	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	TYR	3.8
1	A	2	GLN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	1	12/12	0.94	0.13	21,22,22,24	0
2	GLC	B	2	11/12	0.97	0.08	17,19,19,20	0
2	GLC	B	4	11/12	0.97	0.08	23,24,25,25	0
2	GLC	B	3	11/12	0.98	0.08	21,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	A	452	1/1	0.99	0.06	20,20,20,20	0
3	CA	A	451	1/1	1.00	0.07	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.