



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 10, 2022 – 01:06 pm GMT

PDB ID : 7QIV
Title : Structure of human C3b in complex with the EWE nanobody
Authors : Pedersen, H.; Andersen, G.R.
Deposited on : 2021-12-16
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.26
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	
2	B	915	
3	C	133	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement C3 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	638	4969	3164	843	947	15	0	0	0

- Molecule 2 is a protein called Complement C3b alpha' chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	903	7208	4569	1212	1389	38	4	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1013	GLU	GLN	conflict	UNP P01024

- Molecule 3 is a protein called Nanobody EWE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	124	950	591	170	184	5	1	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	14	0
4	B	1	14	8	1	5	14	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Complement C3 beta chain



- Molecule 2: Complement C3b alpha' chain



- Molecule 3: Nanobody EWE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	255.31Å 64.88Å 144.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.31 – 2.80 48.31 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.31-2.80) 98.6 (48.31-2.80)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.235 , 0.275 0.234 , 0.270	Depositor DCC
R_{free} test set	1471 reflections (2.47%)	wwPDB-VP
Wilson B-factor (Å ²)	83.2	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13155	wwPDB-VP
Average B, all atoms (Å ²)	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/5069	0.55	0/6889
2	B	0.29	0/7352	0.55	1/9954 (0.0%)
3	C	0.28	0/971	0.57	0/1313
All	All	0.29	0/13392	0.55	1/18156 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1202	LEU	CA-CB-CG	5.43	127.78	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4969	0	5031	16	0
2	B	7208	0	7135	35	1
3	C	950	0	901	6	0
4	A	14	0	13	0	0
4	B	14	0	13	0	0
All	All	13155	0	13093	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:835:LEU:HG	2:B:929:LEU:HD23	1.83	0.61
2:B:1046:LEU:HG	2:B:1093:ILE:HD11	1.83	0.61
1:A:141:PRO:O	2:B:930:LYS:NZ	2.37	0.58
2:B:833:LEU:HG	2:B:835:LEU:HD13	1.87	0.57
3:C:102:ALA:HB3	3:C:116:TYR:HB2	1.87	0.57
2:B:1050:LYS:HG3	2:B:1093:ILE:HD13	1.87	0.57
2:B:1399:THR:OG1	2:B:1400:MET:N	2.38	0.56
1:A:594:VAL:HG12	2:B:775:VAL:HG22	1.89	0.56
1:A:474:ASN:HB3	1:A:514:LEU:HD11	1.88	0.54
2:B:862:VAL:HG22	2:B:916:VAL:HG12	1.90	0.53
2:B:1128:GLU:HB3	2:B:1267:GLY:HA2	1.90	0.53
2:B:1151:LEU:HB3	2:B:1165:LEU:HD11	1.91	0.53
2:B:1203:LYS:HB2	2:B:1206:LEU:HG	1.91	0.51
1:A:372:LEU:HB2	1:A:409:LEU:HB2	1.94	0.50
2:B:1494:HIS:HD1	2:B:1497:LYS:HB2	1.76	0.49
2:B:1354:ASP:OD1	2:B:1354:ASP:N	2.41	0.49
1:A:494:ILE:HD12	1:A:502:LYS:HB3	1.95	0.48
2:B:764:ARG:HB3	2:B:797:ASP:HB3	1.95	0.48
1:A:360:THR:HG21	1:A:372:LEU:HD23	1.96	0.47
1:A:291:ILE:HD13	1:A:300:VAL:HB	1.96	0.47
2:B:1029:ASP:OD2	2:B:1042:ARG:NH2	2.44	0.47
3:C:36:ASN:HA	3:C:57:ARG:HB2	1.96	0.47
1:A:628:THR:HB	1:A:641:ASP:HB3	1.96	0.47
2:B:1075:SER:HB3	2:B:1078:LEU:HB3	1.97	0.47
2:B:1060:ARG:NH1	2:B:1099:VAL:HG22	2.31	0.46
2:B:1363:LEU:HD22	2:B:1479:VAL:HG23	1.97	0.46
1:A:46:ALA:HB3	1:A:82:HIS:HB3	1.97	0.46
2:B:1532:ARG:NH2	2:B:1629:GLU:OE2	2.40	0.46
1:A:176:LYS:HA	2:B:1320:ARG:HH12	1.81	0.46
1:A:456:LEU:HB2	1:A:535:TYR:HE2	1.81	0.46
3:C:32:ILE:HD11	3:C:38:MET:HG3	1.98	0.46
1:A:566:LYS:HE3	1:A:584:LYS:HD2	1.98	0.45
3:C:95:THR:HG23	3:C:124:THR:HA	1.98	0.45
2:B:799:ILE:HG23	2:B:826:MET:HA	1.97	0.45
2:B:1601:HIS:ND1	2:B:1633:GLU:OE2	2.49	0.45
1:A:465:LEU:HD21	1:A:521:ILE:HG13	2.00	0.44
3:C:54:THR:HG21	3:C:108:PRO:HB2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:873:CYS:HB3	2:B:901:VAL:HB	2.00	0.44
2:B:1124:VAL:HG11	2:B:1130:ILE:HD11	2.00	0.44
1:A:252:GLU:HG2	1:A:301:VAL:HG22	1.99	0.44
2:B:763:SER:OG	2:B:911:GLU:OE2	2.34	0.43
2:B:1415:THR:HG22	2:B:1419:LYS:HE2	2.01	0.43
2:B:1120:GLU:HG3	2:B:1122:ALA:H	1.82	0.43
2:B:1104:VAL:HG13	2:B:1151:LEU:HD22	2.01	0.42
2:B:1392:TYR:CG	2:B:1398:ALA:HB2	2.55	0.42
3:C:56:ASN:HB2	3:C:61:ARG:H	1.84	0.42
1:A:138:ILE:HG13	1:A:223:GLU:HB3	2.00	0.42
2:B:1142:ALA:HB2	2:B:1188:THR:HG23	2.01	0.41
1:A:180:LEU:HD11	1:A:191:LEU:HD21	2.01	0.41
2:B:1203:LYS:H	2:B:1206:LEU:HD12	1.85	0.41
2:B:1023:ILE:HG13	2:B:1276:PHE:HB2	2.03	0.41
2:B:1422:ALA:HB2	2:B:1429:ILE:HB	2.03	0.41
2:B:1404:ASP:HB3	2:B:1478:LYS:HB3	2.03	0.41
2:B:865:GLU:HB2	2:B:883:GLN:HG2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:959:LYS:NZ	2:B:1635:ASP:OD2[3_455]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	634/645 (98%)	616 (97%)	18 (3%)	0	100 100
2	B	899/915 (98%)	866 (96%)	32 (4%)	1 (0%)	51 81
3	C	122/133 (92%)	117 (96%)	5 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1655/1693 (98%)	1599 (97%)	55 (3%)	1 (0%)	51 81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1199	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	562/567 (99%)	555 (99%)	7 (1%)	71 92
2	B	798/810 (98%)	788 (99%)	10 (1%)	69 91
3	C	98/107 (92%)	97 (99%)	1 (1%)	76 93
All	All	1458/1484 (98%)	1440 (99%)	18 (1%)	71 92

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	LEU
1	A	199	VAL
1	A	365	LYS
1	A	463	THR
1	A	484	GLU
1	A	576	VAL
1	A	628	THR
2	B	759	GLU
2	B	835	LEU
2	B	967	LEU
2	B	1092	LEU
2	B	1188	THR
2	B	1201	ARG
2	B	1275	VAL
2	B	1291	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1456	ASP
2	B	1499	ASP
3	C	54	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	1701	2	14,14,15	0.32	0	17,19,21	0.44	0
4	NAG	A	2000	1	14,14,15	0.35	0	17,19,21	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1701	2	-	1/6/23/26	0/1/1/1
4	NAG	A	2000	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2000	NAG	C4-C5-C6-O6
4	A	2000	NAG	O5-C5-C6-O6
4	B	1701	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	638/645 (98%)	0.40	12 (1%) 66 59	50, 85, 135, 191	0
2	B	903/915 (98%)	0.87	111 (12%) 4 2	49, 115, 202, 271	1 (0%)
3	C	124/133 (93%)	0.68	11 (8%) 9 5	86, 130, 180, 247	0
All	All	1665/1693 (98%)	0.68	134 (8%) 12 6	49, 102, 190, 271	1 (0%)

All (134) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1196	LEU	12.7
2	B	1197	ALA	9.8
2	B	1193	GLY	9.1
2	B	1104	VAL	7.7
2	B	967	LEU	6.8
2	B	1122	ALA	6.1
2	B	1145	ALA	5.7
2	B	1148	LEU	5.7
2	B	1117	VAL	5.5
2	B	1243	LEU	5.3
2	B	1137	ASN	5.2
2	B	1076	THR	5.1
2	B	1159	GLU	5.1
2	B	1524	ASP	4.9
2	B	1080	ALA	4.9
2	B	1120	GLU	4.8
2	B	1216	ASP	4.8
1	A	665	PRO	4.7
2	B	1101	CYS	4.7
2	B	1320	ARG	4.6
2	B	1194	TYR	4.5
2	B	1236	ALA	4.5
2	B	1189	VAL	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	1300	LEU	4.2
2	B	1663	ASN	4.2
2	B	1152	GLN	4.2
2	B	1144	THR	4.2
2	B	1100	LEU	4.1
2	B	1106	TRP	4.1
2	B	1115	ASP	4.1
2	B	987	VAL	4.0
2	B	1142	ALA	4.0
2	B	1146	PHE	3.9
2	B	986	PRO	3.9
2	B	1306	LYS	3.8
2	B	1218	ASN	3.8
2	B	1305	SER	3.8
2	B	1217	LYS	3.6
2	B	1203	LYS	3.6
2	B	1240	LEU	3.6
2	B	1138	GLU	3.6
3	C	39	GLY	3.6
2	B	945	ARG	3.6
2	B	1354	ASP	3.5
2	B	1181	MET	3.5
2	B	1081	TYR	3.5
2	B	1211	LEU	3.4
2	B	1155	LYS	3.4
2	B	1207	LEU	3.4
2	B	1212	THR	3.4
2	B	943	ALA	3.3
3	C	30	GLY	3.3
2	B	1523	SER	3.3
1	A	481	ARG	3.3
2	B	1324	THR	3.3
2	B	1321	SER	3.3
2	B	1208	ASN	3.3
1	A	463	THR	3.2
1	A	461	LEU	3.2
2	B	955	GLU	3.2
1	A	459	SER	3.1
2	B	942	VAL	3.1
2	B	1328	GLU	3.0
3	C	87	MET	3.0
1	A	462	ARG	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	1180	TYR	3.0
2	B	1251	PRO	3.0
2	B	1107	LEU	3.0
2	B	946	THR	2.9
2	B	1158	CYS	2.9
2	B	1162	VAL	2.9
2	B	1215	LYS	2.8
1	A	460	VAL	2.8
3	C	8	LEU	2.8
2	B	1248	PHE	2.8
2	B	1201	ARG	2.8
2	B	985	THR	2.7
2	B	1066	PHE	2.7
2	B	1095	ILE	2.7
2	B	1342	LEU	2.6
2	B	1234	SER	2.6
2	B	1105	LYS	2.6
2	B	1220	TRP	2.6
2	B	1325	LYS	2.6
2	B	1176	LEU	2.6
2	B	1121	ASP	2.6
2	B	1195	ALA	2.6
2	B	1309	HIS	2.6
1	A	64	GLY	2.6
2	B	1498	GLU	2.6
3	C	126	SER	2.6
2	B	1149	ILE	2.5
2	B	1525	ASP	2.5
2	B	1318	LEU	2.5
2	B	1275	VAL	2.5
2	B	1109	LEU	2.5
2	B	1085	VAL	2.5
2	B	1278	ALA	2.4
2	B	990	MET	2.4
1	A	464	GLU	2.4
3	C	17	GLN	2.4
3	C	19	GLY	2.4
1	A	42	MET	2.4
2	B	1135	ASN	2.4
2	B	1237	LEU	2.3
3	C	86	GLN	2.3
2	B	965	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	1340	GLY	2.3
1	A	664	GLN	2.3
2	B	1184	GLN	2.3
3	C	98	TYR	2.3
2	B	1199	MET	2.3
2	B	1200	GLY	2.3
2	B	784	ASN	2.2
2	B	1070	VAL	2.2
2	B	1114	PRO	2.2
2	B	1169	ILE	2.2
2	B	1178	ALA	2.2
2	B	1063	SER	2.2
2	B	950	GLU	2.2
1	A	472	ASN	2.2
2	B	1229	ASN	2.2
2	B	1476	ALA	2.2
2	B	1132	GLY	2.1
2	B	1319	LEU	2.1
3	C	15	LEU	2.1
2	B	1190	ALA	2.1
2	B	1241	LEU	2.0
2	B	1330	PHE	2.0
2	B	1097	SER	2.0
2	B	1316	ALA	2.0
3	C	21	SER	2.0
2	B	1130	ILE	2.0
2	B	1303	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

6.5 Other polymers [i](#)

There are no such residues in this entry.