



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 01:16 PM EDT

PDB ID : 2QJE
Title : Crystal structure of the complex of Bovine C-lobe with Amygdalin at 2.3A resolution
Authors : Mir, R.; Jain, R.; Kumar, S.; Sinha, M.; Singh, N.; Sharma, S.; Kaur, P.; Srinivasan, A.; Bhushan, A.; Singh, T.P.
Deposited on : 2007-07-07
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 73%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 6%; margin-top: 5px;">6% 73% 21% . .</p>
2	B	3	<div style="display: flex; align-items: center;"> <div style="width: 67%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 6%; margin-top: 5px;">67% 33%</p>
2	C	3	<div style="display: flex; align-items: center;"> <div style="width: 33%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: orange; margin-right: 5px;"></div> </div> <p style="margin-left: 6%; margin-top: 5px;">33% 67%</p>
3	D	2	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 10px; background-color: orange; margin-right: 5px;"></div> </div> <p style="margin-left: 6%; margin-top: 5px;">100%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	1	-	-	X	-
3	TVY	D	1	-	-	-	X
3	Z8T	D	2	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 2966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	2593	1613	453	506	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0
2	C	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called beta-L-glucopyranose-(1-6)-(1S)-2-amino-1-phenylethyl alpha-D-mannopyranoside.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	2	32	20	1	11	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
5	A	2	2	2	0	0

- Molecule 6 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
6	A	1	1	1	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	1	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	237	Total 237	O 237	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lactotransferrin




- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-L-glucopyranose-(1-6)-(1S)-2-amino-1-phenylethyl alpha-D-mannopyranoside



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.37Å 50.37Å 65.97Å 90.00° 107.71° 90.00°	Depositor
Resolution (Å)	39.30 – 2.30 39.30 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.0 (39.30-2.30) 92.1 (39.30-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.29Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.223 , 0.227 0.217 , 0.256	Depositor DCC
R_{free} test set	846 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtrriage
Anisotropy	0.212	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.048 for l,-k,h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2966	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TVY, NAG, BMA, CO3, Z8T, FE, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	2/2640 (0.1%)	0.82	7/3575 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	684	CYS	N-CA	6.26	1.58	1.46
1	A	347	TRP	CB-CG	-6.01	1.39	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	684	CYS	N-CA-CB	13.79	135.41	110.60
1	A	683	ALA	CB-CA-C	-8.72	97.02	110.10
1	A	653	GLY	N-CA-C	-6.79	96.12	113.10
1	A	683	ALA	C-N-CA	6.78	138.65	121.70
1	A	684	CYS	N-CA-C	-5.81	95.31	111.00
1	A	545	ASN	CA-CB-CG	-5.54	101.22	113.40
1	A	532	CYS	CA-CB-SG	5.04	123.06	114.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	660	TYR	Sidechain
1	A	683	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2593	0	2510	90	0
2	B	39	0	34	2	0
2	C	39	0	34	10	0
3	D	32	0	0	4	0
4	A	14	0	13	0	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	5	0	0	0	0
8	A	4	0	0	1	0
9	A	237	0	0	22	0
All	All	2966	0	2591	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:ARG:HG2	1:A:344:ARG:HH11	1.14	1.12
1:A:429:PRO:HB2	3:D:2:Z8T:O6	1.54	1.07
1:A:434:LEU:CD1	2:C:1:NAG:H81	1.87	1.05
1:A:434:LEU:HD11	2:C:1:NAG:H81	1.42	1.01
1:A:359:GLN:HG2	9:A:760:HOH:O	1.71	0.91
1:A:343:THR:HA	1:A:606:HIS:NE2	1.85	0.90
1:A:584:ALA:HB3	2:C:1:NAG:H82	1.52	0.89
1:A:671:ASN:HA	9:A:915:HOH:O	1.72	0.89
9:A:733:HOH:O	2:C:3:BMA:H61	1.76	0.85
1:A:430:THR:HB	1:A:594:ASN:ND2	1.92	0.85
1:A:671:ASN:ND2	2:B:1:NAG:H3	1.93	0.84
1:A:663:THR:HG23	9:A:914:HOH:O	1.79	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:THR:HB	1:A:594:ASN:HD22	1.44	0.82
1:A:674:LYS:HD3	9:A:915:HOH:O	1.79	0.82
1:A:565:LYS:HD3	1:A:567:GLU:H	1.45	0.81
1:A:514:LYS:HD3	9:A:755:HOH:O	1.80	0.80
1:A:425:CYS:HB3	9:A:814:HOH:O	1.86	0.74
1:A:434:LEU:HD12	2:C:1:NAG:H81	1.71	0.73
1:A:344:ARG:HH11	1:A:344:ARG:CG	1.98	0.73
1:A:478:THR:HG22	1:A:480:SER:H	1.54	0.72
1:A:434:LEU:HD12	2:C:1:NAG:C8	2.19	0.72
1:A:434:LEU:CD1	2:C:1:NAG:C8	2.66	0.72
1:A:638:ASN:HD22	1:A:643:ASP:H	1.42	0.66
1:A:613:HIS:HB2	9:A:878:HOH:O	1.95	0.66
1:A:684:CYS:O	1:A:685:ALA:HB2	1.96	0.65
1:A:474:ILE:O	1:A:478:THR:HB	1.95	0.65
1:A:654:ARG:HB3	9:A:773:HOH:O	1.97	0.64
1:A:343:THR:HA	1:A:606:HIS:CE1	2.34	0.62
1:A:344:ARG:HG2	1:A:344:ARG:NH1	1.94	0.61
1:A:681:LEU:O	1:A:682:GLU:HB2	1.98	0.61
1:A:638:ASN:ND2	1:A:643:ASP:H	1.98	0.61
1:A:400:TYR:CZ	1:A:404:LYS:HE2	2.36	0.60
1:A:671:ASN:O	1:A:674:LYS:HG2	2.01	0.60
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.37	0.60
1:A:422:SER:HB3	9:A:870:HOH:O	2.02	0.59
1:A:636:THR:O	1:A:636:THR:HG23	2.02	0.58
1:A:426:VAL:HG23	9:A:814:HOH:O	2.02	0.58
1:A:654:ARG:HD3	9:A:773:HOH:O	2.03	0.58
1:A:633:LYS:NZ	9:A:913:HOH:O	2.27	0.57
1:A:461:VAL:O	1:A:462:ASP:HB2	2.04	0.57
1:A:584:ALA:CB	2:C:1:NAG:H82	2.31	0.57
1:A:626:PRO:HA	1:A:630:CYS:SG	2.45	0.57
1:A:671:ASN:HD22	2:B:1:NAG:H3	1.67	0.56
1:A:565:LYS:HD2	1:A:567:GLU:HB2	1.89	0.55
1:A:342:TYR:O	1:A:606:HIS:NE2	2.39	0.55
1:A:682:GLU:OE2	1:A:682:GLU:HA	2.07	0.55
1:A:400:TYR:OH	1:A:404:LYS:HE2	2.07	0.55
1:A:652:GLY:O	3:D:1:TVY:C12	2.55	0.55
1:A:420:HIS:HB3	1:A:423:LEU:HD11	1.89	0.54
1:A:363:GLN:NE2	9:A:886:HOH:O	2.40	0.53
1:A:420:HIS:O	1:A:423:LEU:CD1	2.57	0.53
9:A:733:HOH:O	2:C:3:BMA:C6	2.43	0.53
1:A:603:ARG:HD2	9:A:734:HOH:O	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:546:ASP:HA	2:C:1:NAG:H62	1.90	0.53
1:A:466:GLY:N	8:A:694:CO3:O3	2.40	0.53
1:A:684:CYS:O	1:A:685:ALA:CB	2.56	0.53
1:A:471:MET:HE1	1:A:487:PHE:HE2	1.75	0.52
1:A:653:GLY:O	1:A:654:ARG:C	2.47	0.52
1:A:681:LEU:HD22	1:A:684:CYS:HA	1.92	0.50
1:A:415:ARG:HG2	1:A:415:ARG:HH11	1.75	0.50
1:A:345:VAL:HA	1:A:390:ASP:OD2	2.11	0.50
1:A:478:THR:CG2	1:A:480:SER:HB3	2.42	0.50
1:A:651:LEU:O	3:D:1:TVY:C12	2.59	0.50
1:A:662:GLY:O	1:A:666:VAL:HG23	2.12	0.50
1:A:423:LEU:HD11	1:A:428:ARG:HG3	1.94	0.50
1:A:423:LEU:HD12	1:A:423:LEU:O	2.12	0.50
1:A:565:LYS:HD3	1:A:567:GLU:N	2.22	0.50
1:A:606:HIS:CE1	9:A:727:HOH:O	2.66	0.49
1:A:420:HIS:O	1:A:423:LEU:HD12	2.13	0.49
1:A:455:LYS:NZ	9:A:905:HOH:O	2.45	0.49
1:A:565:LYS:CD	1:A:567:GLU:H	2.22	0.48
1:A:514:LYS:HG2	9:A:825:HOH:O	2.13	0.48
1:A:429:PRO:CB	3:D:2:Z8T:O6	2.45	0.47
1:A:344:ARG:CG	1:A:344:ARG:NH1	2.65	0.47
1:A:636:THR:O	1:A:636:THR:CG2	2.64	0.46
1:A:675:CYS:O	1:A:676:SER:HB3	2.17	0.45
1:A:603:ARG:CD	9:A:734:HOH:O	2.65	0.45
1:A:345:VAL:HG13	1:A:390:ASP:HB2	1.98	0.45
1:A:548:VAL:HG11	1:A:581:VAL:HG11	1.99	0.44
1:A:379:ASP:O	1:A:383:LEU:HG	2.18	0.43
1:A:460:ALA:HB3	1:A:463:ARG:HD3	1.98	0.43
1:A:415:ARG:HH11	1:A:415:ARG:CG	2.31	0.43
1:A:488:SER:HA	1:A:500:ARG:HH21	1.83	0.43
1:A:606:HIS:HB2	9:A:852:HOH:O	2.17	0.43
1:A:478:THR:HG22	1:A:480:SER:N	2.30	0.42
1:A:345:VAL:HG21	1:A:607:VAL:HA	2.02	0.42
1:A:681:LEU:O	1:A:682:GLU:CB	2.66	0.42
1:A:404:LYS:HD3	1:A:685:ALA:C	2.40	0.41
1:A:498:LYS:HE2	1:A:498:LYS:HB2	1.84	0.41
1:A:553:ASN:HD21	1:A:565:LYS:HE2	1.85	0.41
1:A:606:HIS:CG	9:A:852:HOH:O	2.73	0.41
1:A:423:LEU:HD13	1:A:428:ARG:HB2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	336/348 (97%)	320 (95%)	14 (4%)	2 (1%)	25 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	652	GLY
1	A	682	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	281/289 (97%)	262 (93%)	19 (7%)	16 21

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	344	ARG
1	A	363	GLN
1	A	414	ASN
1	A	415	ARG
1	A	423	LEU
1	A	424	ASP
1	A	478	THR
1	A	504	LEU
1	A	515	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	559	ASP
1	A	565	LYS
1	A	583	GLU
1	A	591	VAL
1	A	631	LEU
1	A	633	LYS
1	A	635	GLU
1	A	636	THR
1	A	664	GLU
1	A	682	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	414	ASN
1	A	489	GLN
1	A	594	ASN
1	A	621	ASN
1	A	638	ASN
1	A	671	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.46	0	17,19,21	0.67	0
2	NAG	B	2	2	14,14,15	0.55	0	17,19,21	0.83	0
2	BMA	B	3	2	11,11,12	0.42	0	15,15,17	0.29	0
2	NAG	C	1	1,2	14,14,15	0.61	0	17,19,21	1.04	1 (5%)
2	NAG	C	2	2	14,14,15	0.56	0	17,19,21	0.78	0
2	BMA	C	3	2	11,11,12	1.00	1 (9%)	15,15,17	1.53	3 (20%)
3	TVY	D	1	3	22,22,22	0.63	0	28,30,30	1.02	2 (7%)
3	Z8T	D	2	3	11,11,12	0.66	0	15,15,17	1.10	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
3	TVY	D	1	3	-	4/12/32/32	0/2/2/2
3	Z8T	D	2	3	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	BMA	C4-C5	2.12	1.57	1.53

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	BMA	C1-C2-C3	-4.15	104.57	109.67
2	C	3	BMA	C3-C4-C5	2.91	115.43	110.24
2	C	1	NAG	C3-C4-C5	2.71	115.08	110.24
3	D	2	Z8T	C1-O5-C5	2.67	115.81	112.19
3	D	1	TVY	C1-O5-C5	2.39	118.38	113.69
3	D	1	TVY	O3-C3-C2	-2.34	104.94	110.35
2	C	3	BMA	O5-C1-C2	-2.02	107.65	110.77

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	D	1	TVY	O5-C1-O1-C13
2	C	2	NAG	C8-C7-N2-C2
2	B	3	BMA	O5-C5-C6-O6
2	C	2	NAG	O7-C7-N2-C2
3	D	1	TVY	O1-C13-C15-N17
3	D	1	TVY	C9-C13-O1-C1
3	D	1	TVY	C9-C13-C15-N17
2	B	3	BMA	C4-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	2	0
3	D	2	Z8T	2	0
2	C	3	BMA	2	0
2	C	1	NAG	8	0
3	D	1	TVY	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	CO3	A	694	6	2,3,3	0.74	0	2,3,3	0.05	0
4	NAG	A	1	1	14,14,15	0.49	0	17,19,21	0.80	1 (5%)
7	SO4	A	693	-	4,4,4	0.25	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	NAG	C2-N2-C7	-2.26	119.68	122.90

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1	NAG	C4-C5-C6-O6
4	A	1	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	694	CO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/348 (97%)	0.29	20 (5%) 22 28	16, 32, 59, 76	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	683	ALA	6.6
1	A	342	TYR	6.6
1	A	422	SER	4.7
1	A	482	ALA	4.2
1	A	676	SER	4.1
1	A	420	HIS	3.9
1	A	343	THR	3.6
1	A	416	LYS	3.6
1	A	423	LEU	3.4
1	A	674	LYS	3.0
1	A	344	ARG	2.9
1	A	660	TYR	2.8
1	A	685	ALA	2.7
1	A	623	LYS	2.6
1	A	684	CYS	2.5
1	A	635	GLU	2.4
1	A	483	PHE	2.1
1	A	636	THR	2.1
1	A	345	VAL	2.1
1	A	367	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	Z8T	D	2	11/12	0.44	0.65	20,20,20,20	11
3	TVY	D	1	21/21	0.45	0.62	20,20,20,20	21
2	BMA	B	3	11/12	0.62	0.23	76,77,77,77	0
2	NAG	C	1	14/15	0.71	0.28	61,63,65,68	0
2	BMA	C	3	11/12	0.71	0.35	81,87,89,93	0
2	NAG	C	2	14/15	0.80	0.16	71,72,74,77	0
2	NAG	B	2	14/15	0.85	0.28	69,71,72,74	0
2	NAG	B	1	14/15	0.89	0.17	56,58,61,65	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	FE	A	692	1/1	0.78	0.14	27,27,27,27	0
4	NAG	A	1	14/15	0.82	0.27	58,61,61,61	0
8	CO3	A	694	4/4	0.92	0.20	19,20,21,22	0
5	ZN	A	690	1/1	0.93	0.13	37,37,37,37	0
5	ZN	A	691	1/1	0.96	0.13	39,39,39,39	0
7	SO4	A	693	5/5	0.97	0.14	70,70,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.