



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 09:44 pm GMT

PDB ID : 8QM3  
Title : formaldehyde-inhibited [FeFe]-hydrogenase I from Clostridium pasteurianum (CpI)  
Authors : Duan, J.; Hofmann, E.  
Deposited on : 2023-09-21  
Resolution : 1.53 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	 2% 97%
1	B	584	 7% 96%

## 2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 19244 atoms, of which 8984 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Iron hydrogenase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	574	9012	2841	4473	779	872	47	0	11	0
1	B	572	9055	2861	4488	783	877	46	0	14	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	575	SER	-	expression tag	UNP P29166
A	576	ALA	-	expression tag	UNP P29166
A	577	TRP	-	expression tag	UNP P29166
A	578	SER	-	expression tag	UNP P29166
A	579	HIS	-	expression tag	UNP P29166
A	580	PRO	-	expression tag	UNP P29166
A	581	GLN	-	expression tag	UNP P29166
A	582	PHE	-	expression tag	UNP P29166
A	583	GLU	-	expression tag	UNP P29166
A	584	LYS	-	expression tag	UNP P29166
B	575	SER	-	expression tag	UNP P29166
B	576	ALA	-	expression tag	UNP P29166
B	577	TRP	-	expression tag	UNP P29166
B	578	SER	-	expression tag	UNP P29166
B	579	HIS	-	expression tag	UNP P29166
B	580	PRO	-	expression tag	UNP P29166
B	581	GLN	-	expression tag	UNP P29166
B	582	PHE	-	expression tag	UNP P29166
B	583	GLU	-	expression tag	UNP P29166
B	584	LYS	-	expression tag	UNP P29166

- Molecule 2 is dicarbonyl[bis(cyanide-kappaC)]-mu-(iminodimethanethiolatato-1kappaS:2kappaS)-mu-(oxomethylidene)diiron(2+) (three-letter code: 402) (formula: C<sub>7</sub>H<sub>5</sub>Fe<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	S	0	1
			17	7	2	3	3	2		
2	B	1	Total	C	Fe	N	O	S	0	1
			17	7	2	3	3	2		

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula:  $\text{Fe}_4\text{S}_4$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			4	2	2		
4	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 5 is FORMYL GROUP (three-letter code: FOR) (formula: CH<sub>2</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C 1 1	0	1
5	A	1	Total C 1 1	0	0
5	A	1	Total C O 2 1 1	0	1
5	A	1	Total C O 4 2 2	0	1
5	A	1	Total C O 2 1 1	0	1
5	B	1	Total C O 2 1 1	0	1
5	B	1	Total C 1 1	0	1
5	B	1	Total C 1 1	0	0
5	B	1	Total C O 4 2 2	0	1
5	B	1	Total C O 2 1 1	0	1

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C H O 13 3 7 3	0	0
6	A	1	Total C H O 14 3 8 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C H O 14 3 8 3	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	3	Total Cl 3 3	0	1
7	B	4	Total Cl 4 4	0	1

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total Mg 2 2	0	0
8	B	3	Total Mg 3 3	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	503	Total 503	O 503	0	1
9	B	489	Total 489	O 489	0	4




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Iron hydrogenase 1

Chain A: 2% 97%



- Molecule 1: Iron hydrogenase 1

Chain B: 7% 96%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.29Å 69.77Å 102.64Å 90.00° 102.74° 90.00°	Depositor
Resolution (Å)	44.95 – 1.53 44.95 – 1.53	Depositor EDS
% Data completeness (in resolution range)	97.0 (44.95-1.53) 97.1 (44.95-1.53)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 1.53Å)	Xtrriage
Refinement program	PHENIX (1.19.1_4122: ???)	Depositor
R, $R_{free}$	0.162 , 0.194 0.162 , 0.194	Depositor DCC
$R_{free}$ test set	8794 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtrriage
Anisotropy	0.270	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 57.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	19244	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, SF4, FOR, FES, CL, 402

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/4629	0.69	1/6229 (0.0%)
1	B	0.54	0/4650	0.71	1/6257 (0.0%)
All	All	0.53	0/9279	0.70	2/12486 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	404	ASP	CB-CG-OD2	7.05	124.64	118.30
1	A	404	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4539	4473	4475	0	0
1	B	4567	4488	4502	0	0
2	A	17	0	5	0	0
2	B	17	0	5	0	0
3	A	32	0	0	0	0
3	B	32	0	0	0	0
4	A	4	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	0	0	0
5	A	10	0	0	0	0
5	B	10	0	0	0	0
6	A	12	15	15	0	0
6	B	12	8	16	0	0
7	A	3	0	0	0	0
7	B	4	0	0	0	0
8	A	2	0	0	0	0
8	B	3	0	0	0	0
9	A	503	0	0	0	0
9	B	489	0	0	0	0
All	All	10260	8984	9018	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/584 (100%)	566 (97%)	19 (3%)	0	100	100
1	B	585/584 (100%)	566 (97%)	19 (3%)	0	100	100
All	All	1170/1168 (100%)	1132 (97%)	38 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	503/499 (101%)	493 (98%)	10 (2%)	55	24
1	B	504/499 (101%)	493 (98%)	11 (2%)	52	21
All	All	1007/998 (101%)	986 (98%)	21 (2%)	62	22

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ARG
1	A	117	LYS
1	A	119	ARG
1	A	122	LYS
1	A	211	MET
1	A	299[A]	CYS
1	A	299[B]	CYS
1	A	299[C]	CYS
1	A	366	GLN
1	A	515	LYS
1	B	2	LYS
1	B	4	ILE
1	B	81	GLU
1	B	104	ARG
1	B	299[A]	CYS
1	B	299[B]	CYS
1	B	299[C]	CYS
1	B	387	MET
1	B	417	PHE
1	B	573[A]	LYS
1	B	573[B]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 40 ligands modelled in this entry, 4 are modelled with single atom and 12 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	FOR	B	607[A]	1	0,1,1	-	-	-		
2	402	B	601[A]	1	13,19,19	2.10	5 (38%)	2,36,36	3.10	1 (50%)
5	FOR	A	612[B]	1	0,1,1	-	-	-		
5	FOR	A	611[A]	1	0,1,1	-	-	-		
3	SF4	A	604	1	0,12,12	-	-	-		
5	FOR	B	612[A]	1	0,1,1	-	-	-		
4	FES	A	606	1	0,4,4	-	-	-		
6	GOL	A	610	-	5,5,5	0.78	0	5,5,5	0.87	0
6	GOL	B	609	-	5,5,5	0.60	0	5,5,5	1.03	0
3	SF4	B	602	1	0,12,12	-	-	-		
2	402	A	601[A]	1	13,19,19	2.65	8 (61%)	2,36,36	2.45	1 (50%)
6	GOL	B	611	-	5,5,5	1.21	0	5,5,5	1.30	0
5	FOR	B	613[B]	-	0,1,1	-	-	-		
5	FOR	A	612[A]	1	0,1,1	-	-	-		
3	SF4	B	605	1	0,12,12	-	-	-		
3	SF4	B	603	1	0,12,12	-	-	-		
5	FOR	B	612[B]	1	0,1,1	-	-	-		
3	SF4	A	603	1	0,12,12	-	-	-		
5	FOR	A	613[B]	-	0,1,1	-	-	-		
3	SF4	A	605	1	0,12,12	-	-	-		
6	GOL	A	608	-	5,5,5	1.41	1 (20%)	5,5,5	1.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SF4	B	604	1	0,12,12	-	-	-		
4	FES	B	606	1	0,4,4	-	-	-		
3	SF4	A	602	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	402	A	601[A]	1	-	-	0/5/3/3
3	SF4	A	603	1	-	-	0/6/5/5
6	GOL	B	611	-	-	2/4/4/4	-
2	402	B	601[A]	1	-	-	0/5/3/3
6	GOL	B	609	-	-	2/4/4/4	-
6	GOL	A	610	-	-	0/4/4/4	-
3	SF4	A	604	1	-	-	0/6/5/5
3	SF4	A	605	1	-	-	0/6/5/5
3	SF4	B	602	1	-	-	0/6/5/5
6	GOL	A	608	-	-	2/4/4/4	-
4	FES	A	606	1	-	-	0/1/1/1
4	FES	B	606	1	-	-	0/1/1/1
3	SF4	B	604	1	-	-	0/6/5/5
3	SF4	B	605	1	-	-	0/6/5/5
3	SF4	B	603	1	-	-	0/6/5/5
3	SF4	A	602	1	-	-	0/6/5/5

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601[A]	402	C4-N4	5.03	1.24	1.15
2	B	601[A]	402	S1-FE1	4.45	2.32	2.26
2	A	601[A]	402	O3-C3	4.30	1.26	1.15
2	B	601[A]	402	S2-FE1	3.51	2.31	2.26
2	A	601[A]	402	S1-FE1	3.27	2.31	2.26
2	A	601[A]	402	C2-S2	3.23	1.90	1.85
2	B	601[A]	402	O3-C3	3.04	1.23	1.15
6	A	608	GOL	O2-C2	-3.01	1.34	1.43
2	B	601[A]	402	C4-N4	2.62	1.19	1.15
2	A	601[A]	402	S1-FE2	2.50	2.29	2.26
2	A	601[A]	402	O5-C5	2.44	1.21	1.17
2	A	601[A]	402	C6-N6	-2.20	1.11	1.15

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601[A]	402	C3-FE1	-2.12	1.72	1.79
2	B	601[A]	402	C3-FE1	-2.05	1.73	1.79

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601[A]	402	S2-C2-N1	-4.24	105.51	117.18
2	A	601[A]	402	S2-C2-N1	-3.29	108.12	117.18

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	B	609	GOL	C1-C2-C3-O3
6	B	609	GOL	O2-C2-C3-O3
6	A	608	GOL	O1-C1-C2-O2
6	A	608	GOL	O1-C1-C2-C3
6	B	611	GOL	C1-C2-C3-O3
6	B	611	GOL	O2-C2-C3-O3




There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	574/584 (98%)	-0.17	9 (1%) 72 77	15, 23, 46, 67	0
1	B	572/584 (97%)	0.25	40 (6%) 16 18	14, 21, 64, 94	0
All	All	1146/1168 (98%)	0.04	49 (4%) 35 40	14, 22, 52, 94	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	9	VAL	12.5
1	B	5	ILE	11.3
1	B	3	THR	9.8
1	B	12	ASN	9.1
1	B	4	ILE	8.8
1	B	8	GLY	8.5
1	B	2	LYS	8.4
1	B	10	GLN	8.3
1	B	28	ILE	7.2
1	B	69	GLY	7.0
1	B	11	PHE	6.6
1	B	6	ILE	5.8
1	B	27	ASN	5.7
1	B	71	ILE	5.7
1	B	22	PHE	5.4
1	B	66	ILE	5.0
1	B	53	VAL	4.1
1	B	72	ILE	4.0
1	A	513	ASN	3.7
1	B	14	ASP	3.4
1	B	56	THR	3.4
1	B	459	VAL	3.4
1	B	54	GLU	3.4
1	B	65	LEU	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	70	MET	3.2
1	B	67	GLU	3.1
1	B	518	GLU	3.1
1	B	68	ASP	3.0
1	B	573[A]	LYS	3.0
1	B	57	GLY	2.9
1	A	130	THR	2.9
1	A	562	GLY	2.7
1	A	514	PRO	2.7
1	B	75	ASN	2.7
1	B	24	ARG	2.5
1	A	558	LYS	2.4
1	B	26	ASN	2.3
1	B	13	THR	2.3
1	A	567	ILE	2.3
1	B	29	ASP	2.3
1	B	7	ASN	2.2
1	B	19	ILE	2.2
1	B	58	LEU	2.2
1	A	498	ALA	2.1
1	B	174	LYS	2.1
1	A	563	ARG	2.0
1	B	173	GLY	2.0
1	B	74	THR	2.0
1	A	129	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	GOL	B	609	6/6	0.78	0.12	35,42,49,62	0
5	FOR	A	613[B]	2/2	0.85	0.15	24,24,24,27	2
6	GOL	A	610	6/6	0.91	0.29	27,51,66,66	0
5	FOR	A	609	1/2	0.91	0.12	37,37,37,37	0
5	FOR	A	611[A]	2/2	0.92	0.16	20,20,20,21	2
5	FOR	A	607[C]	1/2	0.93	0.16	24,24,24,24	1
5	FOR	B	608[C]	1/2	0.95	0.11	18,18,18,18	1
5	FOR	B	610	1/2	0.95	0.10	32,32,32,32	0
6	GOL	B	611	6/6	0.95	0.11	19,29,35,37	0
6	GOL	A	608	6/6	0.96	0.06	20,27,33,33	0
5	FOR	B	613[B]	2/2	0.96	0.11	22,22,22,24	2
7	CL	B	619	1/1	0.96	0.04	31,31,31,31	0
5	FOR	A	612[B]	2/2	0.97	0.24	21,21,21,22	2
5	FOR	A	612[A]	2/2	0.97	0.24	22,22,22,27	2
5	FOR	B	612[A]	2/2	0.97	0.14	23,23,23,27	2
5	FOR	B	612[B]	2/2	0.97	0.14	26,26,26,27	2
7	CL	A	618	1/1	0.97	0.04	37,37,37,37	0
5	FOR	B	607[A]	2/2	0.97	0.12	18,18,18,18	2
8	MG	B	616	1/1	0.97	0.04	27,27,27,27	0
8	MG	B	615	1/1	0.98	0.04	29,29,29,29	0
7	CL	A	617	1/1	0.98	0.03	31,31,31,31	0
3	SF4	A	605	8/8	0.99	0.07	18,19,21,21	0
3	SF4	B	603	8/8	0.99	0.10	14,15,15,15	0
4	FES	B	606	4/4	0.99	0.09	20,21,22,22	0
7	CL	A	614[B]	1/1	0.99	0.09	16,16,16,16	1
2	402	A	601[A]	17/17	0.99	0.11	10,18,21,22	0
2	402	B	601[A]	17/17	0.99	0.12	12,15,17,19	0
7	CL	B	614[B]	1/1	0.99	0.10	15,15,15,15	1
7	CL	B	618	1/1	0.99	0.03	27,27,27,27	0
3	SF4	A	602	8/8	0.99	0.12	15,16,17,18	0
7	CL	B	620	1/1	0.99	0.07	27,27,27,27	0
8	MG	A	616	1/1	0.99	0.02	19,19,19,19	0
3	SF4	A	603	8/8	0.99	0.10	15,16,17,17	0
3	SF4	A	604	8/8	0.99	0.10	15,16,16,17	0
8	MG	B	617	1/1	0.99	0.03	24,24,24,24	0
8	MG	A	615	1/1	1.00	0.02	24,24,24,24	0
3	SF4	B	602	8/8	1.00	0.12	13,14,14,15	0
3	SF4	B	604	8/8	1.00	0.09	15,15,16,16	0
3	SF4	B	605	8/8	1.00	0.08	16,17,18,19	0
4	FES	A	606	4/4	1.00	0.09	17,17,17,18	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.















**Electron density around FOR A 607 (C):**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FOR B 608 (C):**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FOR B 613 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FOR A 612 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FOR A 612 (A):**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FOR B 612 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FOR B 612 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FOR B 607 (A):**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 402 A 601 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.