



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 7, 2022 – 12:32 PM EDT

PDB ID : 2QNU  
Title : Crystal structure of PA0076 from Pseudomonas aeruginosa PAO1 at 2.05 Å resolution  
Authors : Filippova, E.V.; Chruszcz, M.; Skarina, T.; Kagan, O.; Cymborowski, M.; Savchenko, A.; Edwards, A.M.; Joachimiak, A.; Minor, W.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2007-07-19  
Resolution : 2.05 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	 2% 83% 11% 7%
1	B	226	 5% 84% 12% ..

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein PA0076.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	211	Total 1580	C 1011	N 273	O 290	S 3	Se 3	0	0	0
1	B	220	Total 1652	C 1063	N 282	O 301	S 3	Se 3	0	0	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total	O	0	0
			68	68		
4	B	52	Total	O	0	0
			52	52		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein PA0076



- Molecule 1: Uncharacterized protein PA0076



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.26Å 125.41Å 49.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.32 – 2.05 43.82 – 2.05	Depositor EDS
% Data completeness (in resolution range)	(Not available) (46.32-2.05) 96.3 (43.82-2.05)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	0.19	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	29.19 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.183 , 0.234 0.214 , 0.253	Depositor DCC
$R_{free}$ test set	1226 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.2	Xtrriage
Anisotropy	0.664	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 57.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3374	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, ACT, MLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.01	0/1610	0.87	2/2193 (0.1%)
1	B	0.98	2/1684 (0.1%)	0.98	6/2293 (0.3%)
All	All	0.99	2/3294 (0.1%)	0.93	8/4486 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	93	CYS	CB-SG	-5.85	1.72	1.81
1	B	130	GLU	CG-CD	5.48	1.60	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	209	ARG	NE-CZ-NH1	10.80	125.70	120.30
1	A	159	ARG	NE-CZ-NH1	9.68	125.14	120.30
1	A	159	ARG	NE-CZ-NH2	-9.20	115.70	120.30
1	B	209	ARG	NE-CZ-NH2	-9.03	115.78	120.30
1	B	13	ARG	NE-CZ-NH2	-8.17	116.21	120.30
1	B	13	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	B	186	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	B	186	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	0	1538	16	0
1	B	1652	0	1615	18	0
2	A	8	0	6	1	0
2	B	4	0	3	0	0
3	A	10	0	14	1	0
4	A	68	0	0	0	0
4	B	52	0	0	1	0
All	All	3374	0	3176	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:LEU:HD12	1:A:17:VAL:HG21	1.37	1.06
1:B:153:ILE:HD13	1:B:157:LEU:HD23	1.60	0.81
1:B:153:ILE:HD13	1:B:157:LEU:CD2	2.26	0.65
1:B:125:PRO:HA	1:B:126:GLU:HG3	1.80	0.64
1:A:58:TRP:CZ3	1:A:196:MSE:HE2	2.35	0.61
1:B:5:GLY:HA3	1:B:21:LEU:HD13	1.87	0.56
1:B:110:GLY:O	1:B:114:GLN:HG3	2.07	0.54
1:A:198:TYR:CD2	1:A:202:PRO:HG3	2.45	0.51
1:A:198:TYR:CE2	1:B:157:LEU:HD13	2.49	0.48
1:B:19:ARG:HD2	4:B:268:HOH:O	2.12	0.48
1:B:170:ALA:O	1:B:174:GLN:HG3	2.14	0.47
1:A:6:PHE:CE2	1:A:19:ARG:HB2	2.49	0.47
1:A:80:ILE:HD12	1:A:86:TYR:CE1	2.51	0.46
1:A:198:TYR:CE1	1:B:157:LEU:HB2	2.50	0.46
1:A:116:GLU:OE1	1:A:186:ARG:NH1	2.48	0.46
1:A:209:ARG:CZ	1:B:153:ILE:CG2	2.94	0.46
1:A:58:TRP:CH2	1:A:196:MSE:HE2	2.51	0.45
3:A:229:PGE:H2	3:A:229:PGE:O3	2.15	0.45
1:A:57:LEU:HD12	1:A:57:LEU:N	2.31	0.45
1:B:151:SER:O	1:B:152:LEU:HD12	2.16	0.45
1:B:6:PHE:CE2	1:B:19:ARG:HG3	2.53	0.44
1:B:13:ARG:HG3	1:B:17:VAL:HG11	2.01	0.43
1:A:196:MSE:HE1	1:A:210:PHE:CD1	2.54	0.43
1:A:139:LEU:HD12	1:A:140:PRO:HD2	2.01	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:GLU:HB3	1:A:28:PRO:HD3	2.02	0.42
1:A:37:MSE:HE2	1:A:87:PHE:HA	2.01	0.42
1:B:62:ILE:HB	1:B:72:VAL:HG13	2.02	0.42
1:B:125:PRO:CA	1:B:126:GLU:HG3	2.49	0.41
1:B:125:PRO:HA	1:B:126:GLU:HA	1.60	0.41
2:A:227:ACT:H1	1:B:175:HIS:HB2	2.03	0.40
1:A:209:ARG:CZ	1:B:153:ILE:HG21	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/226 (92%)	202 (97%)	6 (3%)	0	100	100
1	B	213/226 (94%)	207 (97%)	6 (3%)	0	100	100
All	All	421/452 (93%)	409 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/163 (95%)	154 (99%)	1 (1%)	86	87

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	164/163 (101%)	154 (94%)	10 (6%)	18	10
All	All	319/326 (98%)	308 (97%)	11 (3%)	37	30

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ARG
1	B	10	LEU
1	B	19	ARG
1	B	21	LEU
1	B	53	LEU
1	B	94	LEU
1	B	144	CYS
1	B	147	ARG
1	B	174	GLN
1	B	190	ARG
1	B	224	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	MLZ	B	9	1	8,9,10	0.47	0	4,9,11	1.02	0
1	MLZ	A	9	1	8,9,10	1.17	1 (12%)	4,9,11	2.20	2 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	B	9	1	-	1/7/8/10	-
1	MLZ	A	9	1	-	2/7/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	9	MLZ	CB-CA	2.21	1.56	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	MLZ	CM-NZ-CE	3.33	121.57	111.95
1	A	9	MLZ	CD-CG-CB	2.04	120.85	113.62

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	9	MLZ	CD-CE-NZ-CM
1	B	9	MLZ	CD-CE-NZ-CM
1	A	9	MLZ	CE-CD-CG-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACT	B	227	-	1,3,3	2.35	1 (100%)	0,3,3	-	-
2	ACT	A	227	-	1,3,3	0.08	0	0,3,3	-	-
2	ACT	A	228	-	1,3,3	2.15	1 (100%)	0,3,3	-	-
3	PGE	A	229	-	9,9,9	0.57	0	8,8,8	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	229	-	-	2/7/7/7	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	227	ACT	CH3-C	-2.35	1.45	1.48
2	A	228	ACT	CH3-C	2.15	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	229	PGE	C4-C3-O2-C2
3	A	229	PGE	O2-C3-C4-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	227	ACT	1	0
3	A	229	PGE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/226 (91%)	0.08	4 (1%) 66 71	30, 39, 52, 64	0
1	B	216/226 (95%)	0.22	11 (5%) 28 30	33, 41, 55, 71	0
All	All	423/452 (93%)	0.15	15 (3%) 44 48	30, 40, 54, 71	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	ALA	5.1
1	B	125	PRO	5.1
1	A	214	GLU	2.9
1	B	225	PRO	2.6
1	B	144	CYS	2.6
1	B	3	SER	2.5
1	B	153	ILE	2.4
1	B	224	ILE	2.4
1	A	213	GLY	2.3
1	B	147	ARG	2.2
1	A	190	ARG	2.2
1	B	146	PRO	2.1
1	B	157	LEU	2.1
1	B	152	LEU	2.1
1	B	143	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	MLZ	A	9	10/11	0.93	0.12	36,37,43,43	0
1	MLZ	B	9	10/11	0.94	0.11	34,35,47,48	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PGE	A	229	10/10	0.76	0.15	66,72,73,73	0
2	ACT	A	228	4/4	0.86	0.12	60,61,61,62	0
2	ACT	A	227	4/4	0.97	0.08	43,44,44,45	0
2	ACT	B	227	4/4	0.99	0.07	34,36,36,37	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.