



wwPDB X-ray Structure Validation Summary Report

Jun 17, 2024 – 07:00 PM EDT

PDB ID : 3QPQ
Title : Crystal structure of ANTI-TLR3 antibody C1068 FAB
Authors : Luo, J.; Obmolova, G.; Teplyakov, A.; Gilliland, G.L.
Deposited on : 2011-02-14
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	215	 85% 14% .
1	E	215	 91% 9%
1	I	215	 % 86% 13% .
1	L	215	 87% 13% .
2	D	223	 3% 86% 10% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	223	
2	H	223	
2	J	223	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	L	216	-	-	X	-
4	SO4	L	219	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C1068 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	215	Total 1691	C 1059	N 282	O 343	S 7	0	6	0
1	E	215	Total 1682	C 1054	N 280	O 341	S 7	0	5	0
1	C	215	Total 1677	C 1052	N 279	O 339	S 7	0	4	0
1	I	215	Total 1672	C 1049	N 279	O 337	S 7	0	3	0

- Molecule 2 is a protein called C1068 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	216	Total 1652	C 1049	N 275	O 321	S 7	0	2	0
2	F	221	Total 1710	C 1085	N 289	O 329	S 7	0	4	0
2	D	219	Total 1682	C 1067	N 284	O 324	S 7	0	2	0
2	J	217	Total 1659	C 1053	N 278	O 321	S 7	0	1	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total C O 6 3 3	0	0
3	L	1	Total C O 6 3 3	0	0
3	L	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0
3	F	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	I	1	Total C O 6 3 3	0	0
3	I	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	J	1	Total	C	O	0	0
			6	3	3		
3	J	1	Total	C	O	0	0
			6	3	3		
3	J	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	228	Total	O	0	0
			228	228		
5	H	222	Total	O	0	0
			222	222		
5	E	229	Total	O	0	0
			229	229		
5	F	189	Total	O	0	0
			189	189		
5	C	229	Total	O	0	0
			229	229		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	223	Total 223	O 223	0	0
5	I	184	Total 184	O 184	0	0
5	J	197	Total 197	O 197	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C1068 light chain

Chain L:  87% 13%




- Molecule 1: C1068 light chain

Chain E:  91% 9%




- Molecule 1: C1068 light chain

Chain C:  85% 14%




- Molecule 1: C1068 light chain

Chain I:  86% 13%




- Molecule 2: C1068 heavy chain

Chain H:  86% 10% 4%




HIS
HIS
HIS


- Molecule 2: C1068 heavy chain



- Molecule 2: C1068 heavy chain



- Molecule 2: C1068 heavy chain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.48Å 136.94Å 83.25Å 90.00° 114.95° 90.00°	Depositor
Resolution (Å)	44.55 – 1.90 44.55 – 1.90	Depositor EDS
% Data completeness (in resolution range)	89.2 (44.55-1.90) 89.2 (44.55-1.90)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 1.89Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_601), CNX	Depositor
R, R_{free}	0.201 , 0.256 0.188 , 0.243	Depositor DCC
R_{free} test set	5917 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtrriage
Anisotropy	0.424	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	0.000 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15233	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9489e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.39	0/1727	0.55	0/2342
1	E	0.38	0/1735	0.56	0/2352
1	I	0.38	0/1719	0.54	0/2331
1	L	0.41	0/1744	0.56	0/2365
2	D	0.35	0/1726	0.53	0/2359
2	F	0.34	0/1761	0.53	0/2406
2	H	0.35	0/1693	0.56	1/2314 (0.0%)
2	J	0.33	0/1698	0.55	0/2321
All	All	0.37	0/13803	0.55	1/18790 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	184	LEU	CA-CB-CG	6.23	129.64	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1677	0	1622	30	0
1	E	1682	0	1626	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	1672	0	1618	19	0
1	L	1691	0	1634	21	0
2	D	1682	0	1648	25	0
2	F	1710	0	1679	46	0
2	H	1652	0	1627	27	0
2	J	1659	0	1629	40	0
3	C	36	0	48	6	0
3	D	6	0	8	1	0
3	E	6	0	8	2	0
3	F	6	0	8	0	0
3	I	12	0	16	3	0
3	J	18	0	24	7	0
3	L	18	0	24	6	0
4	L	5	0	0	3	0
5	C	229	0	0	6	1
5	D	223	0	0	3	0
5	E	229	0	0	4	0
5	F	189	0	0	7	0
5	H	222	0	0	7	1
5	I	184	0	0	4	0
5	J	197	0	0	6	1
5	L	228	0	0	9	1
All	All	15233	0	13219	224	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 224 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:216:GOL:H31	5:I:484:HOH:O	1.46	1.15
3:L:216:GOL:H2	5:L:526:HOH:O	0.95	1.12
2:H:97[B]:THR:HG21	2:H:106:PHE:HB3	1.45	0.99
2:J:97[B]:THR:HG21	2:J:106:PHE:HB3	1.47	0.95
2:D:221:HIS:O	2:D:222:HIS:HB2	1.68	0.93

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:510:HOH:O	5:J:483:HOH:O[2_646]	2.06	0.14
5:L:505:HOH:O	5:H:459:HOH:O[2_846]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	217/215 (101%)	211 (97%)	6 (3%)	0	100	100
1	E	218/215 (101%)	214 (98%)	4 (2%)	0	100	100
1	I	216/215 (100%)	211 (98%)	5 (2%)	0	100	100
1	L	219/215 (102%)	214 (98%)	5 (2%)	0	100	100
2	D	217/223 (97%)	207 (95%)	8 (4%)	2 (1%)	17	7
2	F	221/223 (99%)	214 (97%)	7 (3%)	0	100	100
2	H	214/223 (96%)	208 (97%)	6 (3%)	0	100	100
2	J	214/223 (96%)	209 (98%)	5 (2%)	0	100	100
All	All	1736/1752 (99%)	1688 (97%)	46 (3%)	2 (0%)	51	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	221	HIS
2	D	220	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	192/188 (102%)	186 (97%)	6 (3%)	40	32
1	E	193/188 (103%)	188 (97%)	5 (3%)	46	39
1	I	191/188 (102%)	185 (97%)	6 (3%)	40	32
1	L	194/188 (103%)	190 (98%)	4 (2%)	53	48
2	D	192/194 (99%)	189 (98%)	3 (2%)	62	60
2	F	196/194 (101%)	191 (97%)	5 (3%)	46	39
2	H	189/194 (97%)	187 (99%)	2 (1%)	73	73
2	J	189/194 (97%)	183 (97%)	6 (3%)	39	30
All	All	1536/1528 (100%)	1499 (98%)	37 (2%)	47	43

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	76	ASN
2	J	184	LEU
1	I	105	GLU
2	J	111	GLN
2	F	102	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	199	HIS
1	I	38	GLN
2	D	170	HIS
1	I	3	GLN
1	I	148	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	D	1	2	7,8,9	1.89	1 (14%)	9,10,12	2.51	5 (55%)
2	PCA	J	1	2	7,8,9	1.83	1 (14%)	9,10,12	2.17	5 (55%)
2	PCA	H	1	2	7,8,9	1.93	1 (14%)	9,10,12	2.06	5 (55%)
2	PCA	F	1	2	7,8,9	1.95	1 (14%)	9,10,12	2.36	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	D	1	2	-	0/0/11/13	0/1/1/1
2	PCA	J	1	2	-	0/0/11/13	0/1/1/1
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1
2	PCA	F	1	2	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	PCA	CD-N	5.04	1.47	1.34
2	H	1	PCA	CD-N	4.96	1.46	1.34
2	D	1	PCA	CD-N	4.81	1.46	1.34
2	J	1	PCA	CD-N	4.72	1.46	1.34

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	PCA	CB-CA-C	-4.43	106.58	112.66
2	F	1	PCA	CB-CA-C	-3.63	107.68	112.66
2	J	1	PCA	CB-CA-C	-3.29	108.14	112.66
2	F	1	PCA	OE-CD-CG	-3.10	121.19	126.72
2	D	1	PCA	CA-N-CD	-3.05	103.14	113.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	I	216	-	5,5,5	0.48	0	5,5,5	0.40	0
3	GOL	F	224	-	5,5,5	0.41	0	5,5,5	0.36	0
3	GOL	C	217	-	5,5,5	0.39	0	5,5,5	0.53	0
3	GOL	L	217	-	5,5,5	0.32	0	5,5,5	0.52	0
3	GOL	I	217	-	5,5,5	0.43	0	5,5,5	0.20	0
3	GOL	C	220	-	5,5,5	0.44	0	5,5,5	0.32	0
3	GOL	E	216	-	5,5,5	0.48	0	5,5,5	0.15	0
3	GOL	C	216	-	5,5,5	0.41	0	5,5,5	0.43	0
3	GOL	C	221	-	5,5,5	0.45	0	5,5,5	0.52	0
3	GOL	J	226	-	5,5,5	0.42	0	5,5,5	0.48	0
3	GOL	L	216	-	5,5,5	0.36	0	5,5,5	0.38	0
3	GOL	D	224	-	5,5,5	0.35	0	5,5,5	0.51	0
3	GOL	C	219	-	5,5,5	0.37	0	5,5,5	0.39	0
3	GOL	J	224	-	5,5,5	0.39	0	5,5,5	0.37	0
4	SO4	L	219	-	4,4,4	0.28	0	6,6,6	0.14	0
3	GOL	L	218	-	5,5,5	0.38	0	5,5,5	0.33	0
3	GOL	C	218	-	5,5,5	0.41	0	5,5,5	0.32	0
3	GOL	J	225	-	5,5,5	0.37	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	I	216	-	-	2/4/4/4	-
3	GOL	F	224	-	-	2/4/4/4	-
3	GOL	C	217	-	-	1/4/4/4	-
3	GOL	L	217	-	-	1/4/4/4	-
3	GOL	I	217	-	-	4/4/4/4	-
3	GOL	C	220	-	-	2/4/4/4	-
3	GOL	E	216	-	-	4/4/4/4	-
3	GOL	C	216	-	-	0/4/4/4	-
3	GOL	C	221	-	-	4/4/4/4	-
3	GOL	J	226	-	-	2/4/4/4	-
3	GOL	L	216	-	-	1/4/4/4	-
3	GOL	D	224	-	-	0/4/4/4	-
3	GOL	C	219	-	-	4/4/4/4	-
3	GOL	J	224	-	-	2/4/4/4	-
3	GOL	L	218	-	-	0/4/4/4	-
3	GOL	C	218	-	-	2/4/4/4	-
3	GOL	J	225	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	216	GOL	O1-C1-C2-C3
3	F	224	GOL	O1-C1-C2-C3
3	C	218	GOL	O1-C1-C2-C3
3	C	219	GOL	C1-C2-C3-O3
3	C	220	GOL	O1-C1-C2-C3

There are no ring outliers.

13 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	216	GOL	2	0
3	C	217	GOL	1	0
3	L	217	GOL	2	0
3	I	217	GOL	1	0
3	E	216	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	226	GOL	3	0
3	L	216	GOL	4	0
3	D	224	GOL	1	0
3	C	219	GOL	2	0
3	J	224	GOL	3	0
4	L	219	SO4	3	0
3	C	218	GOL	3	0
3	J	225	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	215/215 (100%)	-0.03	0 100 100	17, 26, 41, 63	0
1	E	215/215 (100%)	0.03	1 (0%) 91 92	15, 25, 41, 56	0
1	I	215/215 (100%)	0.12	2 (0%) 84 85	18, 29, 46, 64	0
1	L	215/215 (100%)	-0.04	1 (0%) 91 92	15, 23, 41, 56	0
2	D	218/223 (97%)	0.21	6 (2%) 53 56	18, 28, 53, 86	0
2	F	220/223 (98%)	0.53	14 (6%) 19 22	18, 32, 54, 91	0
2	H	215/223 (96%)	0.20	8 (3%) 41 44	16, 26, 46, 84	0
2	J	216/223 (96%)	0.47	10 (4%) 32 35	19, 31, 58, 82	0
All	All	1729/1752 (98%)	0.19	42 (2%) 59 62	15, 27, 48, 91	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	137	THR	5.5
2	H	138	SER	5.4
2	J	220	HIS	5.3
2	J	42	GLY	5.1
2	H	134	SER	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	F	1	8/9	0.83	0.14	32,42,48,49	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PCA	H	1	8/9	0.87	0.10	27,36,45,51	0
2	PCA	J	1	8/9	0.91	0.15	34,35,37,38	0
2	PCA	D	1	8/9	0.94	0.14	21,32,46,50	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	C	219	6/6	0.51	0.31	51,61,67,71	0
3	GOL	L	216	6/6	0.66	0.24	34,42,45,49	0
3	GOL	E	216	6/6	0.70	0.25	39,45,50,50	0
3	GOL	C	221	6/6	0.70	0.25	30,44,56,56	0
3	GOL	D	224	6/6	0.78	0.24	40,51,61,72	0
3	GOL	I	217	6/6	0.79	0.18	36,42,49,53	0
3	GOL	I	216	6/6	0.82	0.20	30,40,45,47	0
3	GOL	L	217	6/6	0.83	0.17	35,48,49,58	0
3	GOL	J	226	6/6	0.85	0.19	42,51,53,57	0
3	GOL	C	220	6/6	0.86	0.18	47,49,52,55	0
3	GOL	J	225	6/6	0.87	0.17	27,43,45,45	0
3	GOL	C	218	6/6	0.88	0.31	24,38,42,54	0
3	GOL	C	217	6/6	0.89	0.14	33,41,42,48	0
3	GOL	C	216	6/6	0.89	0.12	31,35,41,42	0
3	GOL	J	224	6/6	0.91	0.12	34,48,49,56	0
3	GOL	L	218	6/6	0.92	0.14	37,43,50,55	0
3	GOL	F	224	6/6	0.94	0.09	30,35,40,44	0
4	SO4	L	219	5/5	0.97	0.35	11,32,37,37	5

6.5 Other polymers [i](#)

There are no such residues in this entry.