



## Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 05:06 pm BST

PDB ID : 4R1N  
Title : Crystal structure of (S)-3-hydroxybutylryl-CoA dehydrogenase form the n-butanol synthesizing bacterium, Clostridium butyricum.  
Authors : Kim, E.J.; Kim, S.W.; Kim, K.J.  
Deposited on : 2014-08-07  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	
1	B	282	
1	C	282	
1	D	282	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8756 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxybutyryl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	282	2130	1352	353	410	15	0	0	0
1	B	282	2130	1352	353	410	15	0	0	0
1	C	282	2130	1352	353	410	15	0	0	0
1	D	282	2130	1352	353	410	15	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	56	Total	O	0	0
			56	56		
2	B	60	Total	O	0	0
			60	60		
2	C	63	Total	O	0	0
			63	63		
2	D	57	Total	O	0	0
			57	57		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase




- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase




- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase



- Molecule 1: 3-hydroxybutyryl-CoA dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.47Å 146.47Å 202.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.90 – 1.80 30.92 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.4 (30.90-1.80) 92.4 (30.92-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.09 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.207 , 0.240 0.215 , 0.247	Depositor DCC
$R_{free}$ test set	6846 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8756	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0904e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.97	2/2156 (0.1%)	0.97	6/2901 (0.2%)
1	B	0.93	1/2156 (0.0%)	0.96	3/2901 (0.1%)
1	C	0.95	0/2156	0.98	7/2901 (0.2%)
1	D	0.92	0/2156	0.95	6/2901 (0.2%)
All	All	0.94	3/8624 (0.0%)	0.96	22/11604 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	GLU	CD-OE2	-6.05	1.19	1.25
1	B	34	GLU	CG-CD	5.69	1.60	1.51
1	A	204	GLU	CD-OE1	-5.31	1.19	1.25

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	ARG	NE-CZ-NH2	-7.83	116.39	120.30
1	C	265	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	A	75	MET	CG-SD-CE	-7.05	88.92	100.20
1	D	261	ARG	CG-CD-NE	-6.58	97.97	111.80
1	D	236	ASP	CB-CG-OD1	6.41	124.06	118.30
1	B	261	ARG	CG-CD-NE	-6.39	98.38	111.80
1	D	103	ASP	CB-CG-OD1	6.33	124.00	118.30
1	D	272	LYS	CB-CA-C	-6.26	97.88	110.40
1	C	255	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	C	272	LYS	CB-CA-C	-6.17	98.06	110.40
1	B	82	ASP	CB-CG-OD1	-5.99	112.91	118.30
1	C	231	ASP	CB-CG-OD1	5.89	123.60	118.30
1	B	130	ARG	N-CA-CB	-5.64	100.44	110.60
1	A	82	ASP	CB-CG-OD1	-5.56	113.30	118.30
1	D	103	ASP	CB-CG-OD2	-5.55	113.30	118.30
1	C	74	ASP	CB-CG-OD2	-5.51	113.34	118.30
1	C	236	ASP	CB-CG-OD1	5.47	123.22	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	130	ARG	N-CA-CB	-5.35	100.97	110.60
1	C	77	LEU	CA-CB-CG	5.29	127.47	115.30
1	A	251	ASP	CB-CG-OD2	-5.27	113.56	118.30
1	A	252	THR	OG1-CB-CG2	5.24	122.06	110.00
1	A	265	ARG	NE-CZ-NH2	-5.22	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2130	0	2211	20	0
1	B	2130	0	2211	24	0
1	C	2130	0	2211	17	0
1	D	2130	0	2211	30	0
2	A	56	0	0	0	0
2	B	60	0	0	0	0
2	C	63	0	0	1	0
2	D	57	0	0	2	0
All	All	8756	0	8844	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:MET:HE2	1:B:194:MET:HE2	1.28	1.16
1:C:194:MET:HE2	1:D:194:MET:HE2	1.24	1.15
1:B:22:ALA:HA	1:B:65:ILE:HD12	1.33	1.03
1:C:194:MET:HE2	1:D:194:MET:CE	2.04	0.87
1:A:194:MET:HE2	1:B:194:MET:CE	2.08	0.81
1:C:194:MET:CE	1:D:194:MET:HE2	2.09	0.80
1:D:33:LYS:HE3	1:D:34:GLU:H	1.47	0.79

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:MET:CE	1:B:194:MET:HE2	2.12	0.78
1:D:130:ARG:HG2	1:D:133:LYS:HB2	1.72	0.71
1:C:194:MET:CE	1:D:194:MET:CE	2.69	0.71
1:B:130:ARG:HG2	1:B:133:LYS:HB2	1.75	0.69
1:A:22:ALA:HA	1:A:65:ILE:HD12	1.78	0.66
1:A:194:MET:CE	1:B:194:MET:CE	2.72	0.65
1:C:259:LEU:C	1:C:259:LEU:HD23	2.19	0.63
1:B:259:LEU:C	1:B:259:LEU:HD23	2.20	0.61
1:A:118:SER:HB2	1:A:244:VAL:HG21	1.81	0.61
1:C:89:ILE:HD11	1:C:94:ILE:HD13	1.81	0.61
1:A:282:LYS:OXT	1:A:282:LYS:CG	2.49	0.60
1:D:95:LYS:HE2	1:D:119:LEU:HD12	1.83	0.60
1:A:190:ILE:CG2	1:B:194:MET:HG3	2.34	0.57
1:D:95:LYS:CE	1:D:119:LEU:HD12	2.35	0.56
1:D:118:SER:HB2	1:D:244:VAL:HG21	1.86	0.56
1:C:272:LYS:HG3	2:C:349:HOH:O	2.06	0.56
1:B:118:SER:HB2	1:B:244:VAL:HG21	1.88	0.56
1:D:22:ALA:HA	1:D:65:ILE:HD12	1.87	0.55
1:A:7:LEU:N	1:A:7:LEU:HD12	2.21	0.55
1:C:89:ILE:HD12	1:C:91:ASN:HB3	1.89	0.55
1:C:113:ALA:HA	1:C:135:ILE:O	2.07	0.55
1:D:33:LYS:HE3	1:D:34:GLU:N	2.19	0.55
1:B:201:ILE:HG23	1:B:206:VAL:HB	1.88	0.55
1:C:194:MET:HG3	1:D:190:ILE:CG2	2.37	0.54
1:C:201:ILE:HG23	1:C:206:VAL:HB	1.88	0.54
1:B:56:LYS:HE2	1:B:56:LYS:HA	1.92	0.52
1:C:190:ILE:CG2	1:D:194:MET:HG3	2.39	0.51
1:D:259:LEU:C	1:D:259:LEU:HD23	2.32	0.50
1:B:65:ILE:HD13	1:B:68:ARG:NH2	2.27	0.50
1:B:89:ILE:HD11	1:B:94:ILE:HD13	1.94	0.48
1:D:33:LYS:HE3	1:D:34:GLU:HG3	1.95	0.48
1:D:52:VAL:O	1:D:54:LYS:O	2.31	0.48
1:A:172:ILE:HG13	1:A:174:LYS:HG2	1.96	0.47
1:C:35:GLU:OE2	1:C:39:ARG:NH2	2.46	0.47
1:A:54:LYS:CG	1:A:54:LYS:O	2.63	0.47
1:A:113:ALA:HA	1:A:135:ILE:O	2.15	0.46
1:B:33:LYS:HB2	1:B:36:PHE:CD2	2.50	0.46
1:A:65:ILE:HD13	1:A:68:ARG:NH2	2.30	0.46
1:D:1:MET:SD	1:D:168:MET:HE2	2.54	0.46
1:A:282:LYS:HG2	1:A:282:LYS:OXT	2.16	0.46
1:D:121:ILE:HG12	2:D:349:HOH:O	2.15	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:259:LEU:O	1:B:259:LEU:HD23	2.15	0.46
1:D:33:LYS:CE	1:D:34:GLU:HG3	2.46	0.46
1:B:61:ASP:O	1:B:65:ILE:HG12	2.16	0.45
1:D:113:ALA:HA	1:D:135:ILE:O	2.17	0.44
1:C:253:LYS:HB2	1:D:253:LYS:HA	1.99	0.44
1:C:89:ILE:HD11	1:C:94:ILE:HG21	1.99	0.44
1:B:84:VAL:HG11	1:B:102:LEU:HD13	1.98	0.44
1:D:194:MET:HE2	1:D:194:MET:HB2	1.85	0.44
1:B:74:ASP:OD1	1:B:76:LYS:HD3	2.18	0.44
1:C:194:MET:HG3	1:D:190:ILE:HG23	1.99	0.44
1:B:58:THR:O	1:B:61:ASP:HB2	2.18	0.44
1:A:281:SER:O	1:A:282:LYS:C	2.56	0.43
1:D:135:ILE:HG21	1:D:165:VAL:HG21	2.01	0.43
1:A:135:ILE:HG21	1:A:165:VAL:HG21	2.01	0.42
1:B:152:ILE:HD13	1:B:179:VAL:HB	2.01	0.42
1:D:259:LEU:O	1:D:259:LEU:HD23	2.19	0.42
1:D:52:VAL:C	1:D:54:LYS:O	2.58	0.42
1:A:201:ILE:HG23	1:A:206:VAL:HB	2.02	0.42
1:D:224:MET:HE2	1:D:224:MET:HB2	1.86	0.42
1:D:281:SER:HA	1:D:282:LYS:HA	1.88	0.42
1:B:56:LYS:CE	1:B:56:LYS:HA	2.51	0.41
1:A:8:GLY:O	1:A:13:GLY:HA3	2.21	0.41
1:B:89:ILE:HD11	1:B:94:ILE:CD1	2.51	0.40
1:A:282:LYS:OXT	1:A:282:LYS:HG3	2.22	0.40
1:A:194:MET:HG3	1:B:190:ILE:CG2	2.51	0.40
1:C:143:ALA:N	1:C:144:PRO:CD	2.85	0.40
1:D:272:LYS:HG3	2:D:318:HOH:O	2.21	0.40
1:D:89:ILE:HD11	1:D:94:ILE:CD1	2.51	0.40
1:B:224:MET:HB2	1:B:224:MET:HE2	1.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/282 (99%)	271 (97%)	9 (3%)	0	100	100
1	B	280/282 (99%)	270 (96%)	10 (4%)	0	100	100
1	C	280/282 (99%)	274 (98%)	6 (2%)	0	100	100
1	D	280/282 (99%)	273 (98%)	7 (2%)	0	100	100
All	All	1120/1128 (99%)	1088 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/228 (100%)	217 (95%)	11 (5%)	25	11
1	B	228/228 (100%)	219 (96%)	9 (4%)	32	17
1	C	228/228 (100%)	218 (96%)	10 (4%)	28	14
1	D	228/228 (100%)	221 (97%)	7 (3%)	40	25
All	All	912/912 (100%)	875 (96%)	37 (4%)	30	16

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLU
1	A	46	LYS
1	A	63	GLU
1	A	66	LEU
1	A	75	MET
1	A	90	GLU
1	A	93	LYS
1	A	108	PRO
1	A	168	MET
1	A	194	MET
1	A	252	THR
1	B	1	MET
1	B	2	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	39	ARG
1	B	54	LYS
1	B	66	LEU
1	B	70	SER
1	B	93	LYS
1	B	121	ILE
1	B	224	MET
1	C	12	MET
1	C	63	GLU
1	C	66	LEU
1	C	75	MET
1	C	90	GLU
1	C	93	LYS
1	C	118	SER
1	C	168	MET
1	C	252	THR
1	C	279	ASP
1	D	33	LYS
1	D	66	LEU
1	D	90	GLU
1	D	93	LYS
1	D	149	VAL
1	D	252	THR
1	D	282	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	282/282 (100%)	-0.15	18 (6%) 19 15	20, 32, 65, 94	0
1	B	282/282 (100%)	-0.10	11 (3%) 39 33	20, 33, 57, 104	0
1	C	282/282 (100%)	-0.23	5 (1%) 68 64	22, 30, 51, 103	0
1	D	282/282 (100%)	-0.15	6 (2%) 63 59	22, 34, 57, 95	0
All	All	1128/1128 (100%)	-0.16	40 (3%) 44 38	20, 32, 58, 104	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	281	SER	9.8
1	B	282	LYS	9.1
1	B	281	SER	8.5
1	A	281	SER	7.8
1	C	282	LYS	7.8
1	A	282	LYS	7.2
1	C	281	SER	7.0
1	A	53	ALA	5.8
1	D	282	LYS	5.2
1	D	32	ILE	4.5
1	A	54	LYS	3.7
1	B	39	ARG	3.4
1	A	55	GLU	3.3
1	A	50	LYS	3.3
1	C	53	ALA	3.0
1	A	52	VAL	2.9
1	D	34	GLU	2.9
1	A	35	GLU	2.7
1	A	39	ARG	2.7
1	D	39	ARG	2.7
1	A	58	THR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	53	ALA	2.6
1	B	56	LYS	2.6
1	A	57	ILE	2.5
1	A	63	GLU	2.4
1	A	46	LYS	2.3
1	A	49	SER	2.3
1	A	56	LYS	2.2
1	B	54	LYS	2.2
1	B	55	GLU	2.2
1	C	35	GLU	2.2
1	B	36	PHE	2.1
1	B	35	GLU	2.1
1	A	60	ALA	2.1
1	C	52	VAL	2.1
1	B	52	VAL	2.0
1	D	113	ALA	2.0
1	A	45	THR	2.0
1	A	33	LYS	2.0
1	B	33	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.