



# Full wwPDB NMR Structure Validation Report ⓘ

Jun 6, 2020 – 06:13 am BST

PDB ID : 6R14  
Title : Structure of kiteplatinated dsDNA  
Authors : Margiotta, N.; Papadia, P.; Kubicek, K.; Krejcikova, M.; Gkionis, K.; Sponer, J.  
Deposited on : 2019-03-13

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Ensemble composition and analysis

This entry contains 20 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 780 atoms, of which 285 are hydrogens and 0 are deuteriums.


- Molecule 1 is a DNA chain called Kiteplatinated DNA oligomer, chain A.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	12	371	113	136	40	71	11	0

- Molecule 2 is a DNA chain called Kiteplatinated DNA oligomer, chain B.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	B	12	386	118	135	53	69	11	0

- Molecule 3 is Kiteplatin (three-letter code: LN8) (formula:  $C_6H_{14}Cl_2N_2Pt$ ) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms				
			Total	C	H	N	Pt
3	A	1	23	6	14	2	1

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Kiteplatinated DNA oligomer, chain A



- Molecule 2: Kiteplatinated DNA oligomer, chain B




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Kiteplatinated DNA oligomer, chain A




- Molecule 2: Kiteplatinated DNA oligomer, chain B




### 4.2.2 Score per residue for model 2

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  83% 17%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  8% 67% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

### 4.2.3 Score per residue for model 3

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  83% 17%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  67% 33%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


### 4.2.4 Score per residue for model 4

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

### 4.2.5 Score per residue for model 5

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  83% 17%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  8% 50% 42%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


### 4.2.6 Score per residue for model 6

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  75% 17% 8%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 G12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  83% 17%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


### 4.2.7 Score per residue for model 7

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

### 4.2.8 Score per residue for model 8

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  75% 25%




- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  67% 33%




### 4.2.9 Score per residue for model 9

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  75% 25%



- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%



### 4.2.10 Score per residue for model 10

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%



- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  67% 33%






#### 4.2.11 Score per residue for model 11

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%

G1 C2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


#### 4.2.12 Score per residue for model 12

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  75% 25%

G1 C2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

#### 4.2.13 Score per residue for model 13

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  83% 17%

G1 C2 T3 C4 A5 G6 G7 C8 C9 T10 C11 C12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  67% 33%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


#### 4.2.14 Score per residue for model 14

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%

G1 C2 T3 C4 A5 G6 C7 C8 C9 T10 C11 C12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

#### 4.2.15 Score per residue for model 15

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  100%

G1 C2 T3 C4 A5 G6 C7 C8 C9 T10 C11 C12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  67% 33%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

#### 4.2.16 Score per residue for model 16

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  92% 8%

G1 C2 T3 C4 A5 G6 C7 C8 C9 T10 C11 C12


- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  8% 50% 42%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


#### 4.2.17 Score per residue for model 17

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  83% 17%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 G12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


#### 4.2.18 Score per residue for model 18

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  100%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 G12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  83% 17%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24


#### 4.2.19 Score per residue for model 19

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  67% 25% 8%

G1 G2 T3 C4 A5 G6 G7 C8 C9 T10 C11 G12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  75% 25%

G13 G14 A15 G16 G17 C18 C19 T20 G21 A22 G23 G24

#### 4.2.20 Score per residue for model 20

- Molecule 1: Kiteplatinated DNA oligomer, chain A

Chain A:  100%

G1  
C2  
T3  
C4  
A5  
G6  
G7  
C8  
C9  
T10  
C11  
C12

- Molecule 2: Kiteplatinated DNA oligomer, chain B

Chain B:  8% 83% 8%

G13  
G14  
A15  
G16  
G17  
C18  
C19  
T20  
G21  
A22  
G23  
G24

## 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
Amber	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	200
Number of shifts mapped to atoms	200
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	40%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	235	136	136	0±0
All	All	9900	5700	5420	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:DC:C5	1:A:12:DC:C5	0.42	3.08	6	4

## 5.2 Torsion angles [i](#)

### 5.2.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.2.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.5 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
3	LN8	A	101	1	6,10,12	0.40±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	LN8	A	101	1	6,13,18	0.31±0.05	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LN8	A	101	1	-	-	0±0,1,2,2

There are no bond-length outliers.


There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.6 Other polymers [i](#)

There are no such molecules in this entry.

## 5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 40% for the well-defined parts and 40% for the entire structure.

### 6.1 Chemical shift list 1

File name: input\_cs.cif

Chemical shift list name: *kitePt.STAR*

#### 6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	200
Number of shifts mapped to atoms	200
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 6.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

#### 6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 40%, i.e. 191 atoms were assigned a chemical shift out of a possible 483. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	191/483 (40%)	191/291 (66%)	0/159 (0%)	0/33 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 40%, i.e. 191 atoms were assigned a chemical shift out of a possible 483. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	191/483 (40%)	191/291 (66%)	0/159 (0%)	0/33 (0%)

#### 6.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 6.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (kitePt.STAR). RCI is only applicable to proteins.