



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 08:19 PM EDT

PDB ID : 3R2C
Title : Crystal Structure of Antitermination Factors NusB and NusE in complex with BoxA RNA
Authors : Stagno, J.R.; Ji, X.
Deposited on : 2011-03-14
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)
RNA backbone	3102	1013 (2.42-1.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	 86% 7% • 7%
1	B	148	 86% 6% • 7%
2	J	83	 88% 7% 5%
2	K	83	 86% 10% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	R	12	 50% 25% 25%
3	S	12	 8% 58% 17% 8% 17%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 8328 atoms, of which 4054 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N utilization substance protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	138	2383	759	1209	203	211	1	0	5	0
1	B	137	2280	725	1158	194	202	1	0	0	0

- Molecule 2 is a protein called 30S ribosomal protein S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	J	79	1316	404	684	113	113	2	0	0	0
2	K	80	1355	415	703	118	117	2	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	47	SER	-	SEE REMARK 999	UNP O66430
K	47	SER	-	SEE REMARK 999	UNP O66430

- Molecule 3 is a RNA chain called 5'-R(*GP*GP*CP*UP*CP*CP*UP*UP*GP*GP*CP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				P
3	R	9	253	74	86	25	60	8	0	0	0
3	S	10	317	94	108	35	71	9	0	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	17	4	10	3	0	0
4	A	1	17	4	10	3	0	0
4	A	1	17	4	10	3	0	0
4	A	1	17	4	10	3	0	0
4	B	1	17	4	10	3	0	0
4	B	1	17	4	10	3	0	0
4	J	1	17	4	10	3	0	0
4	R	1	17	4	10	3	0	0
4	S	1	17	4	10	3	0	0
4	S	1	17	4	10	3	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	82	Total	O	0	0
			82	82		
6	B	66	Total	O	0	1
			67	67		
6	J	27	Total	O	0	0
			27	27		
6	K	26	Total	O	0	0
			26	26		
6	R	20	Total	O	0	0
			20	20		
6	S	18	Total	O	0	0
			18	18		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N utilization substance protein B





- Molecule 1: N utilization substance protein B




- Molecule 2: 30S ribosomal protein S10




- Molecule 2: 30S ribosomal protein S10



- Molecule 3: 5'-R(*GP*GP*CP*UP*CP*CP*UP*UP*GP*GP*CP*A)-3'



- Molecule 3: 5'-R(*GP*GP*CP*UP*CP*CP*UP*UP*GP*GP*CP*A)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.53Å 113.01Å 65.76Å 90.00° 102.87° 90.00°	Depositor
Resolution (Å)	35.66 – 1.90 35.66 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.9 (35.66-1.90) 93.9 (35.66-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.91Å)	Xtrriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.169 , 0.202 0.164 , 0.196	Depositor DCC
R_{free} test set	2160 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtrriage
Anisotropy	0.223	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8328	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1202	0.51	0/1613
1	B	0.39	0/1142	0.51	0/1533
2	J	0.32	0/640	0.58	0/861
2	K	0.32	0/660	0.51	0/887
3	R	0.56	0/184	1.11	0/285
3	S	0.60	0/232	1.00	0/360
All	All	0.40	0/4060	0.61	0/5539

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1174	1209	1208	7	0
1	B	1122	1158	1158	7	0
2	J	632	684	684	6	0
2	K	652	703	702	6	0
3	R	167	86	86	2	0
3	S	209	108	109	7	0
4	A	28	40	40	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	14	20	20	0	0
4	J	7	10	10	0	0
4	R	7	10	10	1	0
4	S	14	20	20	2	0
5	A	4	3	3	0	0
5	B	4	3	3	0	0
6	A	82	0	0	1	0
6	B	67	0	0	1	0
6	J	27	0	0	1	0
6	K	26	0	0	0	0
6	R	20	0	0	0	0
6	S	18	0	0	0	0
All	All	4274	4054	4053	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:S:10:G:P	3:S:10:G:H3'	2.13	0.88
3:S:10:G:C3'	3:S:10:G:OP1	2.34	0.76
3:S:10:G:OP1	3:S:10:G:O3'	2.07	0.71
3:S:10:G:H3'	3:S:10:G:OP1	1.92	0.69
2:K:4[B]:GLU:HG2	2:K:4[B]:GLU:O	1.96	0.64
3:S:10:G:P	3:S:10:G:C3'	2.88	0.61
2:J:79:VAL:HG12	2:J:80:LYS:N	2.23	0.52
1:B:99:GLU:O	1:B:99:GLU:HG3	2.09	0.52
1:A:118:LYS:HD3	3:R:8:U:O3'	2.09	0.52
2:K:71:LEU:HD23	2:K:77:VAL:HG21	1.92	0.52
1:B:121:LYS:HG3	3:S:6:C:H5''	1.93	0.51
1:B:132:LYS:HE3	4:S:14:PEG:O4	2.10	0.51
1:B:62:ASP:O	1:B:66:GLU:HG3	2.12	0.49
2:J:79:VAL:CG1	2:J:80:LYS:N	2.74	0.49
1:A:20:LEU:HD13	2:J:39:PRO:HG2	1.94	0.49
1:B:33:VAL:HG22	1:B:38:ILE:HD12	1.96	0.48
2:J:80:LYS:HE3	6:J:116:HOH:O	2.12	0.48
3:S:2:G:H21	4:S:13:PEG:H12	1.78	0.48
1:A:69:LEU:O	1:A:72[A]:TRP:HD1	1.97	0.47
3:R:4:U:O4	4:R:13:PEG:H32	2.15	0.45
2:K:10:LEU:HD12	2:K:10:LEU:N	2.32	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72[A]:TRP:CE3	1:A:76:ARG:HD2	2.53	0.43
2:J:17:LEU:HD21	2:J:72:PRO:HB2	2.01	0.43
2:K:71:LEU:HD12	2:K:71:LEU:HA	1.78	0.42
1:B:59:GLU:OE2	2:J:30:LYS:HE3	2.20	0.42
2:K:66:LEU:O	2:K:69:ILE:HG22	2.20	0.42
1:A:31:GLU:HB2	6:A:154:HOH:O	2.20	0.41
1:A:128:SER:OG	1:A:132:LYS:HE3	2.19	0.41
1:B:35:GLU:HG3	6:B:192:HOH:O	2.20	0.41
2:K:14:ASP:C	2:K:14:ASP:OD1	2.59	0.41
1:A:33:VAL:HG13	1:A:38:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/148 (95%)	140 (99%)	1 (1%)	0	100	100
1	B	135/148 (91%)	134 (99%)	1 (1%)	0	100	100
2	J	77/83 (93%)	77 (100%)	0	0	100	100
2	K	79/83 (95%)	78 (99%)	1 (1%)	0	100	100
All	All	432/462 (94%)	429 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/128 (96%)	121 (98%)	2 (2%)	62	60
1	B	117/128 (91%)	115 (98%)	2 (2%)	60	57
2	J	72/75 (96%)	72 (100%)	0	100	100
2	K	74/75 (99%)	71 (96%)	3 (4%)	30	21
All	All	386/406 (95%)	379 (98%)	7 (2%)	59	55

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	132	LYS
1	B	49	LYS
1	B	132	LYS
2	K	37	LYS
2	K	71	LEU
2	K	80	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	R	7/12 (58%)	1 (14%)	0
3	S	9/12 (75%)	1 (11%)	0
All	All	16/24 (66%)	2 (12%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	R	2	G
3	S	10	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	150	-	6,6,6	0.49	0	5,5,5	0.68	0
4	PEG	A	151	-	6,6,6	0.48	0	5,5,5	0.71	0
4	PEG	A	149	-	6,6,6	0.49	0	5,5,5	0.81	0
4	PEG	B	150	-	6,6,6	0.49	0	5,5,5	0.70	0
5	ACT	B	151	-	3,3,3	0.78	0	3,3,3	1.24	0
4	PEG	B	149	-	6,6,6	0.49	0	5,5,5	0.75	0
4	PEG	J	84	-	6,6,6	0.48	0	5,5,5	0.70	0
4	PEG	A	152	-	6,6,6	0.47	0	5,5,5	0.64	0
4	PEG	R	13	-	6,6,6	0.50	0	5,5,5	0.81	0
4	PEG	S	14	-	6,6,6	0.50	0	5,5,5	0.79	0
5	ACT	A	153	-	3,3,3	0.82	0	3,3,3	1.28	0
4	PEG	S	13	-	6,6,6	0.51	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	150	-	-	2/4/4/4	-
4	PEG	A	151	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	149	-	-	2/4/4/4	-
4	PEG	B	150	-	-	4/4/4/4	-
4	PEG	B	149	-	-	2/4/4/4	-
4	PEG	J	84	-	-	3/4/4/4	-
4	PEG	A	152	-	-	3/4/4/4	-
4	PEG	R	13	-	-	3/4/4/4	-
4	PEG	S	14	-	-	3/4/4/4	-
4	PEG	S	13	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	152	PEG	O2-C3-C4-O4
4	R	13	PEG	O2-C3-C4-O4
4	R	13	PEG	C4-C3-O2-C2
4	A	150	PEG	O1-C1-C2-O2
4	B	149	PEG	O1-C1-C2-O2
4	B	150	PEG	O2-C3-C4-O4
4	A	151	PEG	O2-C3-C4-O4
4	J	84	PEG	O1-C1-C2-O2
4	S	14	PEG	O1-C1-C2-O2
4	A	151	PEG	O1-C1-C2-O2
4	J	84	PEG	O2-C3-C4-O4
4	S	14	PEG	O2-C3-C4-O4
4	A	149	PEG	O2-C3-C4-O4
4	B	150	PEG	O1-C1-C2-O2
4	R	13	PEG	O1-C1-C2-O2
4	S	13	PEG	O2-C3-C4-O4
4	A	152	PEG	C1-C2-O2-C3
4	S	14	PEG	C4-C3-O2-C2
4	S	13	PEG	C1-C2-O2-C3
4	B	149	PEG	C4-C3-O2-C2
4	A	150	PEG	O2-C3-C4-O4
4	B	150	PEG	C1-C2-O2-C3
4	S	13	PEG	O1-C1-C2-O2
4	J	84	PEG	C4-C3-O2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	152	PEG	C4-C3-O2-C2
4	B	150	PEG	C4-C3-O2-C2
4	A	151	PEG	C4-C3-O2-C2
4	A	149	PEG	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	R	13	PEG	1	0
4	S	14	PEG	1	0
4	S	13	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/148 (93%)	-0.24	2 (1%) 75 77	13, 21, 49, 68	0
1	B	137/148 (92%)	-0.21	3 (2%) 62 64	15, 25, 53, 71	0
2	J	79/83 (95%)	-0.01	6 (7%) 13 15	16, 35, 65, 82	0
2	K	80/83 (96%)	0.06	3 (3%) 40 43	20, 39, 75, 78	0
3	R	9/12 (75%)	-0.32	0 100 100	25, 29, 55, 72	0
3	S	10/12 (83%)	-0.20	1 (10%) 7 8	25, 30, 62, 70	0
All	All	453/486 (93%)	-0.14	15 (3%) 46 49	13, 27, 61, 82	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	73	ALA	4.9
2	K	82	ARG	4.5
2	J	81	MET	4.3
2	J	73	ALA	3.5
2	K	4[A]	GLU	3.5
2	J	3	GLN	3.5
1	A	72[A]	TRP	2.9
2	J	71	LEU	2.7
3	S	10	G	2.4
2	J	70	SER	2.3
1	A	98	LYS	2.3
1	B	136	THR	2.3
1	B	98	LYS	2.2
2	J	4	GLU	2.1
1	B	137	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACT	A	153	4/4	0.67	0.14	45,54,61,64	0
4	PEG	A	151	7/7	0.69	0.15	62,75,87,89	0
4	PEG	B	149	7/7	0.77	0.19	52,67,89,89	0
4	PEG	R	13	7/7	0.80	0.28	40,53,69,69	0
4	PEG	A	150	7/7	0.84	0.12	56,68,82,82	0
4	PEG	S	13	7/7	0.85	0.18	35,42,56,64	0
4	PEG	A	149	7/7	0.87	0.17	39,47,64,64	0
5	ACT	B	151	4/4	0.87	0.16	66,73,93,93	0
4	PEG	S	14	7/7	0.88	0.15	42,55,66,66	0
4	PEG	B	150	7/7	0.89	0.19	52,63,83,85	0
4	PEG	A	152	7/7	0.90	0.17	35,47,59,59	0
4	PEG	J	84	7/7	0.93	0.10	47,64,82,98	0

6.5 Other polymers [i](#)

There are no such residues in this entry.