



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2024 – 05:57 PM EST

PDB ID : 3R31  
Title : Crystal structure of betaine aldehyde dehydrogenase from *Agrobacterium tumefaciens*  
Authors : Agarwal, R.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2011-03-15  
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	 5% 70% 20% • 7%
1	B	517	 8% 69% 21% • 7%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Betaine aldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	482	3575	2264	610	685	16	0	0	0
1	B	482	3558	2251	610	681	16	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q8UH56
A	2	VAL	-	expression tag	UNP Q8UH56
A	496	ALA	-	expression tag	UNP Q8UH56
A	497	GLU	-	expression tag	UNP Q8UH56
A	498	ASN	-	expression tag	UNP Q8UH56
A	499	LEU	-	expression tag	UNP Q8UH56
A	500	TYR	-	expression tag	UNP Q8UH56
A	501	PHE	-	expression tag	UNP Q8UH56
A	502	GLN	-	expression tag	UNP Q8UH56
A	503	SER	-	expression tag	UNP Q8UH56
A	504	HIS	-	expression tag	UNP Q8UH56
A	505	HIS	-	expression tag	UNP Q8UH56
A	506	HIS	-	expression tag	UNP Q8UH56
A	507	HIS	-	expression tag	UNP Q8UH56
A	508	HIS	-	expression tag	UNP Q8UH56
A	509	HIS	-	expression tag	UNP Q8UH56
A	510	TRP	-	expression tag	UNP Q8UH56
A	511	SER	-	expression tag	UNP Q8UH56
A	512	HIS	-	expression tag	UNP Q8UH56
A	513	PRO	-	expression tag	UNP Q8UH56
A	514	GLN	-	expression tag	UNP Q8UH56
A	515	PHE	-	expression tag	UNP Q8UH56
A	516	GLU	-	expression tag	UNP Q8UH56
A	517	LYS	-	expression tag	UNP Q8UH56
B	1	MET	-	expression tag	UNP Q8UH56

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	VAL	-	expression tag	UNP Q8UH56
B	496	ALA	-	expression tag	UNP Q8UH56
B	497	GLU	-	expression tag	UNP Q8UH56
B	498	ASN	-	expression tag	UNP Q8UH56
B	499	LEU	-	expression tag	UNP Q8UH56
B	500	TYR	-	expression tag	UNP Q8UH56
B	501	PHE	-	expression tag	UNP Q8UH56
B	502	GLN	-	expression tag	UNP Q8UH56
B	503	SER	-	expression tag	UNP Q8UH56
B	504	HIS	-	expression tag	UNP Q8UH56
B	505	HIS	-	expression tag	UNP Q8UH56
B	506	HIS	-	expression tag	UNP Q8UH56
B	507	HIS	-	expression tag	UNP Q8UH56
B	508	HIS	-	expression tag	UNP Q8UH56
B	509	HIS	-	expression tag	UNP Q8UH56
B	510	TRP	-	expression tag	UNP Q8UH56
B	511	SER	-	expression tag	UNP Q8UH56
B	512	HIS	-	expression tag	UNP Q8UH56
B	513	PRO	-	expression tag	UNP Q8UH56
B	514	GLN	-	expression tag	UNP Q8UH56
B	515	PHE	-	expression tag	UNP Q8UH56
B	516	GLU	-	expression tag	UNP Q8UH56
B	517	LYS	-	expression tag	UNP Q8UH56

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	117	Total	O	0	0
			117	117		
3	B	102	Total	O	0	0
			102	102		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Betaine aldehyde dehydrogenase



- Molecule 1: Betaine aldehyde dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.88Å 98.88Å 174.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.02 – 2.15 43.02 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.02-2.15) 99.9 (43.02-2.15)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.08 (at 2.14Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.194 , 0.245 0.197 , 0.244	Depositor DCC
$R_{free}$ test set	2767 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7360	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.12	6/3648 (0.2%)	0.95	8/4954 (0.2%)
1	B	1.03	4/3630 (0.1%)	0.88	3/4931 (0.1%)
All	All	1.08	10/7278 (0.1%)	0.92	11/9885 (0.1%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	480	GLU	CG-CD	5.90	1.60	1.51
1	A	169	TRP	CB-CG	5.68	1.60	1.50
1	B	452	CYS	CB-SG	5.21	1.91	1.82
1	A	71	MET	CB-CG	5.20	1.68	1.51
1	A	299	PHE	CE1-CZ	5.15	1.47	1.37

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	426	THR	CB-CA-C	-5.91	95.65	111.60
1	A	323	ASP	CB-CG-OD1	5.84	123.56	118.30
1	A	409	VAL	CB-CA-C	-5.72	100.52	111.40
1	A	63	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	468	ARG	NE-CZ-NH1	5.66	123.13	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3575	0	3543	91	0
1	B	3558	0	3519	109	0
2	A	4	0	6	1	0
2	B	4	0	6	0	0
3	A	117	0	0	3	0
3	B	102	0	0	5	0
All	All	7360	0	7074	199	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:MET:CE	1:A:45:LYS:HG2	1.66	1.24
1:A:426:THR:HG21	1:A:431:ARG:HD2	1.16	1.08
1:B:364:PRO:HG2	1:B:367:VAL:HG21	1.40	1.03
1:B:307:ARG:HG3	1:B:307:ARG:HH11	1.25	1.00
1:A:42:MET:HE1	1:A:45:LYS:HG2	1.42	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	480/517 (93%)	463 (96%)	15 (3%)	2 (0%)	34 29
1	B	480/517 (93%)	458 (95%)	21 (4%)	1 (0%)	47 45
All	All	960/1034 (93%)	921 (96%)	36 (4%)	3 (0%)	41 36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	ASP
1	B	113	ASP
1	A	40	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/400 (91%)	337 (93%)	27 (7%)	13	8
1	B	360/400 (90%)	334 (93%)	26 (7%)	14	9
All	All	724/800 (90%)	671 (93%)	53 (7%)	14	8

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	45	LYS
1	B	176	VAL
1	B	409	VAL
1	B	65	GLN
1	B	106	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	12	GLN
1	B	477	HIS
1	B	161	ASN
1	B	414	ASN
1	B	65	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	B	3319	-	3,3,3	1.30	0	2,2,2	0.26	0
2	EDO	A	3318	-	3,3,3	0.90	0	2,2,2	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	3319	-	-	0/1/1/1	-
2	EDO	A	3318	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3318	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	482/517 (93%)	0.09	25 (5%) 27 33	17, 30, 63, 76	0
1	B	482/517 (93%)	0.40	40 (8%) 11 14	17, 37, 75, 87	0
All	All	964/1034 (93%)	0.25	65 (6%) 17 22	17, 34, 71, 87	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	367	VAL	8.2
1	B	369	GLY	5.7
1	B	344	LEU	5.4
1	B	368	ALA	5.3
1	B	293	SER	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	B	3319	4/4	0.74	0.25	37,42,43,44	0
2	EDO	A	3318	4/4	0.92	0.19	29,31,33,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.