



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 07:21 pm BST

PDB ID : 4R3P
Title : Crystal structures of EGFR in complex with Mig6
Authors : Park, E.; Kim, N.; Yi, Z.; Cho, A.; Kim, K.; Ficarro, S.B.; Park, A.; Park, W.Y.; Murray, B.; Meyerson, M.; Beroukim, R.; Marto, J.A.; Cho, J.; Eck, M.J.
Deposited on : 2014-08-17
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	
2	B	7	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	310	2437	1564	410	447	16	9	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	858	ARG	LEU	ENGINEERED MUTATION	UNP P00533

- Molecule 2 is a protein called peptide from ERBB receptor feedback inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	7	68	45	9	13	1	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		
3	B	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



- Molecule 2: peptide from ERBB receptor feedback inhibitor 1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	143.83Å 143.83Å 143.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.48 – 2.90 45.48 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (45.48-2.90) 92.3 (45.48-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.208 , 0.240 0.213 , 0.240	Depositor DCC
R_{free} test set	1097 reflections (9.96%)	wwPDB-VP
Wilson B-factor (Å ²)	76.8	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.032 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2559	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2488	0.48	0/3371
2	B	0.24	0/53	0.95	1/71 (1.4%)
All	All	0.25	0/2541	0.49	1/3442 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	397	LEU	CA-CB-CG	5.83	128.70	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2437	0	2434	48	0
2	B	68	0	58	1	0
3	A	51	0	0	1	1
3	B	3	0	0	0	0
All	All	2559	0	2492	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:832:ARG:NH2	1:A:863:GLY:O	2.17	0.78
1:A:711:GLU:HB3	1:A:731:TRP:HZ3	1.51	0.76
1:A:747:LEU:HD11	1:A:751:THR:HG21	1.67	0.76
1:A:776:ARG:HH21	1:A:1013:ALA:HB1	1.50	0.75
1:A:791:GLN:NE2	1:A:1014:ASP:OD1	2.22	0.70
1:A:783:THR:C	1:A:785:THR:H	1.93	0.69
1:A:884:GLU:OE2	1:A:958:ARG:NH1	2.28	0.67
2:B:397:LEU:HG	2:B:398:PRO:HD2	1.77	0.66
1:A:711:GLU:HB3	1:A:731:TRP:CZ3	2.30	0.66
1:A:783:THR:OG1	1:A:785:THR:O	2.13	0.65
1:A:783:THR:C	1:A:785:THR:N	2.54	0.61
1:A:879:LYS:NZ	1:A:915:TYR:O	2.32	0.61
1:A:795:PHE:HB2	1:A:845:VAL:HB	1.84	0.59
1:A:833:LEU:HD21	1:A:859:ALA:HB1	1.85	0.57
1:A:863:GLY:O	1:A:866:GLU:HG2	2.05	0.57
1:A:936:PRO:HG2	1:A:939:CYS:SG	2.45	0.56
1:A:736:GLU:OE2	1:A:1016:TYR:OH	2.15	0.54
1:A:834:VAL:HG12	1:A:836:ARG:HG3	1.88	0.54
1:A:836:ARG:NH1	1:A:858:ARG:O	2.31	0.54
1:A:783:THR:O	1:A:785:THR:N	2.41	0.53
1:A:805:HIS:O	1:A:809:ILE:HG13	2.08	0.53
1:A:961:PHE:O	1:A:965:ILE:HG13	2.09	0.53
1:A:818:CYS:O	1:A:900:TYR:OH	2.24	0.52
1:A:892:THR:O	1:A:895:SER:OG	2.23	0.49
1:A:778:LEU:HB2	1:A:789:ILE:HG22	1.95	0.49
1:A:846:LYS:HG2	1:A:847:THR:HG23	1.95	0.48
1:A:766:MET:HB3	1:A:777:LEU:HB2	1.97	0.46
1:A:832:ARG:NH2	1:A:862:LEU:HB2	2.31	0.46
1:A:936:PRO:HA	1:A:937:PRO:HD3	1.83	0.45
1:A:924:SER:O	1:A:928:GLU:HG3	2.17	0.44
1:A:737:LYS:N	3:A:1133:HOH:O	2.51	0.44
1:A:832:ARG:NH1	1:A:866:GLU:O	2.51	0.44
1:A:908:MET:HG3	1:A:947:MET:HE1	1.99	0.44
1:A:746:GLU:HA	1:A:787:GLN:HB3	1.99	0.44
1:A:747:LEU:HB3	1:A:785:THR:HB	1.99	0.43
1:A:892:THR:HG23	1:A:894:GLN:H	1.83	0.43
1:A:797:CYS:SG	1:A:841:ARG:HA	2.57	0.43
1:A:733:PRO:O	1:A:736:GLU:HB2	2.18	0.43
1:A:757:LYS:C	1:A:757:LYS:HD2	2.38	0.43
1:A:922:GLU:O	1:A:925:SER:OG	2.37	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:965:ILE:O	1:A:969:SER:OG	2.22	0.43
1:A:832:ARG:NH2	1:A:862:LEU:O	2.52	0.42
1:A:776:ARG:NH2	1:A:1013:ALA:HB1	2.27	0.41
1:A:761:ASP:O	1:A:765:VAL:HG23	2.21	0.41
1:A:935:GLN:HA	1:A:936:PRO:HD2	1.97	0.41
1:A:1012:ASP:OD1	1:A:1013:ALA:N	2.54	0.40
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.55	0.40
1:A:766:MET:HG3	1:A:856:PHE:CD2	2.57	0.40
1:A:889:ARG:HH11	1:A:889:ARG:HG3	1.84	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1103:HOH:O	3:A:1109:HOH:O[2_455]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/323 (94%)	295 (97%)	9 (3%)	0	100	100
2	B	4/7 (57%)	4 (100%)	0	0	100	100
All	All	308/330 (93%)	299 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/284 (92%)	256 (98%)	5 (2%)	57	84
2	B	6/6 (100%)	6 (100%)	0	100	100
All	All	267/290 (92%)	262 (98%)	5 (2%)	57	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	734	GLU
1	A	757	LYS
1	A	761	ASP
1	A	884	GLU
1	A	1006	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	771	ASN
1	A	812	GLN
2	B	393	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PTR	B	395	2	15,16,17	1.24	1 (6%)	19,22,24	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	B	395	2	-	0/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	395	PTR	OH-CZ	-4.33	1.30	1.40

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	310/323 (95%)	0.45	23 (7%) 14 11	46, 72, 132, 156	8 (2%)
2	B	6/7 (85%)	1.72	3 (50%) 0 0	97, 114, 132, 145	0
All	All	316/330 (95%)	0.47	26 (8%) 11 9	46, 72, 136, 156	8 (2%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	746	GLU	5.0
1	A	863	GLY	3.9
1	A	1018	ILE	3.9
1	A	722	ALA	3.8
1	A	747	LEU	3.8
2	B	398	PRO	3.8
1	A	720	SER	3.7
1	A	749	GLU	3.6
1	A	750	ALA	3.5
1	A	727	TYR	3.3
1	A	786	VAL	3.1
1	A	721	GLY	2.6
1	A	867	LYS	2.6
2	B	392	THR	2.5
1	A	719	GLY	2.5
1	A	725	THR	2.5
1	A	730	LEU	2.5
2	B	397	LEU	2.5
1	A	788	LEU	2.5
1	A	744	ILE	2.4
1	A	990	PRO	2.4
1	A	988	HIS	2.4
1	A	869	TYR	2.3
1	A	927	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	758	GLU	2.1
1	A	756	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PTR	B	395	16/17	0.87	0.30	102,109,122,122	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.