



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 11:07 PM EST

PDB ID : 1R53  
Title : Crystal structure of the bifunctional chorismate synthase from *Saccharomyces cerevisiae*  
Authors : Quevillon-Cheruel, S.; Leulliot, N.; Meyer, P.; Graille, M.; Bremang, M.; Blondeau, K.; Sorel, I.; Poupon, A.; Janin, J.; van Tilbeurgh, H.  
Deposited on : 2003-10-09  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chorismate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	287	2191	1376	384	416	15	65	5	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	377	HIS	-	expression tag	UNP P28777
A	378	HIS	-	expression tag	UNP P28777
A	379	HIS	-	expression tag	UNP P28777
A	380	HIS	-	expression tag	UNP P28777
A	381	HIS	-	expression tag	UNP P28777
A	382	HIS	-	expression tag	UNP P28777


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	113	Total	O	0	0
			113	113		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chorismate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.52Å 74.66Å 155.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.32 – 2.20 27.29 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.5 (27.32-2.20) 97.5 (27.29-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.60 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.174 , 0.225 0.180 , 0.223	Depositor DCC
$R_{free}$ test set	852 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.8	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2304	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.99	7/2247 (0.3%)	1.03	15/3028 (0.5%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	353	MET	SD-CE	-12.97	1.05	1.77
1	A	81	MET	SD-CE	-11.23	1.15	1.77
1	A	163	MET	SD-CE	-10.83	1.17	1.77
1	A	246	MET	SD-CE	-7.82	1.34	1.77
1	A	16	SER	CB-OG	5.47	1.49	1.42
1	A	150	GLU	CD-OE2	5.40	1.31	1.25
1	A	232	GLU	CD-OE2	5.29	1.31	1.25

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	ARG	NE-CZ-NH2	-12.72	113.94	120.30
1	A	224	ARG	NE-CZ-NH1	9.22	124.91	120.30
1	A	247	LEU	CA-CB-CG	-8.01	96.88	115.30
1	A	285	ARG	NE-CZ-NH2	7.42	124.01	120.30
1	A	127	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	A	47	ARG	NE-CZ-NH2	7.33	123.97	120.30
1	A	62	ARG	NE-CZ-NH2	7.32	123.96	120.30
1	A	337	ARG	NE-CZ-NH2	7.31	123.96	120.30
1	A	236	ASP	CB-CG-OD2	6.08	123.77	118.30
1	A	1	MET	CG-SD-CE	5.73	109.36	100.20
1	A	9	ARG	NE-CZ-NH2	5.62	123.11	120.30
1	A	38	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	26	ASP	CB-CG-OD1	5.32	123.08	118.30
1	A	276	TYR	CB-CG-CD1	-5.24	117.86	121.00
1	A	169	ASP	CB-CG-OD2	5.23	123.01	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2191	0	2208	49	2
2	A	113	0	0	3	0
All	All	2304	0	2208	49	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:MET:SD	1:A:163:MET:CE	1.17	1.26
1:A:81:MET:CE	1:A:81:MET:SD	1.14	1.24
1:A:353:MET:CE	1:A:353:MET:CG	2.19	1.19
1:A:163:MET:SD	1:A:163:MET:HE2	1.75	1.16
1:A:81:MET:SD	1:A:81:MET:HE2	1.73	1.15
1:A:81:MET:SD	1:A:81:MET:HE1	1.73	1.13
1:A:81:MET:SD	1:A:81:MET:HE3	1.73	1.13
1:A:353:MET:CE	1:A:353:MET:SD	1.05	1.13
1:A:163:MET:SD	1:A:163:MET:HE1	1.75	1.10
1:A:163:MET:SD	1:A:163:MET:HE3	1.75	1.08
1:A:81:MET:CE	1:A:81:MET:CG	2.34	1.05
1:A:353:MET:SD	1:A:353:MET:HE3	1.65	1.05
1:A:353:MET:SD	1:A:353:MET:HE2	1.65	1.02
1:A:353:MET:SD	1:A:353:MET:HE1	1.65	1.02
1:A:163:MET:CE	1:A:163:MET:CG	2.37	1.01
1:A:42[B]:GLN:HE21	1:A:42[B]:GLN:H	1.34	0.74
1:A:39:ILE:HD13	1:A:81:MET:CE	2.30	0.62
1:A:365:GLN:HE22	1:A:366:LYS:HE2	1.65	0.61
1:A:143:LEU:O	1:A:147:SER:OG	2.14	0.60
1:A:353:MET:CE	1:A:353:MET:HG2	2.23	0.60
1:A:312:SER:HB2	2:A:485:HOH:O	2.02	0.59
1:A:353:MET:CG	1:A:353:MET:HE3	2.10	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:LEU:CD1	1:A:63:VAL:HG21	2.35	0.56
1:A:40:GLN:NE2	1:A:62:ARG:O	2.31	0.56
1:A:82:MET:SD	2:A:462:HOH:O	2.59	0.53
1:A:339:ASP:CB	1:A:340:PRO:HA	2.39	0.53
1:A:214:SER:OG	1:A:338:HIS:HD2	1.92	0.52
1:A:214:SER:HB2	1:A:312:SER:OG	2.10	0.52
1:A:63:VAL:HG22	1:A:83:ILE:HD11	1.93	0.50
1:A:353:MET:HE3	1:A:353:MET:HG2	1.89	0.50
1:A:43:LEU:HD11	1:A:63:VAL:HG21	1.94	0.49
1:A:81:MET:HE3	1:A:81:MET:CG	2.25	0.49
1:A:39:ILE:CD1	1:A:81:MET:CE	2.92	0.47
1:A:83:ILE:HD13	1:A:130:ILE:CG2	2.45	0.47
1:A:339:ASP:HB2	1:A:340:PRO:HA	1.96	0.46
1:A:62:ARG:O	1:A:63:VAL:HB	2.16	0.45
1:A:221:CYS:O	1:A:305:TYR:HA	2.16	0.45
1:A:36:GLU:HG2	1:A:63:VAL:O	2.17	0.45
1:A:42[A]:GLN:OE1	1:A:45:ARG:HD3	2.18	0.43
1:A:48:PRO:HB2	1:A:189:PRO:HB3	2.00	0.43
1:A:83:ILE:HD13	1:A:130:ILE:HG21	2.01	0.42
1:A:365:GLN:NE2	1:A:366:LYS:HE2	2.33	0.42
1:A:62:ARG:O	1:A:63:VAL:CB	2.68	0.41
1:A:77:THR:HB	1:A:78:PRO:CD	2.50	0.41
1:A:81:MET:CE	1:A:81:MET:HG2	2.40	0.41
1:A:292:GLY:O	1:A:304:ILE:HA	2.21	0.41
1:A:14:GLY:HA3	1:A:21:VAL:HG12	2.02	0.41
1:A:158:ILE:HG21	1:A:205:ILE:HG13	2.02	0.40
1:A:140:GLU:OE1	2:A:495:HOH:O	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:CG	1:A:9:ARG:NH1[3_555]	1.48	0.72
1:A:127:ARG:NH1	1:A:233:PRO:CB[2_565]	2.04	0.16



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	282/382 (74%)	273 (97%)	6 (2%)	3 (1%)	14 12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	ASP
1	A	63	VAL
1	A	147	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	243/318 (76%)	238 (98%)	5 (2%)	53 67

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	ASP
1	A	82	MET
1	A	247	LEU
1	A	311	LYS
1	A	312	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	GLN
1	A	146	ASN
1	A	244	HIS
1	A	295	GLN
1	A	338	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	287/382 (75%)	-0.06	17 (5%) <span style="border: 1px solid red; padding: 2px;">22</span> <span style="border: 1px solid red; padding: 2px;">21</span>	14, 27, 54, 72	14 (4%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	ASP	6.2
1	A	17[A]	HIS	6.0
1	A	47	ARG	5.3
1	A	48	PRO	4.7
1	A	71	PHE	4.6
1	A	85	ASN	4.6
1	A	370	PHE	4.3
1	A	285	ARG	3.7
1	A	127	ARG	3.7
1	A	286	THR	2.8
1	A	16	SER	2.8
1	A	18	CYS	2.6
1	A	148	ASN	2.5
1	A	339	ASP	2.4
1	A	313	VAL	2.2
1	A	275	PHE	2.2
1	A	19	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.