



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 07:07 PM EST

PDB ID : 3R63
Title : Structure of ERK2 (SPE) mutant (S246E)
Authors : Livnah, O.; Karamansha, Y.
Deposited on : 2011-03-21
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	347	2834	1821	485	515	13	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	246	GLU	SER	engineered mutation	UNP P63086


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	161	161	161	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.77Å 70.00Å 60.12Å 90.00° 109.04° 90.00°	Depositor
Resolution (Å)	50.00 – 1.70 43.39 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.9 (50.00-1.70) 96.9 (43.39-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.188 , 0.245 0.188 , 0.245	Depositor DCC
R_{free} test set	2059 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.196	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2995	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/2902	0.95	10/3931 (0.3%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	ARG	NE-CZ-NH1	18.63	129.62	120.30
1	A	89	ARG	NE-CZ-NH2	-13.76	113.42	120.30
1	A	146	ARG	NE-CZ-NH2	-8.70	115.95	120.30
1	A	89	ARG	CD-NE-CZ	7.97	134.76	123.60
1	A	146	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	A	170	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	74	LEU	CB-CG-CD1	5.86	120.96	111.00
1	A	351	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	223	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	113	LEU	CA-CB-CG	5.20	127.25	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2834	0	2831	48	0
2	A	161	0	0	1	0
All	All	2995	0	2831	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:LEU:HD11	1:A:71:LYS:HD2	1.52	0.91
1:A:51:ILE:HD12	1:A:102:VAL:HG22	1.56	0.87
1:A:78:HIS:HD2	1:A:80:ASN:H	1.23	0.87
1:A:89:ARG:HD3	1:A:349:THR:OG1	1.76	0.85
1:A:78:HIS:CD2	1:A:80:ASN:H	2.00	0.80
1:A:67:LEU:CD1	1:A:71:LYS:HD2	2.16	0.75
1:A:22:ARG:HG2	1:A:23:TYR:CZ	2.23	0.74
1:A:170:ARG:HH11	1:A:170:ARG:HG2	1.54	0.72
1:A:107:GLU:HG3	1:A:157:THR:HG23	1.76	0.67
1:A:170:ARG:HG2	1:A:170:ARG:NH1	2.09	0.67
1:A:170:ARG:HH11	1:A:170:ARG:CG	2.11	0.64
1:A:107:GLU:HG3	1:A:157:THR:CG2	2.29	0.63
1:A:237:HIS:HD2	2:A:454:HOH:O	1.81	0.63
1:A:64:GLN:NE2	1:A:333:LEU:O	2.37	0.57
1:A:69:GLU:HG3	1:A:73:LEU:HD22	1.87	0.56
1:A:107:GLU:HG2	1:A:156:ASN:HA	1.90	0.54
1:A:22:ARG:HG2	1:A:23:TYR:CE2	2.43	0.54
1:A:68:ARG:HD3	1:A:169:ALA:O	2.08	0.53
1:A:146:ARG:HD2	1:A:202:GLY:O	2.10	0.52
1:A:241:ILE:HD12	1:A:276:LEU:HD11	1.92	0.52
1:A:42:ASP:O	1:A:46:LYS:N	2.38	0.51
1:A:356:TYR:C	1:A:356:TYR:CD1	2.85	0.51
1:A:27:SER:HB3	1:A:39:SER:HB3	1.94	0.49
1:A:75:ARG:NH2	1:A:327:PHE:HA	2.27	0.48
1:A:51:ILE:CD1	1:A:102:VAL:HG13	2.43	0.48
1:A:343:GLU:O	1:A:347:GLU:HG2	2.13	0.48
1:A:113:LEU:HD22	1:A:117:GLN:HB2	1.97	0.47
1:A:199:ASN:C	1:A:199:ASN:HD22	2.19	0.47
1:A:107:GLU:N	1:A:107:GLU:CD	2.69	0.46
1:A:78:HIS:HD2	1:A:80:ASN:N	2.03	0.45
1:A:356:TYR:C	1:A:356:TYR:HD1	2.20	0.45
1:A:9:PRO:HG3	1:A:15:GLN:HA	1.98	0.45
1:A:61:THR:O	1:A:65:ARG:HD2	2.17	0.44
1:A:124:ILE:HD13	1:A:220:LEU:HD23	1.98	0.44
1:A:107:GLU:CG	1:A:157:THR:HG23	2.46	0.44
1:A:255:ASN:C	1:A:255:ASN:HD22	2.20	0.44
1:A:107:GLU:CD	1:A:107:GLU:H	2.22	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:ARG:HH21	1:A:68:ARG:NE	2.17	0.43
1:A:239:LEU:HB3	1:A:268:LYS:HE3	2.01	0.42
1:A:161:LEU:C	1:A:161:LEU:HD23	2.39	0.42
1:A:53:LYS:HE3	1:A:100:TYR:OH	2.19	0.42
1:A:196:ILE:HD11	1:A:207:ILE:HD11	2.02	0.42
1:A:152:ASN:HD22	1:A:152:ASN:HA	1.66	0.42
1:A:333:LEU:HD23	1:A:334:ASP:OD1	2.21	0.41
1:A:64:GLN:HE22	1:A:333:LEU:C	2.24	0.41
1:A:255:ASN:ND2	1:A:258:ALA:H	2.19	0.41
1:A:22:ARG:O	1:A:22:ARG:HG3	2.21	0.40
1:A:35:GLY:HA3	1:A:53:LYS:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	342/358 (96%)	326 (95%)	14 (4%)	2 (1%)	25 11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	GLY
1	A	255	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	311/318 (98%)	287 (92%)	24 (8%)	13 3

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ARG
1	A	24	THR
1	A	60	GLN
1	A	64	GLN
1	A	65	ARG
1	A	71	LYS
1	A	73	LEU
1	A	74	LEU
1	A	75	ARG
1	A	89	ARG
1	A	107	GLU
1	A	113	LEU
1	A	155	LEU
1	A	170	ARG
1	A	199	ASN
1	A	223	ARG
1	A	250	LEU
1	A	255	ASN
1	A	256	LEU
1	A	274	ASN
1	A	328	LYS
1	A	340	LYS
1	A	344	LEU
1	A	356	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	60	GLN
1	A	64	GLN
1	A	78	HIS
1	A	85	ASN
1	A	121	ASN
1	A	152	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	178	HIS
1	A	199	ASN
1	A	237	HIS
1	A	251	ASN
1	A	255	ASN
1	A	260	ASN
1	A	274	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/358 (96%)	0.44	32 (9%) 9 10	17, 32, 69, 83	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	356	TYR	13.2
1	A	355	GLY	8.7
1	A	34	TYR	7.7
1	A	357	ARG	5.8
1	A	358	SER	5.3
1	A	33	ALA	5.3
1	A	59	HIS	4.8
1	A	41	TYR	4.7
1	A	20	GLY	4.2
1	A	335	ASP	4.0
1	A	9	PRO	4.0
1	A	62	TYR	4.0
1	A	329	PHE	3.9
1	A	21	PRO	3.6
1	A	32	GLY	3.6
1	A	334	ASP	3.6
1	A	19	VAL	3.5
1	A	26	LEU	3.1
1	A	15	GLN	3.1
1	A	47	VAL	3.0
1	A	27	SER	3.0
1	A	43	ASN	2.9
1	A	336	LEU	2.6
1	A	4	ALA	2.5
1	A	5	ALA	2.5
1	A	333	LEU	2.5
1	A	46	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	38	CYS	2.3
1	A	44	LEU	2.2
1	A	65	ARG	2.2
1	A	28	TYR	2.2
1	A	35	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.