



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 09:05 AM BST

PDB ID : 6R7O  
Title : Crystal structure of the central region of human cohesin subunit STAG1  
Authors : Newman, J.A.; katis, V.L.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.;  
Bountra, C.; Gileadi, O.  
Deposited on : 2019-03-29  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	459	 2% 88% 6% 6%
1	B	459	 5% 81% 13% 6%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cohesin subunit SA-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	3458	2204	574	658	22	0	0	0
1	B	430	3446	2195	577	652	22	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	457	SER	-	expression tag	UNP Q8WVM7
A	458	MET	-	expression tag	UNP Q8WVM7
B	457	SER	-	expression tag	UNP Q8WVM7
B	458	MET	-	expression tag	UNP Q8WVM7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	135	Total	O	0	0
			135	135		
2	B	49	Total	O	0	0
			49	49		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cohesin subunit SA-1



- Molecule 1: Cohesin subunit SA-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	154.65Å 166.72Å 46.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	113.36 – 2.31 113.38 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.6 (113.36-2.31) 99.6 (113.38-2.31)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 2.32Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.226 , 0.264 0.226 , 0.265	Depositor DCC
$R_{free}$ test set	2728 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.1	Xtrriage
Anisotropy	0.370	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7088	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/3520	0.35	0/4762
1	B	0.23	0/3506	0.36	0/4737
All	All	0.23	0/7026	0.36	0/9499

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3425	15	0
1	B	3446	0	3427	32	0
2	A	135	0	0	2	0
2	B	49	0	0	0	0
All	All	7088	0	6852	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:523:SER:OG	1:B:588:LYS:NZ	2.23	0.71

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:613:LEU:HD23	1:B:653:ARG:HD2	1.74	0.70
1:B:589:VAL:HG11	1:B:623:VAL:HG11	1.76	0.68
1:A:601:LEU:HB2	1:A:646:GLU:HG3	1.76	0.67
1:B:550:ARG:NH1	1:B:556:GLU:OE2	2.33	0.61
1:B:752:ILE:HG23	1:B:757:PRO:HG3	1.82	0.61
1:B:676:LEU:HD22	1:B:691:VAL:HG21	1.82	0.61
1:A:855:GLU:HG2	1:A:858:LYS:HE3	1.82	0.60
1:A:493:GLN:HG2	1:A:497:LYS:HE2	1.84	0.58
1:A:648:TYR:HA	1:A:651:GLN:HB3	1.86	0.58
1:B:770:LYS:HG2	1:B:819:PHE:HE1	1.70	0.57
1:A:667:ASP:O	1:A:671:HIS:ND1	2.35	0.56
1:A:766:ARG:NH1	1:A:816:PRO:O	2.41	0.52
1:B:839:ILE:HG22	1:B:840:ASP:H	1.74	0.52
1:B:862:LEU:HD11	1:B:899:ASP:HB3	1.92	0.52
1:B:493:GLN:HG2	1:B:497:LYS:HE2	1.90	0.51
1:B:830:LEU:HD11	1:B:886:ALA:HA	1.92	0.50
1:B:473:LEU:HD13	1:B:524:ALA:HB2	1.95	0.49
1:A:668:ARG:NH2	2:A:1007:HOH:O	2.46	0.48
1:A:489:TRP:CD2	1:A:535:GLN:HG2	2.49	0.48
1:B:600:ASP:HB3	1:B:603:ILE:HD13	1.97	0.47
1:B:753:THR:O	1:B:811:ARG:NH2	2.47	0.47
1:B:752:ILE:HD13	1:B:762:LEU:HD23	1.95	0.47
1:B:826:GLN:HG2	1:B:881:VAL:HG12	1.97	0.47
1:A:800:MET:HE2	1:A:875:LEU:HD23	1.95	0.47
1:B:866:ARG:NH1	1:B:899:ASP:O	2.47	0.46
1:B:721:LYS:NZ	1:B:725:GLU:OE2	2.49	0.46
1:B:489:TRP:CD2	1:B:535:GLN:HG2	2.51	0.45
1:B:770:LYS:HG2	1:B:819:PHE:CE1	2.51	0.45
1:A:859:ILE:H	1:A:859:ILE:HG13	1.57	0.45
1:B:628:VAL:O	1:B:696:LYS:HE3	2.17	0.45
1:B:788:LYS:HB3	1:B:832:PHE:HE1	1.81	0.44
1:B:800:MET:HG2	1:B:874:LYS:HE2	1.98	0.44
1:B:670:ASN:OD1	1:B:715:ASN:ND2	2.51	0.44
1:A:489:TRP:CE2	1:A:535:GLN:HG2	2.53	0.43
1:B:766:ARG:O	1:B:770:LYS:HG3	2.19	0.42
1:A:873:SER:HA	1:A:876:ILE:HD12	2.01	0.42
1:B:489:TRP:O	1:B:493:GLN:HB2	2.20	0.42
1:B:882:ASP:OD2	1:B:885:ALA:N	2.47	0.42
1:A:493:GLN:NE2	2:A:1012:HOH:O	2.52	0.41
1:B:839:ILE:O	1:B:896:TYR:OH	2.31	0.41
1:B:800:MET:HE1	1:B:880:ILE:HD12	2.02	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:815:GLN:N	1:B:816:PRO:HD2	2.36	0.41
1:B:666:VAL:HG21	1:B:710:TRP:CD1	2.56	0.41
1:B:461:ASN:HA	1:B:464:LEU:HB2	2.02	0.41
1:A:633:LEU:HD13	1:A:697:ARG:HA	2.03	0.40
1:A:624:VAL:O	1:A:697:ARG:NE	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/459 (93%)	415 (98%)	10 (2%)	0	100	100
1	B	420/459 (92%)	408 (97%)	12 (3%)	0	100	100
All	All	845/918 (92%)	823 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/415 (93%)	379 (98%)	6 (2%)	62	77
1	B	384/415 (92%)	378 (98%)	6 (2%)	62	77
All	All	769/830 (93%)	757 (98%)	12 (2%)	62	77



All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	641	SER
1	A	646	GLU
1	A	647	GLU
1	A	650	ILE
1	A	795	LEU
1	A	797	ASP
1	B	649	THR
1	B	654	VAL
1	B	839	ILE
1	B	855	GLU
1	B	884	HIS
1	B	888	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	820	ASN
1	B	493	GLN
1	B	748	GLN
1	B	820	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	433/459 (94%)	0.34	7 (1%) 72 78	32, 56, 89, 104	0
1	B	430/459 (93%)	0.56	24 (5%) 24 31	43, 69, 103, 124	0
All	All	863/918 (94%)	0.45	31 (3%) 42 49	32, 62, 98, 124	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	892	HIS	4.8
1	B	887	ALA	4.4
1	B	550	ARG	4.4
1	B	680	GLY	4.4
1	B	460	PRO	4.1
1	B	907	THR	3.8
1	B	872	PHE	3.3
1	B	677	LEU	3.3
1	B	869	LEU	3.3
1	B	904	ILE	3.1
1	B	908	LEU	3.1
1	B	894	MET	3.0
1	A	819	PHE	3.0
1	A	471	PHE	2.7
1	B	877	ILE	2.7
1	B	839	ILE	2.7
1	A	606	THR	2.5
1	B	804	HIS	2.5
1	B	678	GLN	2.4
1	B	551	VAL	2.3
1	B	889	ILE	2.3
1	A	515	GLU	2.2
1	A	551	VAL	2.1
1	B	884	HIS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	886	ALA	2.1
1	A	470	LEU	2.1
1	B	876	ILE	2.1
1	A	895	LYS	2.1
1	B	893	TYR	2.1
1	B	473	LEU	2.0
1	B	665	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.