



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 07:39 AM EDT

PDB ID : 2R8U
Title : Structure of fragment of human end-binding protein 1 (EB1) containing the N-terminal domain at 1.35 Å resolution
Authors : Huang, X.; Lovelace, L.L.; Smith, D.; Lebioda, L.
Deposited on : 2007-09-11
Resolution : 1.35 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	268	
1	B	268	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Microtubule-associated protein RP/EB family member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	131	1067	692	176	192	7	0	0	0
1	B	136	1095	706	182	200	7	0	0	0

- Molecule 2 is MOLYBDATE ION (three-letter code: MOO) (formula: MoO₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Mo	O		
2	A	1	5	1	4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	142	142	142	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	118	Total 118	O 118	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.89Å 47.92Å 100.66Å 90.00° 91.65° 90.00°	Depositor
Resolution (Å)	10.00 – 1.35 31.88 – 1.27	Depositor EDS
% Data completeness (in resolution range)	76.0 (10.00-1.35) 63.1 (31.88-1.27)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.53 (at 1.27Å)	Xtrriage
Refinement program	SHELX, SHELXL-97	Depositor
R, R_{free}	0.164 , 0.227 0.225 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtrriage
Anisotropy	0.468	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.097 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2427	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MOO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/1091	1.16	4/1466 (0.3%)
1	B	0.48	0/1118	1.07	3/1500 (0.2%)
All	All	0.51	0/2209	1.11	7/2966 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	13	ASP	CB-CG-OD1	5.68	123.42	118.30
1	B	44	TYR	CB-CG-CD1	-5.64	117.62	121.00
1	A	71	TYR	CB-CG-CD1	5.61	124.36	121.00
1	A	110	TRP	CG-CD2-CE3	-5.55	128.91	133.90
1	B	88	ASP	CB-CG-OD1	-5.31	113.52	118.30
1	B	55	SER	O-C-N	5.15	130.94	122.70
1	A	6	TYR	CB-CG-CD2	-5.03	117.98	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1067	0	1064	23	0
1	B	1095	0	1082	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	5	0	0	0	0
3	A	142	0	0	6	0
3	B	118	0	0	7	0
All	All	2427	0	2146	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:VAL:H	1:A:118:ASN:HD21	1.10	0.90
1:B:3:VAL:H	1:B:118:ASN:HD21	1.13	0.87
1:B:77:ILE:HD13	3:B:366:HOH:O	1.89	0.72
1:A:98:LYS:HD2	3:A:392:HOH:O	1.90	0.72
1:B:15:LEU:HD11	3:B:355:HOH:O	1.91	0.69
1:A:24:ILE:HD12	1:A:38:LEU:HD21	1.76	0.66
1:A:82:PHE:CE1	1:A:91:ILE:HD11	2.31	0.65
1:A:84:ARG:HD3	3:A:351:HOH:O	2.00	0.62
1:A:3:VAL:N	1:A:118:ASN:HD21	1.92	0.62
1:A:91:ILE:HD12	1:A:107:PHE:CE1	2.35	0.61
1:A:82:PHE:HE1	1:A:91:ILE:HD11	1.65	0.60
1:B:3:VAL:N	1:B:118:ASN:HD21	1.94	0.59
1:B:24:ILE:HD13	1:B:47:PHE:HE1	1.69	0.57
1:B:62:LYS:HE2	1:B:73:GLN:NE2	2.20	0.56
1:A:3:VAL:H	1:A:118:ASN:ND2	1.93	0.55
1:A:59:LYS:HG3	3:A:374:HOH:O	2.08	0.54
1:A:59:LYS:HG3	1:A:59:LYS:O	2.08	0.53
1:B:62:LYS:HG3	3:B:319:HOH:O	2.08	0.53
1:A:62:LYS:HA	3:A:336:HOH:O	2.10	0.52
1:A:91:ILE:HD13	1:A:91:ILE:N	2.25	0.50
1:A:77:ILE:HD13	1:A:77:ILE:N	2.31	0.46
1:A:17:ARG:HD3	3:A:288:HOH:O	2.15	0.45
1:B:188:ASN:HB3	3:B:346:HOH:O	2.16	0.45
1:A:91:ILE:HD12	1:A:107:PHE:HE1	1.82	0.44
1:A:84:ARG:NH2	3:A:343:HOH:O	2.50	0.44
1:A:88:ASP:OD1	1:A:88:ASP:O	2.35	0.44
1:A:24:ILE:HD13	1:A:47:PHE:HE1	1.82	0.44
1:B:33:THR:HB	3:B:340:HOH:O	2.19	0.43
1:B:24:ILE:HD13	1:B:47:PHE:CE1	2.53	0.42
1:B:70:GLU:HA	1:B:73:GLN:HE21	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:GLU:HG3	3:B:311:HOH:O	2.19	0.42
1:B:56:ILE:HD12	1:B:56:ILE:HA	1.87	0.42
1:B:59:LYS:O	1:B:59:LYS:HD2	2.20	0.42
1:B:23:TRP:HA	3:B:342:HOH:O	2.20	0.42
1:A:14:ASN:H	1:B:188:ASN:ND2	2.18	0.41
1:A:66:LYS:HZ3	1:A:66:LYS:HG3	1.67	0.41
1:A:70:GLU:HA	1:A:73:GLN:HE21	1.86	0.41
1:A:62:LYS:HE2	1:A:73:GLN:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/268 (48%)	128 (99%)	1 (1%)	0	100	100
1	B	132/268 (49%)	131 (99%)	1 (1%)	0	100	100
All	All	261/536 (49%)	259 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/228 (50%)	112 (97%)	3 (3%)	46	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	117/228 (51%)	112 (96%)	5 (4%)	29	3
All	All	232/456 (51%)	224 (97%)	8 (3%)	37	7

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	88	ASP
1	A	102	GLN
1	B	31	ASN
1	B	56	ILE
1	B	58	LEU
1	B	59	LYS
1	B	106	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	64	GLN
1	A	73	GLN
1	A	118	ASN
1	B	73	GLN
1	B	118	ASN
1	B	131	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MOO	A	269	-	2,4,4	13.89	2 (100%)	-		

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	269	MOO	O1-MO	14.00	2.03	1.73
2	A	269	MOO	O2-MO	13.78	2.02	1.73

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.